

Table 13. Proportion of persons surviving for more than 12, 24, and 36 months after an AIDS diagnosis in 2001, by selected characteristics—United States and dependent areas

	No. of persons	Survival in months		
		>12	>24	>36
Data for 50 states and the District of Columbia				
Age at diagnosis (yrs)				
<13	110	0.95	0.95	0.95
13–14	77	1.00	1.00	0.99
15–19	259	0.97	0.94	0.92
20–24	1,255	0.96	0.93	0.90
25–29	3,038	0.95	0.92	0.90
30–34	5,878	0.94	0.91	0.88
35–39	8,095	0.93	0.90	0.87
40–44	6,922	0.91	0.87	0.84
45–49	4,969	0.89	0.84	0.81
50–54	2,895	0.87	0.83	0.79
55–59	1,395	0.83	0.77	0.74
60–64	720	0.79	0.73	0.70
≥65	648	0.70	0.63	0.59
Race/ethnicity				
White, not Hispanic	10,482	0.92	0.88	0.86
Black, not Hispanic	18,543	0.90	0.86	0.82
Hispanic	6,512	0.93	0.90	0.88
Asian/Pacific Islander	355	0.90	0.88	0.87
American Indian/Alaska Native	150	0.86	0.82	0.77
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	12,468	0.93	0.90	0.88
Injection drug use	4,384	0.88	0.83	0.79
Male-to-male sexual contact and injection drug use	1,738	0.93	0.89	0.84
High-risk heterosexual contact ^a	2,985	0.91	0.88	0.86
Other ^b	4,977	0.87	0.83	0.81
Subtotal	26,552	0.91	0.87	0.85
Female adult or adolescent				
Injection drug use	2,254	0.90	0.84	0.78
High-risk heterosexual contact ^a	4,587	0.93	0.89	0.86
Other ^b	2,758	0.90	0.86	0.83
Subtotal	9,599	0.91	0.87	0.83
Child (<13 yrs at diagnosis)				
Perinatal	96	0.95	0.95	0.95
Other ^c	14	1.00	1.00	1.00
Subtotal	110	0.95	0.95	0.95
Subtotal for 50 states and the District of Columbia	36,261	0.91	0.87	0.84
Data for U.S. dependent areas	1,013	0.82	0.78	0.75
Total	37,325^d	0.91	0.87	0.84

Note. Excludes persons whose date of death was before, or in the same month as, date of diagnosis. Excludes persons whose month of diagnosis or month of death is unknown.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^c Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^d Includes 221 persons of unknown race or multiple races.