

**Table 3. Estimated numbers of AIDS cases, by year of diagnosis and selected characteristics, 2001–2005 and cumulative—United States and dependent areas (Revised June 2007)**

	Year of diagnosis					Cumulative <sup>a</sup>
	2001	2002	2003	2004	2005	
<b>Data for 50 states and the District of Columbia</b>						
<b>Age at diagnosis (yrs)</b>						
<13	121	105	71	50	68	9,112
13–14	82	67	78	83	86	1,062
15–19	266	318	309	341	447	5,289
20–24	1,305	1,404	1,595	1,703	1,836	34,795
25–29	3,143	3,149	3,121	3,327	3,407	114,141
30–34	6,101	5,737	5,663	5,355	5,122	193,926
35–39	8,419	8,288	8,294	7,448	7,246	208,505
40–44	7,287	7,472	7,839	8,080	8,210	164,697
45–49	5,257	5,406	5,804	5,763	6,418	102,732
50–54	3,096	3,293	3,470	3,674	3,935	56,950
55–59	1,492	1,603	1,737	1,914	2,064	30,424
60–64	790	877	889	939	967	16,493
≥65	720	689	795	845	801	14,503
<b>Race/ethnicity</b>						
White, not Hispanic	11,001	11,274	11,254	11,520	11,780	385,537
Black, not Hispanic	19,465	19,559	20,224	19,983	20,187	397,548
Hispanic	6,847	6,711	7,291	7,099	7,676	155,179
Asian/Pacific Islander	374	427	461	471	483	7,659
American Indian/Alaska Native	166	181	186	188	182	3,238
<b>Transmission category</b>						
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	15,294	15,764	16,448	16,660	17,230	452,111
Injection drug use	5,948	5,682	5,579	5,243	5,441	168,314
Male-to-male sexual contact and injection drug use	2,104	1,990	2,002	1,942	2,018	65,881
High-risk heterosexual contact <sup>b</sup>	4,293	4,567	4,616	4,707	4,797	61,438
Other <sup>c</sup>	269	272	246	265	280	13,978
Subtotal	27,908	28,276	28,891	28,817	29,766	761,723
<b>Female adult or adolescent</b>						
Injection drug use	3,099	2,911	2,971	2,961	2,940	73,050
High-risk heterosexual contact <sup>b</sup>	6,730	6,895	7,501	7,447	7,591	102,171
Other <sup>c</sup>	220	221	232	248	243	6,582
Subtotal	10,049	10,027	10,704	10,656	10,774	181,802
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	118	103	71	50	67	8,460
Other <sup>d</sup>	3	2	1	0	1	641
Subtotal	121	105	71	50	68	9,101
<b>Region of residence</b>						
Northeast	11,273	10,292	10,955	10,452	11,529	300,963
Midwest	3,929	4,126	4,282	4,225	4,862	97,930
South	16,571	17,301	18,014	18,761	18,115	359,725
West	6,306	6,689	6,414	6,086	6,102	194,011
<b>Subtotal for 50 states and the District of Columbia</b>	<b>38,079</b>	<b>38,408</b>	<b>39,666</b>	<b>39,524</b>	<b>40,608</b>	<b>952,629</b>
<b>Data for U.S. dependent areas</b>	<b>1,189</b>	<b>1,046</b>	<b>1,060</b>	<b>923</b>	<b>982</b>	<b>30,386</b>
<b>Total<sup>e</sup></b>	<b>39,327</b>	<b>39,512</b>	<b>40,834</b>	<b>40,665</b>	<b>41,897</b>	<b>984,155</b>

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

<sup>a</sup> From the beginning of the epidemic through 2005.

<sup>b</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>c</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>d</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>e</sup> Includes persons of unknown race or multiple races and persons of unknown sex. Cumulative total includes 3,469 persons of unknown race or multiple races, 3 persons of unknown sex, 1,136 persons of unknown state of residence, and 3 persons who were residents of other areas. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.