

Table 35. Tuberculosis Cases and Percentages by Initial Drug Regimen: Reporting Areas, 2006

Reporting Area	Total Cases	Cases in Persons Alive at Diagnosis	Cases with Information on Initial Drug Regimen		Percentage of Cases in Persons with Initial Drug Regimen ^{1,2}			
			No.	(%)	IR	IRZ	IRZ,E/S	IRZE ³
United States	13,779	13,458	13,402	(99.6)	(1.2)	(4.8)	(83.0)	(82.9)
Alabama	196	192	192	(100.0)	(0.5)	(2.1)	(89.6)	(89.6)
Alaska	70	68	68	(100.0)	(1.5)	(10.3)	(80.9)	(80.9)
Arizona	315	308	306	(99.4)	(0.7)	(3.9)	(86.3)	(84.6)
Arkansas	102	99	99	(100.0)	(18.2)	(27.3)	(37.4)	(37.4)
California	2,779	2,714	2,712	(99.9)	(0.8)	(2.9)	(86.9)	(86.8)
Colorado	124	121	121	(100.0)	(0.8)	(3.3)	(79.3)	(79.3)
Connecticut	89	85	85	(100.0)	(1.2)	(7.1)	(82.4)	(82.4)
Delaware	29	28	28	(100.0)	(0.0)	(3.6)	(92.9)	(92.9)
District of Columbia	72	72	64	(88.9)	(1.6)	(3.1)	(93.8)	(93.8)
Florida	1,038	1,013	1,012	(99.9)	(0.2)	(5.5)	(85.1)	(85.1)
Georgia	504	489	488	(99.8)	(0.4)	(5.9)	(75.0)	(75.0)
Hawaii	115	115	115	(100.0)	(1.7)	(5.2)	(81.7)	(80.9)
Idaho	20	20	19	(95.0)	(5.3)	(10.5)	(78.9)	(78.9)
Illinois	569	562	552	(98.2)	(1.1)	(5.8)	(84.8)	(84.8)
Indiana	125	123	122	(99.2)	(0.0)	(9.0)	(84.4)	(84.4)
Iowa	40	39	39	(100.0)	(0.0)	(10.3)	(82.1)	(82.1)
Kansas	82	78	78	(100.0)	(0.0)	(3.8)	(84.6)	(84.6)
Kentucky	84	83	83	(100.0)	(0.0)	(2.4)	(88.0)	(88.0)
Louisiana	207	196	195	(99.5)	(3.1)	(7.7)	(87.2)	(87.2)
Maine	16	16	16	(100.0)	(0.0)	(18.8)	(81.3)	(81.3)
Maryland	253	244	244	(100.0)	(0.0)	(4.5)	(93.4)	(93.4)
Massachusetts	259	257	255	(99.2)	(0.4)	(0.8)	(85.1)	(85.1)
Michigan	221	218	218	(100.0)	(5.0)	(19.7)	(69.7)	(69.7)
Minnesota	217	213	213	(100.0)	(0.5)	(4.2)	(87.3)	(87.3)
Mississippi	115	112	112	(100.0)	(1.8)	(13.4)	(80.4)	(80.4)
Missouri	104	102	102	(100.0)	(2.0)	(15.7)	(78.4)	(78.4)
Montana	13	12	12	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
Nebraska	25	24	24	(100.0)	(0.0)	(20.8)	(79.2)	(79.2)
Nevada	101	100	100	(100.0)	(0.0)	(0.0)	(94.0)	(94.0)
New Hampshire	17	17	17	(100.0)	(0.0)	(5.9)	(88.2)	(88.2)
New Jersey	508	497	497	(100.0)	(1.0)	(4.4)	(82.7)	(82.7)
New Mexico	48	46	46	(100.0)	(0.0)	(10.9)	(82.6)	(82.6)
New York State ⁴	317	310	309	(99.7)	(1.0)	(5.5)	(84.8)	(84.8)
New York City	954	939	939	(100.0)	(1.6)	(1.8)	(86.5)	(86.5)
North Carolina	374	367	367	(100.0)	(0.5)	(1.9)	(90.7)	(90.2)
North Dakota	9	9	9	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
Ohio	239	234	234	(100.0)	(1.3)	(3.4)	(88.0)	(87.6)
Oklahoma	144	141	141	(100.0)	(12.8)	(12.8)	(63.1)	(63.1)
Oregon	81	78	78	(100.0)	(0.0)	(2.6)	(89.7)	(89.7)
Pennsylvania	341	331	331	(100.0)	(0.6)	(3.6)	(42.6)	(42.6)
Rhode Island	26	26	26	(100.0)	(7.7)	(3.8)	(76.9)	(76.9)
South Carolina	222	213	213	(100.0)	(0.9)	(3.8)	(81.2)	(81.2)
South Dakota	14	14	14	(100.0)	(0.0)	(14.3)	(71.4)	(71.4)
Tennessee	279	273	273	(100.0)	(1.5)	(5.5)	(61.9)	(61.9)
Texas	1,585	1,541	1,518	(98.5)	(1.1)	(4.2)	(84.4)	(84.4)
Utah	34	34	34	(100.0)	(0.0)	(5.9)	(88.2)	(88.2)
Vermont	8	7	6	(85.7)	(0.0)	(50.0)	(50.0)	(50.0)
Virginia	332	321	321	(100.0)	(0.9)	(5.3)	(83.5)	(83.5)
Washington	262	257	257	(100.0)	(1.2)	(4.7)	(86.8)	(86.8)
West Virginia	22	22	21	(95.5)	(0.0)	(0.0)	(90.5)	(90.5)
Wisconsin	75	74	73	(98.6)	(0.0)	(6.8)	(84.9)	(84.9)
Wyoming	4	4	4	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
American Samoa ⁵
Fed. States of Micronesia ⁵	71	69	68	(98.6)	(0.0)	(13.2)	(80.9)	(76.5)
Guam ⁵	53	53	53	(100.0)	(0.0)	(1.9)	(96.2)	(96.2)
Marshall Islands ⁵	27	27	27	(100.0)	(0.0)	(11.1)	(81.5)	(81.5)
N. Mariana Islands ⁵	35	35	35	(100.0)	(0.0)	(2.9)	(97.1)	(97.1)
Puerto Rico ⁵	112	105	105	(100.0)	(2.9)	(1.9)	(92.4)	(92.4)
Republic of Palau ⁵	5	5	4	(80.0)	(0.0)	(25.0)	(75.0)	(75.0)
U.S. Virgin Islands ⁵

¹Overall U.S. percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥75% of cases.

²I=isoniazid; R=rifampin; Z=pyrazinamide; E=ethambutol; S=streptomycin.

³Streptomycin is no longer considered a first-line drug. CDC. Treatment of Tuberculosis. American Thoracic Society, CDC, and Infectious Diseases Society of America. MMWR 2003;52(No. RR-11): 1–77.

⁴Excludes New York City.

⁵Not included in U.S. totals. Data for 2006 are interim as of April 6, 2007.

Note: Excluding cases with no information on drug regimen, 421 (3.1%) persons were not started on any drugs, 12 (<0.1%) were started on one drug, and 1,313 (9.6%) had an initial multidrug regimen other than IR, IRZ, or IRZ,E/S.

Ellipses indicate data not available.