

**All pneumoconioses: Proportionate mortality ratio (PMR) adjusted for age, sex, and race by usual industry, U.S. residents age 15 and over, selected states and years, 1990–1999**

CIC	Industry	Number of Deaths	PMR	95% Confidence Interval	
				LCL	UCL
041	Coal mining	4,623	33.1	32.2	34.1
262	Miscellaneous nonmetallic mineral and stone products	127	7.9	6.6	9.4
040	Metal mining	131	5.2	4.4	6.2
360	Ship and boat building and repairing	176	4.5	3.9	5.2
261	Pottery and related products	27	3.8	2.5	5.6
050	Nonmetallic mining and quarrying, except fuel	69	3.6	2.8	4.6
271	Iron and steel foundries	71	2.8	2.2	3.6
361	Railroad locomotives and equipment	7	2.7	1.1	5.5
252	Structural clay products	29	2.6	1.8	3.8
502	Lumber and construction materials	23	2.3	1.4	3.4
192	Industrial and miscellaneous chemicals	159	1.8	1.5	2.1
211	Other rubber products, and plastics footwear and belting	48	1.5	1.1	1.9
250	Glass and glass products	55	1.4	1.0	1.8

CIC - Census Industry Code      LCL - lower confidence limit      UCL - upper confidence limit

NOTE: The total number of deaths with all pneumoconioses reported was 10,014 in these same selected states and years, and the comparable number of all pneumoconioses deaths in the entire United States for this same time period was 31,455. See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, ICD codes, industry and occupation codes, and list of selected states and years.

SOURCE: National Center for Health Statistics multiple cause-of-death data.