

All pneumoconioses: Years of potential life lost to age 65 and to life expectancy by race and sex, U.S. residents age 15 and over, 1996–2005

Year	White		Black		Other		Total
	Male	Female	Male	Female	Male	Female	
Years of Potential Life Lost to Age 65							
1996	1,815	115	140	-	55	-	2,125
1997	1,810	105	130	-	-	-	2,045
1998	1,665	60	170	30	10	5	1,940
1999	1,340	60	260	25	30	-	1,715
2000	1,745	80	185	25	25	-	2,060
2001	1,610	145	220	35	30	-	2,040
2002	1,535	240	150	10	5	5	1,945
2003	1,735	195	175	55	55	-	2,215
2004	1,790	45	210	25	30	-	2,100
2005	1,820	295	165	5	35	-	2,320
TOTAL	16,865	1,340	1,805	210	275	10	20,505
Years of Potential Life Lost to Life Expectancy							
1996	27,385	1,004	1,566	28	195	8	30,186
1997	25,884	865	1,305	63	111	-	28,228
1998	24,977	746	1,213	95	112	51	27,194
1999	23,921	673	1,397	51	243	-	26,285
2000	24,784	960	1,405	58	198	-	27,405
2001	24,408	899	1,481	87	181	-	27,056
2002	24,061	1,155	1,294	66	105	22	26,703
2003	23,838	1,171	1,373	142	254	-	26,778
2004	23,127	813	1,582	75	146	7	25,750
2005	22,415	1,291	1,214	82	197	-	25,199
TOTAL	244,800	9,577	13,830	747	1,742	88	270,784

- indicates no deaths listed.

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data.