

**All pneumoconioses: Percent of deaths by condition and state, U.S. residents age 15 and over, 1996–2005**

<b>State</b>	<b>Asbestosis</b>	<b>Silicosis</b>	<b>Coal Workers' Pneumoconiosis</b>	<b>Byssinosis</b>	<b>Unspecified/Other Pneumoconioses</b>
Alabama	75.9	4.6	14.4	0.7	5.6
Alaska	86.2	6.9	3.4	-	3.4
Arizona	67.3	15.5	14.1	-	4.5
Arkansas	64.4	8.6	23.9	1.2	3.1
California	85.6	5.5	5.4	0.2	3.7
Colorado	39.1	23.1	24.7	0.7	13.7
Connecticut	85.9	7.8	2.9	-	3.4
Delaware	90.5	3.4	2.0	-	5.4
District of Columbia	60.0	20.0	10.0	-	10.0
Florida	76.6	4.6	14.9	0.3	4.9
Georgia	69.8	10.6	9.4	2.9	8.2
Hawaii	90.0	3.3	3.3	-	3.3
Idaho	72.1	18.6	7.0	-	2.3
Illinois	43.5	9.3	35.3	0.2	14.3
Indiana	33.1	12.6	44.0	-	11.3
Iowa	52.9	14.0	28.9	-	5.8
Kansas	72.8	4.9	16.0	1.2	4.9
Kentucky	9.4	2.9	69.8	0.1	18.6
Louisiana	82.9	7.7	7.7	-	2.3
Maine	95.1	2.8	0.7	0.7	1.4
Maryland	83.8	4.8	8.7	-	4.1
Massachusetts	91.6	4.3	0.2	-	4.6
Michigan	63.9	15.5	15.1	0.2	7.7
Minnesota	78.5	17.5	1.8	0.9	2.2
Mississippi	72.5	4.0	23.3	-	0.8
Missouri	70.5	15.0	10.4	-	5.2
Montana	80.7	11.4	6.1	0.9	1.8
Nebraska	87.0	7.4	5.6	-	-
Nevada	80.9	8.8	4.4	-	5.9
New Hampshire	87.3	7.0	4.2	-	4.2
New Jersey	90.2	3.8	3.7	0.1	2.9
New Mexico	28.1	19.0	39.7	-	13.2
New York	78.6	10.4	5.9	0.3	6.1
North Carolina	70.5	9.1	11.7	5.1	4.8
North Dakota	92.1	2.6	2.6	-	2.6
Ohio	42.5	13.5	31.3	-	14.8
Oklahoma	65.0	9.8	21.1	-	4.1
Oregon	90.2	7.4	2.1	-	0.9
Pennsylvania	16.9	4.0	67.7	0.1	12.9
Rhode Island	78.9	19.7	1.4	1.4	-
South Carolina	81.7	6.8	6.5	2.9	2.5
South Dakota	50.0	35.0	10.0	-	5.0
Tennessee	43.7	6.8	43.4	-	8.5
Texas	86.6	7.6	4.3	0.1	2.8
Utah	30.3	13.3	36.4	-	21.0
Vermont	54.3	31.4	-	2.9	14.3
Virginia	30.3	1.9	54.8	0.2	13.6
Washington	92.9	3.2	3.2	0.1	0.7
West Virginia	11.3	1.5	66.4	-	22.5
Wisconsin	61.9	30.7	4.5	0.8	4.1
Wyoming	30.1	4.1	23.3	-	42.5
<b>TOTAL</b>	<b>49.5</b>	<b>6.3</b>	<b>35.1</b>	<b>0.3</b>	<b>10.1</b>

- indicates no deaths listed.

NOTE: Percentages may total more than 100% due to deaths with multiple pneumoconioses, or to less than 100% due to rounding. See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data.