

Bison *PRNP* genotyping and potential association with *Brucella* spp. seroprevalence

C. M. Seabury*, N. D. Halbert*, P. J. P. Gogan[†], J. W. Templeton* and J. N. Derr*

*Department of Veterinary Pathobiology, College of Veterinary Medicine, Texas A&M University, College Station, TX, USA. [†]Forestry Sciences Laboratory, USGS-Northern Rocky Mountain Science Center, Montana State University, Bozeman, MT, USA

Summary

The implication that host cellular prion protein (PrP^C) may function as a cell surface receptor and/or portal protein for *Brucella abortus* in mice prompted an evaluation of nucleotide and amino acid variation within exon 3 of the prion protein gene (*PRNP*) for six US bison populations. A non-synonymous single nucleotide polymorphism (T50C), resulting in the predicted amino acid replacement M17T (Met → Thr), was identified in each population. To date, no variation (T50; Met) has been detected at the corresponding exon 3 nucleotide and/or amino acid position for domestic cattle. Notably, 80% (20 of 25) of the Yellowstone National Park bison possessing the C/C genotype were *Brucella* spp. seropositive, representing a significant ($P = 0.021$) association between seropositivity and the C/C genotypic class. Moreover, significant differences in the distribution of *PRNP* exon 3 alleles and genotypes were detected between Yellowstone National Park bison and three bison populations that were either founded from seronegative stock or previously subjected to test-and-slaughter management to eradicate brucellosis. Unlike domestic cattle, no indel polymorphisms were detected within the corresponding regions of the putative bison *PRNP* promoter, intron 1, octapeptide repeat region or 3'-untranslated region for any population examined. This study provides the first evidence of a potential association between nucleotide variation within *PRNP* exon 3 and the presence of *Brucella* spp. antibodies in bison, implicating PrP^C in the natural resistance of bison to brucellosis infection.

Keywords bison, brucellosis, natural resistance, *PRNP*.

Introduction

Brucella abortus is a gram-negative facultative intracellular pathogen associated with brucellosis infection in mammals, including undulant fever in humans and abortion and/or infertility in many domestic and wild animals (Acha & Szylres 1980). Although classical breeding studies in domestic cattle (*Bos taurus*) have indicated that natural resistance to *B. abortus* is controlled by two or more genes (Templeton *et al.* 1988), few genes have been investigated and only one, *NRAMP1*, has been implicated in natural resistance to brucellosis in domestic cattle (Feng *et al.* 1996). *NRAMP1* is an integral protein hypothesized to be

involved in phagosome acidification and/or phagosome-lysosome fusion processes within macrophages and has been associated with infectious and autoimmune diseases (Blackwell *et al.* 2001). To date, no association between *NRAMP1* and natural resistance to brucellosis has been demonstrated in American bison (*Bison bison*), a member of the closely related sister genus to *Bos*.

Recently, *B. abortus* heat shock protein 60 was demonstrated to interact with host-encoded cellular prion protein (PrP^C), aiding in the establishment of infection into mouse macrophages and implicating PrP^C as a potential receptor for *B. abortus* (Watarai *et al.* 2003; Watarai 2004). While no consensus regarding the physiological function of PrP^C has been reached to date (Collinge 2001; Aguzzi & Hardt 2003), it may function as a cell-surface receptor for signal transduction (Mouillet-Richard *et al.* 2000). Moreover, the rapid cycling of PrP^C between the cell surface and the early endosome via clathrin-coated vesicles is a physiological process shared by many other cell-surface receptors (Shyng *et al.* 1994). Notably, amino acid polymorphisms associated with the *PRNP* gene are known to influence the phenotypic

Address for correspondence

J.N. Derr, Department of Veterinary Pathobiology, College of Veterinary Medicine, Texas A&M University, College Station, TX 77843-4467, USA.
E-mail: jderr@cvm.tamu.edu

C.M. Seabury and N.D. Halbert contributed equally to this work.

Accepted for publication 5 January 2005

expression of scrapie in sheep and goats (Belt *et al.* 1995; Billinis *et al.* 2002), Creutzfeldt Jakob disease, Gerstman Sträussler Scheinker disease, and Kuru in humans (Collinge 2001), and chronic wasting disease in deer and elk (O'Rourke *et al.* 1999; Johnson *et al.* 2003; O'Rourke *et al.* 2004). Additionally, an association was recently demonstrated between susceptibility to bovine spongiform encephalopathy (BSE) and specific insertion/deletion (indel) polymorphisms within the putative PRNP promoter and intron 1 for several German cattle breeds (Sander *et al.* 2004), supporting the hypothesis that mutations potentially influencing the level of bovine PRNP expression may also influence incubation time and susceptibility to BSE (Bossers *et al.* 1996; Sander *et al.* 2004).

While brucellosis has been effectively eliminated from most bison populations in North America, two substantial reservoirs of brucellosis-infected bison remain in Yellowstone National Park (NP; ID, MT, WY, USA) and Wood Buffalo NP (AB, Canada; Meyer & Meagher 1995; Joly & Messier 2004). Following the initial identification of brucellosis in Yellowstone NP bison (Mohler 1917), multiple studies have documented *Brucella* spp. antibodies (Rush 1932; Tunnicliff & Marsh 1935) as well as *B. abortus* biovars 1 and 2 (Roffe *et al.* 1999; Rhyan *et al.* 2001) in Yellowstone NP bison. Moreover, serological evidence of exposure to *Brucella* spp. has remained relatively high (50–60%) among Yellowstone NP bison for the past 70 years (Tunnicliff & Marsh 1935; Rhyan *et al.* 2001).

Herein, we evaluate the frequencies of bison PRNP exon 3 allelic variants and genotypes for six US bison populations (four federal, one private, and one state) and investigate whether variation within exon 3 is significantly associated with *Brucella* spp. antibodies in Yellowstone NP bison. The distribution of PRNP exon 3 alleles and genotypes for three bison populations that were either founded from

seronegative stock or subjected to test-and-slaughter management to eradicate brucellosis are compared with those observed for Yellowstone NP bison. Additionally, we investigate the frequencies of known bovine PRNP indel polymorphisms within the putative bison PRNP promoter, intron 1 and 3'-untranslated region (UTR; Hills *et al.* 2001, 2003; Sander *et al.* 2004).

Materials and methods

Bison sampling and serology

Collectively, 198 bison from four US national parks, one private, and one state herd were utilized to investigate the frequencies of PRNP allelic variants and genotypes (Table 1). Yellowstone NP samples were obtained non-selectively with regard to pregnancy and/or *Brucella* spp. serology from bison subjected to management actions as they moved to the park boundaries near West Yellowstone, Montana (MT). From these samples, the following selection criteria were used for the current study: approximately equal sex ratio, proportionate sampling from ages 1–10⁺ as determined by established aging methods (Fuller 1959; Moffitt 1998), sampling from various collection dates and West Yellowstone NP locals, and approximately equal distributions of *Brucella* spp. serological phenotypes (seropositive/seronegative). Samples from the remaining five herds were selected randomly from previously established repositories.

Brucella spp. antibodies were detected in serum samples from Yellowstone NP bison postmortem from 1997 ($n = 27$) and 1999 ($n = 39$) using the complement fixation, Rivanol precipitation, buffered acidified plate antigen, standard plate agglutination, standard tube agglutination, and card tests (Nielsen 2002). Because the effectiveness of

Table 1 Distribution of bison samples by local and sex.

Bison herd	Abbreviation	Location	Total	Male	Female
Castle Rock ¹	CR	NM	20	5	15
Grand Teton National Park ²	GT	WY	15	4	11
Henry Mountains ³	HM	UT	20	7	13
Theodore Roosevelt National Park ⁴	TR	ND	10	3	7
Wind Cave National Park ⁵	WC	SD	16	3	13
Yellowstone National Park ⁶	YNP	ID, MT, WY	117	58	59
Total			198	80	118

¹Private herd primarily established from seronegative YNP bison (D. Hunter, pers. comm.).

²Established from YNP bison in 1948 and supplemented with TR bison in 1964 (NPS 1996); population chronically infected with brucellosis (Meyer & Meagher 1995).

³Founded exclusively from YNP bison in 1940s; brucellosis eradicated through test-and-slaughter management in 1960s (Dowling 1990).

⁴South unit; no current or historical evidence of brucellosis (M. Oehler, pers. comm.).

⁵Founded partially from YNP (Coder 1975); brucellosis eradicated through test-and-slaughter management during 1950–1970 (B. Muenchau, pers. comm.).

⁶Samples collected near West Yellowstone, Montana; population chronically infected with brucellosis (Meyer & Meagher 1995).

individual serological tests vary in bison (Williams *et al.* 1997), only samples with consensus serological results were utilized. Fluorescence polarization assay (Gall *et al.* 2000) was used to evaluate the presence of *Brucella* spp. antibodies in samples collected in 2002 ($n = 44$) and 2003 ($n = 7$).

PRNP exon 3 amplification and sequencing

Genomic DNA was isolated by spotting whole blood on Whatman Bioscience FTA[®] Classic Cards following the recommended protocol (Whatman Inc., Clifton, NJ, USA), by using the SUPER QUICK-GENE DNA Isolation kit (AGTC Inc., Denver, CO, USA) on white blood cell isolates or via pulverization/proteinase K treatment of tissues (Halbert *et al.* 2004). Flanking primers SAF1 and SAF2 (Prusiner *et al.* 1993) were used to PCR amplify and sequence bison PRNP exon 3 (Seabury & Derr 2003). Bidirectional sequencing of 96 bison samples from four US federal and one private bison herd was previously employed to evaluate the number of variable sites within bison PRNP exon 3 (Seabury *et al.* 2004a), with 71 of those samples included in the present study. Herein, 127 additional samples were sequenced using SAF1 only. PRNP exon 3 genotypes and allelic variants were determined using SeqScape version 1.01 (Applied Biosystems, Foster City, CA, USA), and representative exon 3 alleles were validated via cloning and subsequent bidirectional sequencing following Seabury & Derr (2003).

PRNP promoter, intron 1 and 3'-UTR genotyping

Fluorescent dye-labelled primer combinations (Seabury *et al.* 2004b) targeting known indel polymorphisms in the bovine PRNP promoter (+23 bp; Sander *et al.* 2004), intron 1 (+12 bp; Hills *et al.* 2001) and 3'-UTR (+14 bp; Hills *et al.* 2003) were utilized to generate genotypic data for bison from Yellowstone NP ($n = 64$; 24 seropositive, 40 seronegative), Wind Cave National Park NP ($n = 20$), Castle Rock ($n = 20$), and Henry Mountains ($n = 20$). PCR reactions and thermal cycling parameters followed Seabury *et al.* (2004b) with the following modifications: 5- μ l reaction with primers PRNP 47784F, 47803R, 67976F and 68070R with a 52 °C annealing temperature, and a second

5- μ l reaction using primers PRNP 49686F and 49777R with the cycling profile of Sander *et al.* (2004). Resulting fragments were co-loaded into a single injection for separation on an ABI 3100 (Applied Biosystems) using an internal size standard (Mapmarker LOW; Bioventures, Inc., Murfreesboro, TN, USA).

Statistical analysis

Bison PRNP allele frequencies, genic and genotypic differentiation employing both the probability test (Raymond & Rousset 1995) and the G-based test of Goudet *et al.* (1996) as well as Hardy-Weinberg equilibrium were calculated using GENEPOP 3.1d (Raymond & Rousset 1995). PRNP exon 3 data for Yellowstone NP bison were subdivided into two groups based on serology (Table 2) and tested for genic and genotypic differentiation. Population differentiation was examined among all bison herds sampled, including Yellowstone NP both with and without subdivision. Because disease cases often differ from controls within individual genotypic classes, we also used a variation of the Z_{\max} test to evaluate associations between specific genotypic classes and serological status in Yellowstone NP bison (Lange 2002). $P < 0.05$ was considered statistically significant.

Results

All bison PRNP exon 3 alleles examined possessed six octapeptide repeats and were identical in sequence with the exception of one non-synonymous single nucleotide polymorphism (SNP) at nucleotide position 50 (T50C). The T50C SNP is predicted to result in the amino acid replacement M17T. None of the tested populations deviated from Hardy-Weinberg expectations for this SNP. Observed PRNP exon 3 allelic and genotypic frequencies for each bison population examined as well as tests of genic and genotypic differentiation are depicted in Table 3. Significant differences in the distributions of PRNP exon 3 alleles and genotypes were detected between Yellowstone NP bison and three other bison populations, each either founded from *Brucella* spp. seronegative stock or subjected to test-and-slaughter management (Table 1). Notably, our samples for

	Age class ¹										
	Total	1	2	3	4	5	6	7	8	9	10+
Yellowstone											
Female seronegative	23	1	1	8	3	3	1	4	1		1
Female seropositive	36	1	1	7	5	7		3	2	1	9
Male seronegative	21	4	1	7	3	1		1		2	2
Male seropositive	37	1	1	16	9	1	1	5		1	2
Sum	117	7	4	38	20	12	2	13	3	4	14

Table 2 Distribution of Yellowstone bison samples by sex, *Brucella* spp. serological status and age.

¹Bison scored in field as 'adult' considered here as 3-year olds

Table 3 Observed *PRNP* exon 3 allelic and genotypic frequencies from six bison populations. The Yellowstone NP population (YNP) is further subdivided by *Brucella* spp. serological phenotypes into seropositive (YNP+) and seronegative (YNP-) classes. Probabilities for population tests of genic and genotypic differentiation are depicted above and below the diagonal, respectively, in the last seven columns ($P < 0.05$ illustrated in bold; all standard errors were <0.01).

Herd	Allele ¹		Genotype ¹				Genic and genotypic differentiation							
	<i>n</i>	T	C	TT	TC	CC	YNP +	YNP -	YNP	GT	TR	CR	WC	HM
YNP +	73	0.514	0.486	0.301	0.425	0.274	-	0.05359	NA ²	0.42297	0.47863	0.00026	0.00583	0.01257
YNP -	44	0.648	0.352	0.409	0.477	0.114	0.06994	-	NA ²	0.66664	0.79848	0.05440	0.18667	0.30602
YNP	117	0.564	0.436	0.342	0.444	0.214	NA ²	NA ²	-	0.84522	0.81513	0.00208	0.02145	0.03399
GT	15	0.600	0.400	0.267	0.667	0.067	0.43923	0.65242	0.85227	-	1.00000	0.05578	0.16801	0.20469
TR	10	0.600	0.400	0.300	0.600	0.100	0.51240	0.78990	0.82476	1.00000	-	0.11272	0.21174	0.25001
CR	20	0.825	0.175	0.700	0.250	0.050	0.00183	0.06113	0.00233	0.04905	0.12022	-	0.76515	0.57814
WC	16	0.781	0.219	0.625	0.313	0.063	0.01063	0.18886	0.03151	0.14260	0.21646	0.78500	-	0.78844
HM	20	0.750	0.250	0.600	0.300	0.100	0.01307	0.31785	0.04908	0.19035	0.38893	0.61824	0.80418	-

See Table 1 for herd abbreviations and histories.

¹Frequencies rounded to three decimal places.

²Not an appropriate statistical test.

the Castle Rock, Wind Cave NP and Henry Mountains bison herds all possessed a significantly higher frequency of the T allele than bison sampled from Yellowstone NP (Table 3). No significant genic or genotypic differentiation was detected in comparisons between Yellowstone NP, Grand Teton NP and Theodore Roosevelt NP bison (Table 3).

A significant overabundance (20 of 25; 80%; $Z_{\max} P = 0.021$) of *Brucella* spp. seropositive bison were detected within the C/C genotypic class in Yellowstone NP. However, no significant overall genic or genotypic ($P = 0.05359$; 0.06994) differentiation was detected between seropositive and seronegative Yellowstone NP bison (Table 3). No significant sex- and/or age-specific associations were detected for seropositive and/or seronegative Yellowstone NP bison (data not shown), although small sample sizes may have precluded detection (Table 2). Significant differences in the distributions of *PRNP* exon 3 alleles and genotypes were detected between seropositive Yellowstone NP bison and our samples for the Castle Rock, Wind Cave NP and Henry Mountains bison populations (Table 3).

No indel polymorphisms were detected within the targeted regions of the putative bison *PRNP* promoter, intron 1 or 3'-UTR. Based on our fluorescent genotyping assay, all bison were fixed for the following alleles (compared directly with sizes in domestic cattle; Seabury *et al.* 2004b): *PRNP* promoter, 100-bp allele; intron 1, 103-bp allele; 3'-UTR, 97-bp allele.

Discussion

The pattern of nucleotide variation observed for bison *PRNP* exon 3, consisting of one non-synonymous SNP of moderate frequency within the signal sequence, is in marked contrast to that of domestic cattle (Heaton *et al.* 2003; Seabury *et al.* 2004a). Furthermore, while no bison *PRNP* indel variation was detected, exon 3 alleles with four to seven octapeptide

repeats (Goldmann *et al.* 1991; Schläpfer *et al.* 1998; Seabury *et al.* 2004a) and indel variation within the promoter, intron 1 and 3'-UTR (Hills *et al.* 2001, 2003; Sander *et al.* 2004) have been documented for domestic cattle. The bison *PRNP* exon 3 T50C non-synonymous SNP and predicted amino acid substitution M17T have not been reported in domestic cattle (Heaton *et al.* 2003; Seabury *et al.* 2004a) or any other species of Bovini (Wopfner *et al.* 1999; Takasuga *et al.* 2003; Seabury *et al.* 2004a). However, the amino acid threonine is widely distributed at the corresponding signal peptide residue for most other mammalian species surveyed (van Rheede *et al.* 2003).

Previous studies have illustrated the sensitivity of PrP^C topology to mutations in the signal sequence and/or transmembrane domain (Hegde *et al.* 1998, 1999; Kim *et al.* 2001; Kim & Hegde 2002), with signal sequence mutations primarily altering the ratio of three topological forms (Hegde *et al.* 1998, 1999) in which PrP^C is synthesized at the endoplasmic reticulum (Kim *et al.* 2001). Furthermore, an association has been suggested between one topological form (PrP^C), an uncleaved signal peptide, and neurodegenerative disease (Hegde *et al.* 1998, 1999; Stewart *et al.* 2001; Kim & Hegde 2002). In addition, Lundberg *et al.* (2002) demonstrated that the N-terminal region of mouse PrP^C (residues 1–28), when uncleaved, is a cell-penetrating peptide capable of transporting large hydrophilic cargoes through a cell membrane. Therefore, the predicted amino acid polymorphism M17T may have functional implications in PrP^C biogenesis and/or entry of *Brucella* spp. into bison host cells.

The collective body of evidence presented herein is consistent with a tentative association between the T/C bison *PRNP* exon 3 alleles and *Brucella* spp. serological status (Table 3). The genotypic differentiation noted between seropositive Yellowstone NP bison and the Castle Rock, Wind Cave NP and Henry Mountains bison populations is largely influenced by

high T/T genotype frequencies observed in these populations (Table 3), consistent with a significant overabundance of seropositive Yellowstone NP bison within the C/C genotypic class. Because the Grand Teton NP, Wind Cave NP, Castle Rock, and Henry Mountains populations were all founded either in part or entirely from Yellowstone NP bison (Table 1), it is unlikely that unique genetic backgrounds lead to the observed differences in PRNP exon 3 allele and genotype frequencies among populations. However, marked differences do exist in the incidence of brucellosis and previous disease management strategies among these populations (Table 1). It is possible that test-and-slaughter management in both the Wind Cave NP and Henry Mountains bison populations, and the seronegative founder event establishing the Castle Rock population, effectively altered the genetic composition of these herds with respect to genes involved in natural resistance to brucellosis.

To date, a cellular receptor for *Brucella* spp. organisms has not been conclusively identified (Gorvel & Moreno 2002). The association noted herein between bison PRNP exon 3 and *Brucella* spp. serological status indirectly supports the findings of Watarai *et al.* (2003), further implicating PrP^C as a potential cellular receptor and/or portal protein for *B. abortus*. However, much remains to be understood about the potential relationship between PrP^C and *B. abortus* infection. The opportunistic sampling of Yellowstone NP bison introduced several uncontrollable factors such as variation in the timing and rate of exposure and use of different serological tests among collection years, perhaps somewhat obscuring the true level of association between PRNP exon 3 variation and immune response to *Brucella* spp. organisms. Moreover, while both the susceptibility and immune response to brucellosis infection in bison and domestic cattle have traditionally been assumed equivalent, differences in the pathogenesis of *B. abortus* infection (Rhyan *et al.* 2001) and in the efficacy of vaccination with strains 19 (Davis *et al.* 1991) and RB51 (Olsen *et al.* 2003) are known to exist. Additionally, while other loci are likely involved in the establishment of brucellosis infection in bison and other bovids, future experimental challenge and association studies aimed at evaluating variation within the PRNP gene for domestic and wild bovids are necessary to further evaluate the magnitude of the relationship between PrP^C and *B. abortus* infection.

Acknowledgements

We thank J.E. Womack, L.G. Adams and T.A. Ficht for insightful discussion, D. Hunter, B. Muenchau, Jim Karpowitz and M. Oehler for herd information, and C. Schutta for SeqScape software. We also thank former Montana State Veterinarian A. Gertonson and current Montana State Veterinarian T. Linfield for permission to collect samples from slaughtered Yellowstone NP bison as well as serological test results.

References

- Acha P. & Szylres B. (1980) Zoonoses. Bacterioses listed B (botulism, brucellosis). In: *Zoonosis and Communicable Diseases Common to Man and Animals*. pp. 28–45. Pan American Health Organization, Washington, DC.
- Aguzzi A. & Hardt W.-D. (2003) Dangerous liaisons between a microbe and the prion protein. *Journal of Experimental Medicine* **198**, 1–4.
- Belt P.B., Muileman I.H., Schreuder B.E., Bos-de Ruijter J., Gielkens A.L. & Smits M.A. (1995) Identification of five allelic variants of the sheep PrP gene and their association with natural scrapie. *Journal of General Virology* **76**, 509–17.
- Billinis C., Panagiotidis C.H., Psychas V., Argyroudis S., Nicolaou A., Leontides S., Papadopoulos O. & Sklaviadis T. (2002) Prion protein gene polymorphisms in natural goat scrapie. *Journal of General Virology* **83**, 713–21.
- Blackwell J.M., Goswami T., Evans C.A.W. *et al.* (2001) SLC11A1 (formerly NRAMP1) and disease resistance. *Cellular Microbiology* **3**, 773–84.
- Bossers A., Schreuder B.E., Muileman I.H., Belt P.B. & Smits M.A. (1996) PrP genotype contributes to determining survival times of sheep with natural scrapie. *Journal of General Virology* **77**, 2669–73.
- Coder G.D. (1975) The national movement to preserve the American buffalo in the United States and Canada between 1880 and 1920. PhD dissertation. The Ohio State University, Columbus, OH.
- Collinge J. (2001) Prion diseases of humans and animals: their causes and molecular basis. *Annual Review of Neuroscience* **24**, 519–50.
- Davis D.S., Templeton J.W., Ficht T.A., Huber J.D., Angus R.D. & Adams L.G. (1991) *Brucella abortus* in bison. II. Evaluation of strain 19 vaccination of pregnant cows. *Journal of Wildlife Diseases* **27**, 258–64.
- Dowling K. (1990) *Buffalo Producer's Guide to Management and Marketing*. R. R. Donneley & Sons, Chicago, IL, USA.
- Feng J., Li Y., Hashad M., Schurr E., Gros P., Adams L.G. & Templeton J.W. (1996) Bovine natural resistance associated macrophage protein 1 (*Nramp1*) gene. *Genome Research* **6**, 956–64.
- Fuller W.A. (1959) The horns and teeth as indicators of age in bison. *Journal of Wildlife Management* **23**, 342–44.
- Gall D., Nielsen K., Forbes L. *et al.* (2000) Validation of the fluorescence polarization assay and comparison to other serological assays for the detection of serum antibodies to *Brucella abortus* in bison. *Journal of Wildlife Diseases* **36**, 469–76.
- Goldmann W., Hunter N., Martin T., Dawson M. & Hope J. (1991) Different forms of the bovine PrP gene have five or six copies of a short, G-C-rich element within the protein-coding exon. *Journal of General Virology* **72**, 201–4.
- Gorvel J.P. & Moreno E. (2002) *Brucella* intracellular life: from invasion to intracellular replication. *Veterinary Microbiology* **90**, 281–97.
- Goudet J., Raymond M., De Meeüs T. & Rousset F. (1996) Testing differentiation in diploid populations. *Genetics* **144**, 1933–40.
- Halbert N.D., Raudsepp T., Chowdhary B.P. & Derr J.N. (2004) Conservation genetic analysis of the Texas State Bison Herd. *Journal of Mammalogy* **85**, 924–31.

- Heaton M.P., Leymaster K.A., Freking B.A., Hawk D.A., Smith T.P.L., Keele J.W., Snelling W.M., Fox J.M., Chitko-McKown C.G. & Laegreid W.W. (2003) Prion gene sequence variation within diverse groups of U.S. sheep, beef cattle, and deer. *Mammalian Genome* **14**, 765–777.
- Hegde R.S., Mastrianni J.A., Scott M.R., Defea K.A., Tremblay P., Torchia M., DeArmond S.J., Prusiner S.B. & Lingappa V.R. (1998) A transmembrane form of the prion protein in neurodegenerative disease. *Science* **279**, 827–34.
- Hegde R.S., Tremblay P., Groth D., DeArmond S.J., Prusiner S.B. & Lingappa V.R. (1999) Transmembrane and genetic prion diseases share a common pathway of neurodegeneration involving transmembrane prion protein. *Nature* **402**, 822–6.
- Hills D., Comincini S., Schlaepfer J., Dolf G., Ferretti L. & Williams J.L. (2001) Complete genomic sequence of the bovine prion gene (PRNP) and polymorphism in its promoter region. *Animal Genetics* **32**, 231–32.
- Hills D., Schlaepfer J., Comincini S., MacLean I., Dolf G., Ferretti L., Olsaker I. & Williams J.L. (2003) Sequence variation in the bovine and ovine PRNP genes. *Animal Genetics* **34**, 183–90.
- Johnson C., Johnson J., Clayton M., McKenzie D. & Aiken J. (2003) Prion protein heterogeneity in free-ranging white-tailed deer within the chronic wasting disease affected region of Wisconsin. *Journal of Wildlife Diseases* **39**, 576–81.
- Joly D.O. & Messier F. (2004) Factors affecting apparent prevalence of tuberculosis and brucellosis in wood bison. *Journal of Animal Ecology* **73**, 623–31.
- Kim S.J. & Hegde R. (2002) Cotranslational partitioning of nascent prion protein into multiple populations at the translocation channel. *Molecular Biology of the Cell* **13**, 3775–86.
- Kim S.J., Rahbar R. & Hegde R. (2001) Combinatorial control of prion protein biogenesis by the signal sequence and transmembrane domain. *Journal of Biological Chemistry* **276**, 26132–40.
- Lange K. (2002) *Mathematical and Statistical Methods for Genetic Analysis*, 2nd edn. Springer-Verlag, New York, USA.
- Lundberg P., Magzoub M., Lindberg M., Hallbrink M., Jarvet J., Eriksson L.E.G., Langel U. & Graslund A. (2002) Cell membrane translocation of the N-terminal (1–28) part of the prion protein. *Biochemical and Biophysical Research Communications* **299**, 85–90.
- Meyer M.E. & Meagher M. (1995) Brucellosis in free-ranging bison (*Bison bison*) in Yellowstone, Grand Teton, and Wood Buffalo National Parks: a review. *Journal of Wildlife Diseases* **31**, 579–98.
- Moffitt S.A. (1998) Aging bison by the incremental cementum growth layers in teeth. *Journal of Wildlife Management* **62**, 1276–80.
- Mohler J.R. (1917) Report of the chief of the Bureau of Animal Industry, pathological division, abortion disease. In: *Annual Reports of the Department of Agriculture*. pp. 105–6. United States Department of Agriculture, Washington, DC, USA.
- Mouillet-Richard S., Ermonval M., Chebassier C., Laplanche J.L., Lehmann S., Launay J.M. & Kellerman O. (2000) Signal transduction through prion protein. *Science* **289**, 1925–28.
- National Park Service (NPS) (1996) *The Jackson Bison Herd: Long Term Management Plan and Environmental Assessment*. National Park Service, Washington, DC.
- Nielsen K. (2002) Diagnosis of brucellosis by serology. *Veterinary Microbiology* **90**, 447–59.
- O'Rourke K.I., Besser T.E., Miller M.W., Cline T.F., Spraker T.R., Jenny A.L., Wild M.A., Zebarth G.L. & Williams E.S. (1999) PrP genotypes of captive and free-ranging Rocky Mountain elk (*Cervus elaphus nelsoni*) with chronic wasting disease. *Journal of General Virology* **80**, 2765–679.
- O'Rourke K.I., Spraker T.R., Hamburg L.K., Besser T.E., Brayton K.A. & Knowles D.P. (2004) Polymorphisms in the prion precursor functional gene but not the pseudogene are associated with susceptibility to chronic wasting disease in white-tailed deer. *Journal of General Virology* **85**, 1339–46.
- Olsen S.C., Jensen A.E., Stoffregen W.C. & Palmer M.V. (2003) Efficacy of calfhood vaccination with *Brucella abortus* strain RB51 in protecting bison against brucellosis. *Research in Veterinary Science* **74**, 17–22.
- Prusiner S.B., Fuzi M., Scott M. *et al.* (1993) Immunologic and molecular biologic studies of prion proteins in bovine spongiform encephalopathy. *Journal of Infectious Diseases* **167**, 602–13.
- Raymond M. & Rousset F. (1995) GENEPOP (Version 1.2): population genetics software for exact tests and ecumenicism. *Journal of Heredity* **86**, 248–9.
- van Rheede T., Smolenaars M.M.W., Madsen O. & de Jong W.W. (2003) Molecular evolution of the mammalian prion protein. *Molecular Biology and Evolution* **20**, 111–21.
- Rhyan J.C., Gidlewski T., Roffe T.J., Aune K., Philo L.M. & Ewalt D.R. (2001) Pathology of brucellosis in bison from Yellowstone National Park. *Journal of Wildlife Diseases* **37**, 101–9.
- Roffe T.J., Rhyan J.C., Aune K., Philo L.M., Ewalt D.R., Gidlewski T. & Hennager S.G. (1999) Brucellosis in Yellowstone National Park bison: quantitative serology and infection. *Journal of Wildlife Management* **63**, 1132–7.
- Rush W.M. (1932) Bang's disease in the Yellowstone National Park buffalo and elk herds. *Journal of Mammalogy* **13**, 371–2.
- Sander P., Hamann H., Pfeiffer I., Wemheuer W., Brenig B., Groschup M., Ziegler U., Distl O. & Leeb T. (2004) Analysis of sequence variability of the bovine prion protein gene (PRNP) in German cattle breeds. *Neurogenetics* **5**, 19–25.
- Schläpfer J., Saitbekova N., Gaillard C. & Dolf G. (1998) A new allelic variant in the bovine prion protein gene (PRNP) coding region. *Animal Genetics* **30**, 386–87.
- Seabury C.M. & Derr J.N. (2003) Identification of a novel ovine PrP polymorphism and scrapie resistant genotypes for St. Croix White and a related composite breed. *Cytogenetic and Genome Research* **102**, 85–8.
- Seabury C.M., Honeycutt R.L., Rooney A.P., Halbert N.D. & Derr J.N. (2004a) Prion protein gene (PRNP) variants and evidence for strong purifying selection in functionally important regions of bovine exon 3. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 15142–7.
- Seabury C.M., Womack J.E., Piedrahita J. & Derr J.N. (2004b) Comparative PRNP genotyping of U.S. cattle sires for potential association with BSE. *Mammalian Genome* **15**, 828–33.
- Shyng S.-L., Heuser J.E. & Harris D.A. (1994) A glycolipid-anchored prion protein is endocytosed via clathrin-coated pits. *Journal of Cellular Biology* **125**, 1239–50.
- Stewart R.S., Drisaldi B. & Harris D.A. (2001) A transmembrane form of the prion protein contains an uncleaved signal peptide and is retained in the endoplasmic reticulum. *Molecular Biology of the Cell* **12**, 881–89.

- Takasuga A., Abe T., Ito T., Watanabe T., Kamatani N. & Sugimoto Y. (2003) Novel prion protein polymorphisms in cattle. *Animal Genetics* **34**, 396–97.
- Templeton J.W., Smith III R. & Adams L.G. (1988) Natural disease resistance in domestic animals. *Journal of the American Veterinary Medical Association* **192**, 1306–15.
- Tunncliffe E.A. & Marsh H (1935) Bang's disease in bison and elk in the Yellowstone National Park and on the National Bison Range. *Journal of the American Veterinary Medical Association* **86**, 745–52.
- Watarai M. (2004) Interaction between *Brucella abortus* and cellular prion protein in lipid raft microdomains. *Microbes and Infection* **6**, 93–100.
- Watarai M., Kim S., Erdenebaatar J., Makino S., Horiuchi M., Shirahata T., Sakaguchi S. & Katamine S. (2003) Cellular prion protein promotes *Brucella* infection into macrophages. *Journal of Experimental Medicine* **198**, 5–17.
- Williams E.S., Cain S.L. & Davis D.S. (1997) Brucellosis: the disease in bison. In: *Brucellosis, Bison, Elk, and Cattle in the Greater Yellowstone Area: Defining the Problem, Exploring Solutions* (Ed. by E.T. Thorne, M.S. Boyce, P. Nicoletti & T.J. Kreeger), pp. 27–32. Greater Yellowstone Interagency Brucellosis Committee and Wyoming Game and Fish Department, Cheyenne, WY.
- Wopfner F., Weidenhöfer G., Schneider R., von Brunn A., Gilch S., Schwarz T.F., Werner T. & Schätzl H.M. (1999) Analysis of 27 mammalian and 9 avian PrPs reveals high conservation of flexible regions of the prion protein. *Journal of Molecular Biology* **289**, 1163–78.