

Minnesota, 2006

Primary and secondary syphilis

- **CASES:** Minnesota reported 47 cases in 2006.
- **RATES:** Minnesota ranked 42 among 50 states, DC, and 3 territories with 0.9 cases per 100,000 population compared to the total rate of 3.3 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Minnesota, the rate among males was 1.7 per 100,000 population compared to the total male rate of 5.7 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the total female rate of 1.0 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


In Minnesota, the race/ethnicity adjusted rates per 100,000 population were 0.8 among whites, 3.0 among blacks, 1.6 among Hispanics, 0.5 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

- The rate among blacks was 3.8 times that of whites.
- The rate among Hispanics was 2 times that of whites.

Congenital syphilis

- **CASES:** Minnesota reported 1 case in 2006.
- **RATES:** Minnesota ranked 28 among 26 states, DC, and 1 territory reporting congenital syphilis cases with a rate of 1.4 cases per 100,000 live births compared to the total rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Minnesota, 1997-2006



Total P&S Syphilis Rates by County, Minnesota, 2006

