TABLE SNR13. Nonfatal occupational injury and illness incidence rates¹ by State and case types, private industry, 2005

| State | Total recordable cases | Cases with days away from work, job transfer, or restriction | | | Other |
|-------------------------------|------------------------------|--|--|---|------------------------------|
| | | Total | Cases with days away from work ² | Cases with job transfer or restriction | Other recordable cases |
| Private industry ³ | 4.6 | 2.4 | 1.4 | 1.0 | 2.2 |
| Alabama | 4.6 | 2.2 | 1.1 | 1.1 | 2.4 |
| Alaska | 6.2 | 3.0 | 2.4 | .6 | 3.2 |
| Arizona | 4.8 | 2.4 | 1.4 | 1.0 | 2.4 |
| Arkansas | 5.0 | 2.4 | 1.2 | 1.1 | 2.6 |
| California | 4.7 | 2.7 | 1.3 | 1.4 | 2.0 |
| California | 4.7 | 2.1 | 1.5 | 1.4 | 2.0 |
| Connecticut | 5.0 | 2.8 | 1.6 | 1.2 | 2.2 |
| Delaware | 3.7 | 1.9 | 1.3 | .6 | 1.8 |
| District of Columbia | 2.0 | .9 | | .2 | 1.0 |
| Florida | 4.5 | 2.4 | 1.2 | 1.2 | 2.1 |
| Georgia | 4.3 | 2.0 | 1.1 | .9 | 2.3 |
| Georgia | 4.3 | 2.0 | 1.1 | .9 | 2.3 |
| Guam | 4.4 | 2.6 | 2.3 | .3 | 1.8 |
| Hawaii | 4.9 | 3.1 | 2.7 | .4 | 1.8 |
| Illinois | 4.1 | 2.4 | 1.3 | 1.0 | 1.8 |
| Indiana | 5.8 | 2.8 | 1.3 | 1.5 | 2.9 |
| lowa | 6.5 | 3.0 | 1.5 | 1.6 | 3.4 |
| 10wa | 0.5 | 3.0 | 1.5 | 1.0 | 5.4 |
| Kansas | 5.3 | 2.4 | 1.2 | 1.2 | 2.9 |
| Kentucky | 6.2 | 3.1 | 1.8 | 1.3 | 3.0 |
| Louisiana | 3.1 | 1.5 | 1.0 | .5 | 1.6 |
| Maine | 7.2 | 3.9 | 1.7 | 2.2 | 3.3 |
| Maryland | 4.2 | 2.2 | 1.5 | .6 | 2.0 |
| Waryland | | 2.2 | 1.0 | .0 | 2.0 |
| Massachusetts | 4.2 | 2.3 | 1.7 | .6 | 1.9 |
| Michigan | 5.3 | 2.7 | 1.2 | 1.4 | 2.6 |
| Minnesota | 5.0 | 2.5 | 1.3 | 1.2 | 2.6 |
| Missouri | 5.4 | 2.6 | 1.2 | 1.4 | 2.8 |
| Montana | 6.6 | 2.9 | 2.2 | .8 | 3.7 |
| | | | | | |
| Nebraska | 5.0 | 2.4 | 1.4 | 1.0 | 2.6 |
| Nevada | 5.7 | 2.9 | 1.5 | 1.4 | 2.8 |
| New Jersey | 3.8 | 2.0 | 1.4 | .6 | 1.8 |
| New Mexico | 4.4 | 2.2 | 1.4 | .8 | 2.2 |
| New York | 3.2 | 1.7 | 1.5 | .2 | 1.5 |
| Nauth Caralla | 4.0 | | | | 4.0 |
| North Carolina | 4.0 | 2.1 | 1.1 | 1.0 | 1.9 |
| Oklahoma | 4.6 | 2.6 | 1.4 | 1.2 | 2.1 |

See footnotes at end of table.

TABLE SNR13. Nonfatal occupational injury and illness incidence rates by State and case types, private industry, 2005 — Continued

| State | Total recordable cases | Cases with days away from work, job transfer, or restriction | | | Other |
|---|---------------------------------|--|--|--|---|
| | | Total | Cases with days away from work ² | Cases with job transfer or restriction | Other recordable cases |
| Oregon Puerto Rico Rhode Island South Carolina Tennessee Texas Utah Vermont | 5.5 3.6 4.8 3.6 5.6 | 2.9 3.4 2.8 1.9 2.5 2.0 2.6 2.8 | 1.7 3.3 2.1 1.1 1.2 1.0 1.2 1.9 | 1.2 (⁴) .7 .9 1.3 1.0 1.4 | 2.5 .9 2.6 1.7 2.3 1.6 3.0 3.4 |
| Virginia | 2.2 6.1 5.5 5.8 | 2.0 1.3 2.9 3.1 3.0 | 1.3 1.2 2.0 2.7 1.7 | .7 .1 .9 .4 1.3 | 2.0 .9 3.2 2.5 2.8 |

 $^{^{1}\,}$ The incidence rates represent the number of injuries and illnesses per 100 full-time workers and were calculated as: (N/EH) x 200,000, where

= number of injuries and illnesses

EΗ = total hours worked employees during the

calendar year

200,000 = base for 100 equivalent full-time workers (working 40 hours per week, 50 weeks per year)

NOTE: Because of rounding, components may not add to totals. SOURCE: Bureau of Labor Statistics, U.S. Department of Labor

October 2006

Days-away-from-work cases include those that result in days away from work with or without job transfer or restriction.
 Data cover all 50 States.
 Incidence rate less than 0.05.