


Figure 6.1. Prevalence of obesity among adults aged 20 years and over: United States, 1997–March 2006




NOTES: Obesity is defined as a Body Mass Index (BMI) of 30 kg/m² or more. The measure is based on self-reported height and weight. The analyses excluded people with unknown height or weight (about 4% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See “About This Release” and Table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2006 NHIS. The estimate for 2006 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2006, 24.8% (95% confidence interval = 23.36–26.34%) of U.S. adults aged 20 years and over were obese, which was not significantly different from the 2005 estimate of 25.4%.

■ The annual prevalence of obesity among U.S. adults aged 20 years and over has increased over time from 19.4% in 1997 to 25.4% in 2005.

Figure 6.2. Prevalence of obesity among adults aged 20 years and over, by age group and sex: United States, January–March 2006




NOTES: Obesity is defined as a Body Mass Index (BMI) of 30 kg/m² or more. The measure is based on self-reported height and weight. The analyses excluded 325 people (5.4%) with unknown height or weight.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2006 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the prevalence of obesity was highest among adults aged 40–59 years (28.5%) and lowest among adults aged 20–39 years (21.7%).
- For adults aged 60 years and over, the prevalence of obesity was higher for women than for men.

Figure 6.3. Age-adjusted prevalence of obesity among adults aged 20 years and over, by sex and race/ethnicity: United States, January–March 2006



NOTES: Obesity is defined as a Body Mass Index (BMI) of 30 kg/m² or more. The measure is based on self-reported height and weight. The analyses excluded 325 people (5.4%) with unknown height or weight. Estimates are age adjusted using the projected 2000 U.S. population as the standard population using five age groups: 20–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2006 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ Non-Hispanic black women were more likely than Hispanic and non-Hispanic white women to be obese.

■ Within the three race/ethnicity groups, the age-adjusted prevalence of obesity was not significantly different for men than for women.

Data tables for figures 6.1–6.3:

Data table for figure 6.1. Prevalence of obesity among adults aged 20 years and over: United States, 1997–March 2006

Year	Percent (95% confidence interval)	
	Crude ¹	Age-adjusted ²
1997	19.4 (18.9-19.9)	19.5 (18.9-20.0)
1998	20.6 (20.1-21.1)	20.6 (20.0-21.1)
1999	21.5 (20.9-22.1)	21.5 (20.9-22.1)
2000	21.8 (21.2-22.4)	21.8 (21.2-22.3)
2001	23.0 (22.4-23.6)	22.9 (22.3-23.5)
2002	23.9 (23.3-24.6)	23.8 (23.2-24.5)
2003	23.7 (23.1-24.3)	23.5 (22.9-24.2)
2004	24.5 (23.9-25.1)	24.3 (23.8-25.0)
2005	25.4 (24.77-26.09)	25.3 (24.66-25.96)
January–March 2006	24.8 (23.36-26.34)	24.7 (23.23-26.17)

¹Crude estimates are presented in the figure.

²Estimates for this Healthy People 2010 Leading Health Indicator are age adjusted using the projected 2000 U.S. population as the standard population using seven age groups: 20–29 years, 30–39 years, 40–49 years, 50–59 years, 60–69 years, 70–79 years, and 80 years and over.

NOTES: Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See “About This Release” and Table III in the Appendix for more details.

DATA SOURCE: NHIS, 1997–March 2006. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for figure 6.2. Prevalence of obesity among adults aged 20 years and over, by age group and sex: United States, January–March 2006

Age and sex	Percent	95% confidence interval
20–39 years		
Total	21.7	19.57-23.76
Male	21.7	18.66-24.64
Female	21.7	18.75-24.61
40–59 years		
Total	28.5	26.07-30.83
Male	30.1	26.48-33.75
Female	26.8	23.64-30.06
60 years and over		
Total	23.7	21.19-26.28
Male	21.0	17.22-24.78
Female	26.1	22.65-29.46
20 years and over: crude¹		
Total	24.8	23.36-26.34
Male	24.9	22.87-26.95
Female	24.8	22.71-26.87
20 years and over: age-adjusted²		
Total	24.7	23.23-26.17
Male	24.5	22.52-26.48
Female	24.8	22.75-26.89

¹Crude estimates are presented in the figure.

²Estimates for this Healthy People 2010 Leading Health Indicator are age adjusted using the projected 2000 U.S. population as the standard population using seven age groups: 20–29 years, 30–39 years, 40–49 years, 50–59 years, 60–69 years, 70–79 years, and 80 years and over.

DATA SOURCE: National Health Interview Survey, 2006. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for figure 6.3. Age-adjusted prevalence of obesity among adults aged 20 years and over, by sex and race/ethnicity: United States, January–March 2006

Sex and race/ethnicity	Percent¹	95% confidence interval
Male		
Hispanic or Latino	26.6	21.17-32.09
Not Hispanic or Latino:		
White, single race	23.1	20.63-25.56
Black, single race	32.3	27.20-37.31
Female		
Hispanic or Latino	25.4	21.30-29.46
Not Hispanic or Latino:		
White, single race	23.2	20.66-25.81
Black, single race	37.1	32.68-41.44

¹Estimates for this Healthy People 2010 Leading Health Indicator are age adjusted using the projected 2000 U.S. population as the standard population using five age groups: 20–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2006. Data are based on household interviews of a sample of the civilian noninstitutionalized population.