


**Figure 14.1. Prevalence of diagnosed diabetes among adults aged 18 years and over: United States, 1997–March 2006**




NOTES: Prevalence of diagnosed diabetes is based on self-report of ever having been diagnosed with diabetes by a doctor or other health professional. Persons reporting “borderline” diabetes status and women reporting diabetes only during pregnancy were not coded as having diabetes in the analyses. The analyses excluded persons with unknown diabetes status (about 0.1% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See “About This Release” and Table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2006 NHIS. The estimate for 2006 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- In early 2006, 8.0% (95% confidence interval = 7.23–8.81%) of adults aged 18 years and over had ever been diagnosed as having diabetes, which was not significantly different from the 2005 estimate of 7.4%.
- From 1997 to early 2006, there was an increasing trend in the prevalence of diagnosed diabetes among adults aged 18 years and over, from 5.1% in 1997 to 8.0% in early 2006.

**Figure 14.2. Prevalence of diagnosed diabetes among adults aged 18 years and over, by age group and sex: United States, January–March 2006**




NOTES: Prevalence of diagnosed diabetes is based on self-report of ever having been diagnosed with diabetes by a doctor or other health professional. Persons reporting “borderline” diabetes status and women reporting diabetes only during pregnancy were not coded as having diabetes in the analyses. The analyses excluded 6 persons (0.1%) with unknown diabetes status.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2006 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the prevalence of diagnosed diabetes increased with age, with the highest rate among adults aged 65 years and over (18.1%) and the lowest rate among adults aged 18–44 years (2.8%).
- For all four age groups, there was no significant difference between women and men in the prevalence of diagnosed diabetes.

**Figure 14.3. Age-sex-adjusted prevalence of diagnosed diabetes among adults aged 18 years and over, by race/ethnicity: United States, January–March 2006**



NOTES: Prevalence of diagnosed diabetes is based on self-report of ever having been diagnosed with diabetes by a doctor or other health professional. Persons reporting “borderline” diabetes status and women reporting diabetes only during pregnancy were not coded as having diabetes in the analyses. The analyses excluded 6 persons (0.1%) with unknown diabetes status. Estimates are age-sex adjusted using the projected 2000 U.S. population as the standard population using four age groups: 18–44 years, 45–54 years, 55–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2006 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ The age-sex-adjusted prevalence of diagnosed diabetes was 10.5% for Hispanic persons, 6.8% for non-Hispanic white persons, and 12.3% for non-Hispanic black persons.

■ The prevalence of diagnosed diabetes was higher among non-Hispanic black persons and Hispanic persons than among non-Hispanic white persons.

## Data tables for figures 14.1–14.3:

**Data table for figure 14.1. Prevalence of diagnosed diabetes among adults aged 18 years and over: United States, 1997–March 2006**

Year	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
1997	5.1 (4.9-5.4)	5.3 (5.1-5.6)
1998	5.3 (5.0-5.6)	5.4 (5.1-5.7)
1999	5.4 (5.1-5.7)	5.5 (5.2-5.8)
2000	5.9 (5.6-6.2)	6.0 (5.7-6.3)
2001	6.4 (6.1-6.7)	6.4 (6.1-6.7)
2002	6.5 (6.2-6.8)	6.5 (6.2-6.8)
2003	6.6 (6.3-6.9)	6.5 (6.2-6.9)
2004	7.0 (6.7-7.4)	6.9 (6.6-7.3)
2005	7.4 (7.10-7.78)	7.3 (6.95-7.57)
January–March 2006	8.0 (7.23-8.81)	7.8 (7.07-8.54)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population using four age groups: 18–44 years, 45–54 years, 55–64 years, and 65 years and over.

NOTES: Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See “About This Release” and Table III in the Appendix for more details.

DATA SOURCE: NHIS, 1997–March 2006. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 14.2. Prevalence of diagnosed diabetes among adults aged 18 years and over, by age group and sex: United States, January–March 2006**

Age and sex	Percent	95% confidence interval
<b>18–44 years</b>		
Total	2.8	2.08-3.48
Male	2.9	1.95-3.88
Female	2.6	1.70-3.59
<b>45–54 years</b>		
Total	7.8	6.19-9.41
Male	8.6	6.05-11.10
Female	7.0	4.93-9.17
<b>55–64 years</b>		
Total	15.6	12.33-18.91
Male	17.2	12.07-22.27
Female	14.2	10.82-17.57
<b>65 years and over</b>		
Total	18.1	15.77-20.40
Male	17.2	13.09-21.24
Female	18.8	15.62-21.91
<b>18 years and over: crude<sup>1</sup></b>		
Total	8.0	7.23-8.81
Male	8.0	6.83-9.27
Female	8.0	7.11-8.89
<b>18 years and over: age-adjusted<sup>2</sup></b>		
Total	7.8	7.07-8.54
Male	8.0	6.87-9.22
Female	7.5	6.70-8.40

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population using four age groups:

18–44 years, 45–54 years, 55–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2006. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 14.3. Age-sex-adjusted prevalence of diagnosed diabetes among adults aged 18 years and over, by race/ethnicity: United States, January–March 2006**

Race/ethnicity	Percent <sup>1</sup>	95% confidence interval
Hispanic or Latino	10.5	8.37-12.64
<b>Not Hispanic or Latino:</b>		
White, single race	6.8	5.96-7.69
Black, single race	12.3	10.10-14.48

<sup>1</sup>Estimates are age-sex adjusted using the projected 2000 U.S. population as the standard population using four age groups: 18–44 years, 45–54 years, 55–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2006. Data are based on household interviews of a sample of the civilian noninstitutionalized population.