

Apache County

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Apache (Arizona) trout	<i>Oncorhynchus gilae apache</i>	Threatened	Yellowish to yellow-olive cutthroat-like trout with large dark spots on body. Dorsal, anal, and caudal fins edged with white. No red lateral band.	Apache, Coconino, Gila, Graham, Greenlee, Navajo	> 5,000 ft	Streams and rivers generally above 6,000 ft. elevation with adequate stream flow and shading; temperatures below 77 degrees F; and substrate composed of boulders, rocks, gravel and some sand and silt.	Presently restricted to drainages in the White Mountains. Hybridization with introduced trout has complicated efforts to maintain the genetic purity of some populations. Special regulations (4d Rule) allow Arizona to manage species as a sport fish.
Black-footed ferret	<i>Mustela nigripes</i>	Endangered	Weasel-like, yellow buff coloration with black on feet, tail tip, and eye mask. It has a blunt light colored nose and is 15-18 inches long and tail length is 5-6 inches.	Apache, Coconino, Navajo, Yavapai	< 10,500	Grassland plains generally found in association with prairie dogs.	Unsurveyed prairie dog towns may be occupied by ferrets or may be appropriate for future reintroduction efforts. The Service developed guidelines for surveying prairie dog towns which are available upon request. No wild populations of this species are currently known to exist in Arizona. Reintroduced population exists in Aubrey Valley (Coconino County), Arizona.
California condor	<i>Gymnogyps californianus</i>	Endangered	Very large vulture (47 in., wingspan to 9 1/2 ft, weight to 22 lbs); adult plumage blackish, immature more brownish; adult wing linings white, immature mottled; head and upper parts of neck bare; yellow-orange in adults, grayish in mature.	Apache, Coconino, Mohave, Navajo, Yavapai	Varies	High desert canyons and plateaus.	Recovery program has reintroduced condors to Northern Arizona, with the first release (6 birds) in December 1996. The release site is located at the Vermillion Cliffs (Coconino County), with an experimental/nonessential area designated for most of Northern Arizona and Southern Utah. The area in Arizona is within a polygon formed by Hwy 191, Interstate 40, and Hwy 93, and extends north of the Arizona-Utah and Nevada borders. Breeding is documented in Arizona.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Chiricahua leopard frog	<i>Lithobates [Rana] chiricahuensis</i>	Threatened	Cream colored tubercles (spots) on a dark background on the rear of the thigh, dorsolateral folds that are interrupted and deflected medially, and a call given out of water distinguish this spotted frog from other leopard frogs.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, Navajo, Pima, Santa Cruz, Yavapai	3,300-8,900 ft	Streams, rivers, backwaters, ponds, and stock tanks that are mostly free from introduced fish, crayfish, and bullfrogs.	Require permanent or nearly permanent water sources. Populations north of the Gila River may be a closely-related, but distinct, undescribed species. A special rule allows take of frogs due to operation and maintenance of livestock tanks on State and private lands.
Little Colorado spinedace	<i>Lepidomeda vittata</i>	Threatened	Small (<4 inches long) silvery minnow.	Apache, Coconino, Navajo	4,000-8,000 ft	Moderate to small streams; found in pools and riffles with water flowing over fine gravel and silt substrate.	Critical habitat includes eighteen miles of East Clear Creek, eight miles of Chevelon Creek, and five miles of Nutrioso Creek.
Loach minnow	<i>Tiaroga cobitis</i>	Threatened	Small (<3 inches) slender, elongated fish, olive colored with dirty white spots at the base of the dorsal and caudal fins. Breeding males vivid red on mouth and base of fins.	Apache, Cochise, Gila, Graham, Greenlee, Navajo, Pinal	< 8,000 ft	Benthic species of small to large perennial streams with swift shallow water over cobble and gravel. Recurrent flooding and natural hydrograph important.	Presently found in Aravaipa Creek, Deer Creek, Turkey Creek, Blue River, Campbell Blue Creek, San Francisco River, Eagle Creek, North Fork of the East Fork Black River, and White River in Arizona, and Dry Blue Creek, Pace Creek, Frieborn Creek, the Tularosa River, West Fork Gila River, and the mainstem upper Gila River in New Mexico. Populations have been recently reintroduced in Hot Springs and Redfield canyons in Cochise and Graham counties; Fossil Creek in Gila County; and Bonita Creek in Graham County Arizona. Critical habitat (72 FR 13356-13422, March 21, 2007) found in Apache, Graham, Greenlee, and Pinal counties, Arizona, as well as portions of the Blue River, San Francisco River, Tularosa River, Negrito Creek, Pace Creek, Dry Blue Creek, Frieborn Creek, Whitewater Creek, Gila River, and its West, Middle, and East Forks in Catron, Grant, and Hidalgo counties in New Mexico.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Mexican gray wolf	<i>Canis lupus baileyi</i>	Endangered	Large dog-like carnivore. Head and feet are large in proportion to rest of body. Coat color varies with mix of brown, rust, black, gray, and white. Distinct white lip line around mouth. Adults weigh between 60-90 pounds.	Apache, Gila, Greenlee, Navajo	4,000-12,000 ft	Chaparral, woodland, and forested areas. May cross desert areas.	In March 1998, Mexican gray wolves were reintroduced as an experimental nonessential (10j) population under a program to re-establish the subspecies to a portion of its historical range. Wolves are released within the experimental boundary into a designated area known as the "Blue Range Wolf Recovery Area" (BRWRA) located in the Apache National Forest in Greenlee and Apache counties. Mexican gray wolves found outside of the experimental nonessential boundary are considered endangered. In 2002, the White Mountain Apache tribe (WMAT) became one of the lead agencies for the reintroduction and allowed wolves on their lands. This effectively expanded the experimental nonessential population into Navajo, Apache, and Gila counties on WMAT lands.
Mexican spotted owl	<i>Strix occidentalis lucida</i>	Threatened	Medium sized with dark eyes and no ear tufts. Brownish and heavily spotted with white or beige.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai	4,100-9,000 ft	Nests in canyons and dense forests with multi-layered foliage structure.	Generally nest in older forests of mixed conifer or ponderosa pine/gambel oak type, in canyons, and use variety of habitats for foraging. Sites with cool microclimates appear to be of importance or are preferred. Critical habitat was finalized on August 31, 2004 (69 FR 53182) in Arizona in Apache, Cochise, Coconino, Gila, Graham, Greenlee, Maricopa, Navajo, Pima, Pinal, Santa Cruz, and Yavapai counties.
Navajo sedge	<i>Carex specuicola</i>	Threatened	Perennial forb with triangular stems, elongated rhizomes. Flower: white June and July.	Apache, Coconino, Navajo	5,700-6,000 ft	Silty soils at shady seeps and springs.	Designated critical habitat is on the Navajo Nation near Inscription House Ruins. Found at seep springs on vertical cliffs of pink-red Navajo sandstone.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Southwestern willow flycatcher	<i>Empidonax traillii extimus</i>	Endangered	Small passerine (about 6 inches) grayish-green back and wings, whitish throat, light olive-gray breast and pale yellowish belly. Two wingbars visible. Eye-ring faint or absent.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 8,500 ft	Cottonwood/willow and tamarisk vegetation communities along rivers and streams.	Migratory riparian-obligate species that occupies breeding habitat from late April to September. Distribution within its range is restricted to riparian corridors. Difficult to distinguish from other members of the <i>Empidonax</i> complex by sight alone. Training seminar required for those conducting flycatcher surveys. Critical habitat was finalized on October 19, 2005 (50 CFR 60886). In Arizona there are critical habitat segments in Apache, Cochise, Gila, Graham, Greenlee, Maricopa, Mohave, Pima, Pinal, and Yavapai counties.
Zuni fleabane	<i>Erigeron rhizomatus</i>	Threatened	Herbaceous perennial that grows in clusters of numerous erect unbranched stems up to 2.0 feet tall. Flower heads solitary; pale blue ray flowers and yellow disk flowers.	Apache	7,300-8,000 ft	Selenium-rich red or gray detrital clay soils derived from the Chinle and Baca formations.	Only one Arizona location; other 28 sites in Sawtooth Mountains and northwestern part of the Datil Mountains in Catron County, New Mexico. Two sites also on the northwest side of the Zuni Mountains in McKinley County, New Mexico.
New Mexican Jumping Mouse	<i>Zapus hudsonius luteus</i>	Candidate	Small rodent with extremely long tail and long hind feet. Pelage is coarse with a broad dorsal band of brown or yellowish brown darkened with brownish black hairs; sides paler; under parts white or sometimes suffused with yellowish color. Back of the forefeet and hind feet are grayish white; tail is sparsely haired and distinctly bicolor (dark brown above and yellowish white below). The head is small, narrow, and relatively high crowned. The nose is short and pointed. They are the only mammal with 18 teeth.	Apache	< 8,000 ft	Nests in dry soils but also uses moist, streamside, dense riparian/wetland vegetation.	The New Mexican jumping mouse is diminished to 6 populations in the White Mountains, Arizona. A inventory and status assessment throughout its historical range in Arizona is currently being conducted.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Northern Mexican Gartersnake	<i>Thamnophis eques megalops</i>	Candidate	Background color ranges from olive, olive-brown, to olive-gray. Body has three yellow or light colored stripes running down the length of the body, darker towards tail. Species distinguished from other native gartersnakes by the lateral stripes reaching the 3rd and 4th scale rows. Paired black spots extend along dorsolateral fields.	Apache, Cochise, Coconino, Gila, Graham, Navajo, Pima, Pinal, Santa Cruz, Yavapai	130-8,500 ft	Cienegas, stock tanks, large-river riparian woodlands and forests, streamside gallery forests.	Core population areas in the U.S. include mid/upper Verde River drainage, mid/lower Tonto Creek, and the San Rafael Valley and surrounding area. Status on tribal lands unknown. Distributed south into Mexico along the Sierra Madre Occidental and Mexican Plateau. Strongly associated with the presence of a native prey base including leopard frogs and native fish.
Three Forks springsnail	<i>Pyrgulopsis trivialis</i>	Candidate	Minute hydrobiid snail; shell ovate to narrowly conic; height 0.05 -0.17 inches; whorls 2.5-5.0	Apache	8,000-8,500 ft	Rheocrene springs, seeps, marshes, spring pools, outflows and diverse lotic waters commonly referred to as cienegas.	Distribution limited to Three Forks and Boneyard Spring complexes in the North Fork of the East Fork Black River watershed.
Yellow-billed cuckoo	<i>Coccyzus americanus</i>	Candidate	Medium-sized bird with a slender, long-tailed profile, slightly down-curved bill that is blue-black with yellow on the lower half. Plumage is grayish-brown above and white below, with rufous primary flight feathers.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 6,500 ft	Large blocks of riparian woodlands (cottonwood, willow, or tamarisk galleries).	Neotropical migrant that winters primarily in South America and breeds primarily in the U.S. (but also in southern Canada and northern Mexico). As a migrant it is rarely detected; can occur outside of riparian areas. Cuckoos are found nesting statewide, mostly below 5,000 feet in central, western, and southeastern Arizona. Concern for cuckoos are primarily focused upon alterations to its nesting and foraging habitat. Nesting cuckoos are associated with relatively dense, wooded, streamside riparian habitat, with varying combinations of Fremont cottonwood, willow, velvet ash, Arizona walnut, mesquite, and tamarisk. Some cuckoos have also been detected nesting in velvet mesquite, netleaf hackberry, Arizona sycamore, Arizona alder, and some exotic neighborhood shade trees.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Zuni bluehead sucker	<i>Catostomus discorbulus yarrowi</i>	Candidate	Fusiform, slender, with a terminal mouth. Bluish head, silvery tan to dark green above, silvery to yellowish or dirty-white below. Sexually mature bluehead suckers range between 3.5 to 8 inches in length.	Apache	> 6,000 ft	Small streams in low-velocity, moderate deep pools, and pool-runs with seasonal dense algae. Young prefer quieter shallow areas near shoreline.	Limited to possibly one creek in Arizona and to the headwaters of Zuni River drainage in New Mexico. A Statewide Conservation Agreement between Arizona Game and Fish Department for six non-listed native fish, was finalized in December 2006.
Arizona willow	<i>Salix arizonica</i>	Conservation Agreement	Woody, perennial shrub reaching up to 8.5 feet tall; grows as a prostrate mat to large hedge or thicket plant; has small, egg-shaped leaves; new branches are yellow-green, previous years branches are bright red.	Apache	> 8,000 ft	Unshaded or partially shaded wet meadows, streambanks and cienegas; typically found in or adjacent to perennial water.	Known in the vicinity of Mount Baldy, on the Apache-Sitgreaves National Forest, and private land. Conservation agreement between the Service, Forest Service, and National Park Service finalized in April 1995.
Gooddings onion	<i>Allium gooddingii</i>	Conservation Agreement	Herbaceous perennial plant; broad, flat, rather blunt leaves; flowering stalk 14-18 inches tall, flattened, and narrowly winged toward apex; fruit is broader than long; seeds are short and thick.	Apache, Greenlee, Pima	7,500-11,250 ft	Shaded sites on north-trending drainages, on slopes, or in narrow canyons, within mixed conifer and spruce fir forests.	Known from the White, Santa Catalina, and Chuska Mountains. Also found in New Mexico on the Lincoln and Gila National Forests. A Conservation Agreement between the Service and the Forest Service signed in February 1998.
American peregrine falcon	<i>Falco peregrinus anatum</i>	Delisted	A crow-sized falcon with slate blue-gray on the back and wings, and white on the underside; a black head with vertical "bandit's mask" pattern over the eyes; long pointed wings; and a long wailing call made during breeding. Very adept flyers and hunters, reaching diving speeds of 200 mph.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	3,500-9,000 ft	Areas with rocky, steep cliffs, primarily near water, where prey (primarily shorebirds, songbirds, and waterfowl) concentrations are high. Nests are found on ledges of cliffs, and sometimes on man-made structures such as office towers and bridge abutments.	Species recovered with over 1,650 breeding birds in the US and Canada.