

La Paz County

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Bald eagle	<i>Haliaeetus leucocephalus</i>	Threatened	Large, adults have white head and tail. Height 28-38 inches; wingspan 66-96 inches. Dark with varying degrees of mottled brown plumage. Feet bare of feathers.	Gila, Graham, La Paz, Maricopa, Mohave, Pinal, Yavapai, Yuma	Varies	Large trees or cliffs near water (reservoirs, rivers, and streams) with abundant prey.	Some birds are nesting residents while a larger number winters along rivers and reservoirs. Once endangered (32 FR 4001, 03-11-1967; 43 FR 6233, 02-14-78) because of reproductive failures from pesticide poisoning and loss of habitat, this species was downlisted to threatened on August 11, 1995, and delisted August 8, 2007. Threatened status reinstated for Desert nesting bald eagles.
Bonytail chub	<i>Gila elegans</i>	Endangered	Large (12-14 up to 24 inches) minnow characterized by small head, large fins, slightly humped back and long thin caudal peduncle.	La Paz, Mohave	< 4,000 ft	Warm, swift, turbid mainstem rivers of the Colorado River basin, reservoirs in lower basin.	Endemic to Colorado River Basin. Rarest of Colorado River fish. Population augmentation is ongoing in Lake Mohave and Lake Havasu. Critical habitat includes the Colorado River from Hoover Dam to Davis Dam and another section of the Colorado River from the northern boundary of Havasu National Wildlife Refuge to Parker Dam including Lake Havasu in Mohave County, Arizona. Additional critical habitat is located in Colorado, Utah, Nevada, and California.
California Brown pelican	<i>Pelecanus occidentalis californicus</i>	Proposed delisted	Large, dark gray-brown water bird with webbed feet, pouch underneath its long bill, and wingspan of 7 ft. Adults have a white head and neck, brownish black breast, and silver gray upper parts.	La Paz, Maricopa, Yuma	Varies	Coastal land and islands; species found occasionally around Arizona's lakes and rivers.	Subspecies endangered due to pesticides. Considered an uncommon transient in Arizona. Most observations recorded along the Colorado River and in the Gila Valley. Individuals known to wander up from Mexico in summer and fall. There are no breeding records for Arizona. Populations also exist in California and Mexico.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Razorback sucker	<i>Xyrauchen texanus</i>	Endangered	Large, up to 3 feet long and up to 6 lbs, high sharp-edged keel-like hump behind the head. Head flattened on top. Olive-brown above to yellowish below.	Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Pinal, Yavapai, Yuma	< 6,000 ft	Riverine and lacustrine areas, generally not in fast moving water and may use backwaters.	Big River fish also found in Horseshoe reservoir (Maricopa County). Critical habitat includes the 100-year floodplain of the river through the Grand Canyon from confluence with Paria River to Hoover Dam Hoover Dam to Davis Dam; Parker Dam to Imperial Dam. Also Gila River from Arizona/New Mexico border to Coolidge Dam; and Salt River from Hwy 60/SR77 Bridge to Roosevelt Dam; Verde River from FS boundary to Horseshoe Lake.
Southwestern willow flycatcher	<i>Empidonax traillii extimus</i>	Endangered	Small passerine (about 6 inches) grayish-green back and wings, whitish throat, light olive-gray breast and pale yellowish belly. Two wingbars visible. Eye-ring faint or absent.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 8,500 ft	Cottonwood/willow and tamarisk vegetation communities along rivers and streams.	Migratory riparian-obligate species that occupies breeding habitat from late April to September. Distribution within its range is restricted to riparian corridors. Difficult to distinguish from other members of the Empidonax complex by sight alone. Training seminar required for those conducting flycatcher surveys. Critical habitat was finalized on October 19, 2005 (50 CFR 60886). In Arizona there are critical habitat segments in Apache, Cochise, Gila, Graham, Greenlee, Maricopa, Mohave, Pima, Pinal, and Yavapai counties.
Yuma clapper rail	<i>Rallus longirostris yumanensis</i>	Endangered	Water bird with long legs and short tail. Long, slender decurved bill. Mottled brown or gray on its rump. Flanks and undersides are dark gray with narrow vertical stripes producing a barring effect.	Gila, La Paz, Maricopa, Mohave, Pinal, Yuma	< 4,500 ft	Fresh water and brackish marshes.	Species is associated with dense emergent riparian vegetation. Requires wet substrate (mudflat, sandbar) with dense herbaceous or woody vegetation for nesting and foraging. Channelization and marsh destruction are primary sources of habitat loss.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Yellow-billed cuckoo	<i>Coccyzus americanus</i>	Candidate	Medium-sized bird with a slender, long-tailed profile, slightly down-curved bill that is blue-black with yellow on the lower half. Plumage is grayish-brown above and white below, with rufous primary flight feathers.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 6,500 ft	Large blocks of riparian woodlands (cottonwood, willow, or tamarisk galleries).	Neotropical migrant that winters primarily in South America and breeds primarily in the U.S. (but also in southern Canada and northern Mexico). As a migrant it is rarely detected; can occur outside of riparian areas. Cuckoos are found nesting statewide, mostly below 5,000 feet in central, western, and southeastern Arizona. Concern for cuckoos are primarily focused upon alterations to its nesting and foraging habitat. Nesting cuckoos are associated with relatively dense, wooded, streamside riparian habitat, with varying combinations of Fremont cottonwood, willow, velvet ash, Arizona walnut, mesquite, and tamarisk. Some cuckoos have also been detected nesting in velvet mesquite, netleaf hackberry, Arizona sycamore, Arizona alder, and some exotic neighborhood shade trees.
American peregrine falcon	<i>Falco peregrinus anatum</i>	Delisted	A crow-sized falcon with slate blue-gray on the back and wings, and white on the underside; a black head with vertical "bandit's mask" pattern over the eyes; long pointed wings; and a long wailing call made during breeding. Very adept flyers and hunters, reaching diving speeds of 200 mph.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	3,500-9,000 ft	Areas with rocky, steep cliffs, primarily near water, where prey (primarily shorebirds, songbirds, and waterfowl) concentrations are high. Nests are found on ledges of cliffs, and sometimes on man-made structures such as office towers and bridge abutments.	Species recovered with over 1,650 breeding birds in the US and Canada.