

# Yavapai County

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Arizona cliffrose	<i>Purshia subintegra</i>	Endangered	Evergreen shrub of the rose family (Roseaceae). Bark pale gray and shreddy. Young twigs covered with dense hairs. Leaves have 1-5 lobes and edges curl downward (revolute). Flowers: 5 petals, white or yellow <0.5 inches long.	Graham, Maricopa, Mohave, Yavapai	< 4,000 ft	White limestone soils derived from tertiary lakebed deposits.	Occurs in central Arizona at Horseshoe Lake, in the Burro Creek drainage, and near Cottonwood in the Verde Valley.
Bald eagle	<i>Haliaeetus leucocephalus</i>	Threatened	Large, adults have white head and tail. Height 28-38 inches; wingspan 66-96 inches. Dark with varying degrees of mottled brown plumage. Feet bare of feathers.	Gila, Graham, La Paz, Maricopa, Mohave, Pinal, Yavapai, Yuma	Varies	Large trees or cliffs near water (reservoirs, rivers, and streams) with abundant prey.	Some birds are nesting residents while a larger number winters along rivers and reservoirs. Once endangered (32 FR 4001, 03-11-1967; 43 FR 6233, 02-14-78) because of reproductive failures from pesticide poisoning and loss of habitat, this species was downlisted to threatened on August 11, 1995, and delisted August 8, 2007. Threatened status reinstated for Desert nesting bald eagles.
Black-footed ferret	<i>Mustela nigripes</i>	Endangered	Weasel-like, yellow buff coloration with black on feet, tail tip, and eye mask. It has a blunt light colored nose and is 15-18 inches long and tail length is 5-6 inches.	Apache, Coconino, Navajo, Yavapai	< 10,500	Grassland plains generally found in association with prairie dogs.	Unsurveyed prairie dog towns may be occupied by ferrets or may be appropriate for future reintroduction efforts. The Service developed guidelines for surveying prairie dog towns which are available upon request. No wild populations of this species are currently known to exist in Arizona. Reintroduced population exists in Aubrey Valley (Coconino County), Arizona.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
California condor	<i>Gymnogyps californianus</i>	Endangered	Very large vulture (47 in., wingspan to 9 1/2 ft, weight to 22 lbs); adult plumage blackish, immature more brownish; adult wing linings white, immature mottled; head and upper parts of neck bare; yellow-orange in adults, grayish in mature.	Apache, Coconino, Mohave, Navajo, Yavapai	Varies	High desert canyons and plateaus.	Recovery program has reintroduced condors to Northern Arizona, with the first release (6 birds) in December 1996. The release site is located at the Vermillion Cliffs (Coconino County), with an experimental/nonessential area designated for most of Northern Arizona and Southern Utah. The area in Arizona is within a polygon formed by Hwy 191, Interstate 40, and Hwy 93, and extends north of the Arizona-Utah and Nevada borders. Breeding is documented in Arizona.
Chiricahua leopard frog	<i>Lithobates [Rana] chiricahuensis</i>	Threatened	Cream colored tubercles (spots) on a dark background on the rear of the thigh, dorsolateral folds that are interrupted and deflected medially, and a call given out of water distinguish this spotted frog from other leopard frogs.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, Navajo, Pima, Santa Cruz, Yavapai	3,300-8,900 ft	Streams, rivers, backwaters, ponds, and stock tanks that are mostly free from introduced fish, crayfish, and bullfrogs.	Require permanent or nearly permanent water sources. Populations north of the Gila River may be a closely-related, but distinct, undescribed species. A special rule allows take of frogs due to operation and maintenance of livestock tanks on State and private lands.
Colorado pikeminnow	<i>Ptychocheilus lucius</i>	Endangered	Largest American minnow (up to 6 feet and 80 lbs) dusky-green, slender body with gold flecks on the dorsal surface. Head long and slender.	Gila, Yavapai	< 4,000 ft	Warm, swift, turbid mainstem rivers. Prefers eddies and pools.	Experimental non-essential population (treated as proposed threatened species). No critical habitat in Arizona.
Desert pupfish	<i>Cyprinodon macularius</i>	Endangered	Small (2 inches) smoothly rounded body shape with narrow vertical bars on the sides. Breeding males blue on head and sides with yellow on tail. Females and juveniles tan to olive colored back and silvery sides.	Cochise, Graham, Maricopa, Pima, Santa Cruz, Yavapai	< 4,000 ft	Shallow springs, small streams, and marshes. Tolerates saline and warm water.	Two subspecies are recognized: Desert Pupfish ( <i>C.m. macularis</i> ) and Quitobaquito Pupfish ( <i>C.m. eremus</i> ). Critical habitat includes Quitobaquito Springs, Pima County, portions of San Felipe Creek, Carrizo Wash, and Fish Creek Wash, Imperial County, California.
Gila chub	<i>Gila intermedia</i>	Endangered	Deep compressed body, flat head. Dark olive-gray color above, silver sides. Endemic to Gila River Basin.	Cochise, Gila, Graham, Greenlee, Pima, Pinal, Santa Cruz, Yavapai	2,000-5,500 ft	Pools, springs, cienegas, and streams.	Found on multiple private lands, including the Nature Conservancy and the Audubon Society. Also occurs on Federal and state lands and in Sonora, Mexico. Critical habitat occurs in Cochise, Gila, Graham, Greenlee, Pima, Pinal, Santa Cruz, and Yavapai counties.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Gila topminnow	<i>Poeciliopsis occidentalis occidentalis</i>	Endangered	Small (2 inches), guppy-like, live bearing, lacks dark spots on its fins. Breeding males are jet black with yellow fins.	Cochise, Gila, Graham, Maricopa, Pima, Santa Cruz, Yavapai	< 4,500 ft	Small streams, springs, and cienegas vegetated shallows.	Species historically also occurred in backwaters of large rivers but is currently isolated to small streams and springs.
Mexican spotted owl	<i>Strix occidentalis lucida</i>	Threatened	Medium sized with dark eyes and no ear tufts. Brownish and heavily spotted with white or beige.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai	4,100-9,000 ft	Nests in canyons and dense forests with multi-layered foliage structure.	Generally nest in older forests of mixed conifer or ponderosa pine/gambel oak type in canyons, and use variety of habitats for foraging. Sites with cool microclimates appear to be of importance or are preferred. Critical habitat was finalized on August 31, 2004 (69 FR 53182) in Arizona in Apache, Cochise, Coconino, Gila, Graham, Greenlee, Maricopa, Navajo, Pima, Pinal, Santa Cruz, and Yavapai counties.
Razorback sucker	<i>Xyrauchen texanus</i>	Endangered	Large, up to 3 feet long and up to 6 lbs, high sharp-edged keel-like hump behind the head. Head flattened on top. Olive-brown above to yellowish below.	Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Pinal, Yavapai, Yuma	< 6,000 ft	Riverine and lacustrine areas, generally not in fast moving water and may use backwaters.	Big River fish also found in Horseshoe reservoir (Maricopa County). Critical habitat includes the 100-year floodplain of the river through the Grand Canyon from confluence with Paria River to Hoover Dam Hoover Dam to Davis Dam; Parker Dam to Imperial Dam. Also Gila River from Arizona/New Mexico border to Coolidge Dam; and Salt River from Hwy 60/SR77 Bridge to Roosevelt Dam; Verde River from FS boundary to Horseshoe Lake.
Southwestern willow flycatcher	<i>Empidonax traillii extimus</i>	Endangered	Small passerine (about 6 inches) grayish-green back and wings, whitish throat, light olive-gray breast and pale yellowish belly. Two wingbars visible. Eye-ring faint or absent.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 8,500 ft	Cottonwood/willow and tamarisk vegetation communities along rivers and streams.	Migratory riparian-obligate species that occupies breeding habitat from late April to September. Distribution within its range is restricted to riparian corridors. Difficult to distinguish from other members of the Empidonax complex by sight alone. Training seminar required for those conducting flycatcher surveys. Critical habitat was finalized on October 19, 2005 (50 CFR 60886). In Arizona there are critical habitat segments in Apache, Cochise, Gila, Graham, Greenlee, Maricopa, Mohave, Pima, Pinal, and Yavapai counties.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Spikedace	<i>Meda fulgida</i>	Threatened	Small (<3 inches) slim fish with silvery sides and "spine" on dorsal fin. Breeding males are a brassy golden color.	Cochise, Graham, Greenlee, Gila, Navajo, Pinal, Yavapai	< 6,000 ft	Medium to large perennial streams with moderate to swift velocity waters over cobble and gravel substrate. Recurrent flooding and natural hydrograph important to withstand invading exotic species.	Presently found in Aravaipa Creek, Eagle Creek, Verde River, and the Gila River from the San Pedro River to Ashurst-Hayden Dam in Arizona, and the Gila River and its East and West Forks in New Mexico. Populations have been recently reintroduced in Hot Springs and Redfield canyons in Cochise and Graham counties; Fossil Creek in Gila County; and Bonita Creek in Graham County Arizona. Critical habitat (72 FR 13356-13422, March 21, 2007) in Graham, Greenlee, Pinal, and Yavapai counties in Arizona, and in Catron Grant, and Hidalgo counties in New Mexico.
Headwater chub	<i>Gila nigra</i>	Candidate	Streamlined, dark gray to brown fish, often with longitudinal stripes on the sides. Adults reach a maximum size of about 12 inches.	Yavapai, Gila, Graham	3,000-6,700 ft	Medium-sized streams in large, deep pools often associated with cover such as undercut banks or deep places created by trees or rocks.	Occurs in the East Verde River and tributaries, Fossil Creek, Wet Bottom Creek, Deadman Creek, Tonto Creek and tributaries, San Carlos River, Ash Creek, and the upper Gila River in New Mexico. Statewide Conservation Agreement with Arizona Game and Fish was signed on December 2003.
Northern Mexican Gartersnake	<i>Thamnophis eques megalops</i>	Candidate	Background color ranges from olive, olive-brown, to olive-gray. Body has three yellow or light colored stripes running down the length of the body, darker towards tail. Species distinguished from other native gartersnakes by the lateral stripes reaching the 3rd and 4th scale rows. Paired black spots extend along dorsolateral fields.	Apache, Coconino, Cochise, Gila, Graham, Navajo, Pima, Pinal, Yavapai	130-8,500 ft	Cienegas, stock tanks, large-river riparian woodlands and forests, streamside gallery forests.	Core population areas in the U.S. include mid/upper Verde River drainage, mid/lower Tonto Creek, and the San Rafael Valley and surrounding area. Status on tribal lands unknown. Distributed south into Mexico along the Sierra Madre Occidental and Mexican Plateau. Strongly associated with the presence of a native prey base including leopard frogs and native fish.
Page springsnail	<i>Pyrgulopsis morrisoni</i>	Candidate	Small (<0.11 inches) snail with ovate shell with fine growth lines.	Yavapai	3,300-3,600 ft	Permanently saturated cienegas, firm substrate like cobble, gravel, woody debris, and aquatic vegetation.	Locally endemic to the Upper Verde River drainage. All extant populations exist in a series of springs within a one mile area west of Oak Creek near Page Springs. A Candidate Conservation Agreement with the Arizona Game and Fish Department is currently under development.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Yellow-billed cuckoo	<i>Coccyzus americanus</i>	Candidate	Medium-sized bird with a slender, long-tailed profile, slightly down-curved bill that is blue-black with yellow on the lower half. Plumage is grayish-brown above and white below, with rufous primary flight feathers.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 6,500 ft	Large blocks of riparian woodlands (cottonwood, willow, or tamarisk galleries).	Neotropical migrant that winters primarily in South America and breeds primarily in the U.S. (but also in southern Canada and northern Mexico). As a migrant it is rarely detected; can occur outside of riparian areas. Cuckoos are found nesting statewide, mostly below 5,000 feet in central, western, and southeastern Arizona. Concern for cuckoos are primarily focused upon alterations to its nesting and foraging habitat. Nesting cuckoos are associated with relatively dense, wooded, streamside riparian habitat, with varying combinations of Fremont cottonwood, willow, velvet ash, Arizona walnut, mesquite, and tamarisk. Some cuckoos have also been detected nesting in velvet mesquite, netleaf hackberry, Arizona sycamore, Arizona alder, and some exotic neighborhood shade trees.
American peregrine falcon	<i>Falco peregrinus anatum</i>	Delisted	A crow-sized falcon with slate blue-gray on the back and wings, and white on the underside; a black head with vertical "bandit's mask" pattern over the eyes; long pointed wings; and a long wailing call made during breeding. Very adept flyers and hunters, reaching diving speeds of 200 mph.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	3,500-9,000 ft	Areas with rocky, steep cliffs, primarily near water, where prey (primarily shorebirds, songbirds, and waterfowl) concentrations are high. Nests are found on ledges of cliffs, and sometimes on man-made structures such as office towers and bridge abutments.	Species recovered with over 1,650 breeding birds in the US and Canada.
Arizona agave	<i>Agave arizonica</i>	Delisted	Member of the agave family. Has rosettes of bright green leaves, 17-24cm long and 2-4cm wide, broadest in the middle. Flowers are small, pale yellow, and jar shaped.	Gila, Maricopa, Yavapai	3,600-5,800 ft	Occurs on open slopes in chaparral or juniper grasslands. Prefers shallow, cobbled, and gravelly soils on steep slopes.	Arizona agave is a hybrid produced by a crossing of two other common agave species ( <i>A. chrysantha</i> x <i>A. toumeyana</i> ssp. <i>toumeyana</i> ).