

Table XXIII-4
PROSTATE CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence ^a			US Death ^b		
	All Races, Males	White Males	Black Males	All Races, Males	White Males	Black Males
<u>AGE-ADJUSTED RATES, 2001-2005</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	163.0	156.7	248.5	26.7	24.6	59.4
Under 65	57.7	55.3	102.7	1.9	1.6	5.2
65 and over	890.8	857.7	1,256.4	198.6	183.6	433.6
All ages(IARC world std) ^c	109.1	105.1	171.5	11.5	10.4	26.9
<u>AGE-SPECIFIC RATES, 2001-2005</u>						
AGE AT DIAGNOSIS/DEATH:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	-	-	-	-	-	-
35-39	0.8	0.7	1.6	-	-	-
40-44	8.7	7.4	21.9	0.2	0.2	0.5
45-49	40.2	36.3	88.4	0.9	0.7	2.4
50-54	136.4	129.7	255.4	3.3	2.7	9.7
55-59	328.2	314.8	591.4	9.1	7.6	25.2
60-64	573.9	556.4	945.0	23.5	19.9	64.8
65-69	877.4	854.2	1,320.4	50.1	43.7	127.9
70-74	983.7	957.1	1,373.9	98.7	87.4	253.3
75-79	970.8	929.2	1,335.0	183.3	166.5	439.2
80-84	816.3	765.0	1,081.2	324.3	301.1	722.5
85+	676.3	643.7	939.3	613.5	584.6	1,136.3

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.