


Partition of Trend in Incidence Rates for the Time Period 1999-2006 All Races, Both Sexes

Overall Decreasing Regression Coefficient : -4.43




Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).

Percents may not add to 100 due to rounding.

* Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Partition of Trend in Incidence Rates for the Time Period 1999-2006 All Races, Males

Overall Decreasing Regression Coefficient : -6.53




Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Percents may not add to 100 due to rounding.

Figure 1.27

Partition of Trend in Incidence Rates for the Time Period 1999-2006 All Races, Females

Overall Decreasing Regression Coefficient : -3.39



Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).

Percents may not add to 100 due to rounding.

* Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.