

## *Landscape Stability Indicators for Sagebrush Steppe Ecosystems*

by Bruce P. Van Haveren, Ph.D.,  
Senior Scientist, National Science  
and Technology Center

### Background

Sagebrush steppe ecosystems are found primarily within the Intermountain Semidesert Province (Bailey, 1995). It is nearly impossible to find examples of pristine sagebrush steppe. Most of these areas are small relict areas, topographically isolated from human disturbance or protected within special-use areas. West (2000) suggests that none of the original pre-Euroamerican sagebrush steppe remains. Only about one percent of the sagebrush steppe is in late seral stage, consisting of sagebrush-grass communities dominated by sagebrush with a healthy understory of grasses and forbs (Laycock 1991; West 2000). Another 25 to 30 percent of the original sagebrush steppe is in a stagnant late seral stage, close to a threshold of site change. This stage is characterized by relatively high densities of sagebrush (canopy cover >25%), large areas of bare soil in the shrub interspaces, and decreased grass and forb composition. West (2000) claims that most of the land area of the sagebrush steppe has experienced threshold-exceeding changes. Once a threshold is crossed, it becomes very difficult and expensive to move a sagebrush steppe community back across the threshold to a late seral stage. Indicators are needed that will help land managers identify these thresholds.

### Discussion

An ecological indicator is a characteristic of the environment, both abiotic and biotic, that may provide quantitative information on the status of ecological systems. An indicator may be a single environmental variable or




Figure 1. Valley profile for North Draw, North Fork of Savery Creek, southcentral Wyoming


an aggregation of variables expressed as an index. The indicator concept is related to the concept of landscape sensitivity, the capacity of a landscape to resist or absorb impulses of change. Abiotic system stability, as applied to landscapes, is an example of a parameter that can be used to help quantify the ecological condition of wildlands. A recent National Science and Technology Center (NSTC) project identified abiotic indicators of upland landscapes in the Intermountain Semidesert Province, with an initial focus on sagebrush steppe ecosystems.

Success in managing and restoring upland components of landscapes hinges on the ability to read and react to the ecological clues, both biotic and abiotic, that such landscapes reveal. Ecologically responsible land managers seek early warning signals of change on the landscape. They want to identify land-use thresholds or endpoints of ecological condition, they want to know where a watershed or ecosystem lies with respect to those endpoints, and they wish to learn whether the ecological trend is moving towards or away from the endpoints. They need to know when a landscape, such as a headwater basin, is stable or close to a threshold of change or whether it is moving in a downward or upward ecological trend. An example of such an indicator for the sagebrush steppe is hydrologic cover. Hydrologic cover is defined as the sum of live vegetal cover and litter cover

(Sturges, 1986). It is very closely related, in an inverse manner, to bare ground (Van Haveren, 2000). It is also inversely related to runoff potential, in that a high hydrologic cover value would indicate low runoff potential. Based on numerous studies on a variety of plant communities around the western United States, we know that hydrologic cover must be at least 70 percent to reduce runoff from rainfall and to protect soils from erosion. It turns out that hydrologic cover is an excellent indicator of upland watershed stability for sagebrush steppe and is very easy to measure. A review of ground cover studies in sagebrush steppe revealed that litter consistently is the dominant cover component (Van Haveren, in press). This suggests that land managers should pay particular attention to the litter balance on sagebrush steppe rangelands.

Steep headwater basins in the sagebrush steppe are often close to a threshold of geomorphic instability. The presence of gullies in these basins is an indication that the threshold was crossed. The threshold of instability appears to be a function of valley slope, drainage area, and valley width (Patton and Schumm, 1975; Elliott, 1989).

My observations of headwater areas in sagebrush steppe suggest there are two principal valley-floor types that illustrate the importance of valley width to geomorphic instability. Type I valleys are V-shaped, as shown in Figure 1



**Figure 2.** Valley profile for Middle Draw, North Fork of Savery Creek, southcentral Wyoming

for a headwater basin in southcentral Wyoming. Valley floors are very narrow and hydrologically connected to the adjacent hillslopes. Type II valleys, illustrated in Figure 2, are characterized by an inflected-U shape and often have wide, swale-like bottoms. Because of the slope breaks, the valley floors are hydrologically disconnected from the hillslopes. Type I valleys tend to be more unstable than Type II valleys. This classification is similar to that proposed for alluvial valleys by Bisson and Montgomery (1996), who differentiated between confined and unconfined valleys.

## Conclusion

Landscape analysis and classification hold considerable promise as tools for land managers concerned with the integrity and health of ecosystems. It may be possible, within a landscape classification system, to identify landform units that are susceptible to ecological change, or close to a threshold of change, and to combine this information with indicators of ecological condition.

## Literature Cited

Bailey, Robert G. 1995. Description of the ecoregions of the United States. 2nd edn. Miscellaneous Publication 1391. Washington DC: USDA Forest Service. 108p.

Bisson, Peter A. and Montgomery, David R. 1996. Valley segments,

stream reaches, and channel units. Pp. 23-51 in *Methods in Stream Ecology*, Hauer, F. Richard and Lamberti, Gary A. (eds). Chapter 2. 1996. London: Academic Press.

Elliott, John G. 1989. A geomorphically determined valley-erosion threshold for reclaimed surface-mined drainage basins, northwestern Colorado. Pp. 419-429 in *Proceedings of the Symposium on Headwaters Hydrology*, Woessner, William W. and Potts, Donald F. (eds). Bethesda, MD: American Water Resources Association. 708p.

Laycock, W. A. 1991. Stable states and thresholds of range condition on North American rangelands: a viewpoint. *Journal of Range Management* 44(5):427-433.

Patton, Peter C. and Schumm, Stanley A. 1975. Gully erosion, northwestern Colorado: a threshold phenomenon. *Geology* 3:88-90.

Sturges, David L. 1986. Responses of vegetation and ground cover to spraying a high elevation, big sagebrush watershed with 2,4-D. *Journal of Range Management* 39(2):141-146.

Van Haveren, Bruce P. 2000. Ground cover dynamics in a sagebrush steppe community. *Resource Notes* 19:1-2.

Van Haveren, Bruce P. In press. Characteristics of litter cover at a high elevation sagebrush steppe site. Submitted to the *Journal of Soil and Water Conservation*.

West, Neil E. 2000. Synecology and disturbance regimes of sagebrush steppe ecosystems. Pp. 15-26 in *Proceedings: Sagebrush Steppe Ecosystems Symposium*, Entwistle, P. G.; DeBolt, A. M.; Kaltenecker, J. H. and Steenhof, K. (comp.). Publication No. BLM/ID/PT-001001+1150. Boise, Idaho: Bureau of Land Management. Idaho State Office. 145p.

## Contact

Bruce P. Van Haveren, Ph.D.,  
Science and Research Advisor, BLM  
NSTC, ST-131  
Building 50, Denver Federal Center  
P.O. Box 25047  
Denver, CO 80225-0047  
phone (303) 236-0161  
fax (303) 236-6450  
email [bvanhave@blm.gov](mailto:bvanhave@blm.gov)

RESOURCE NOTES are intended to be early announcements of technical and informational topics for Bureau of Land Management personnel and some of their customers. Information in this RESOURCE NOTE is based on the opinion and experience of the author and has not been peer-reviewed. Conclusions and opinions expressed herein do not necessarily represent those of BLM. Use of trade names does not imply U.S. Government endorsement of commercial products.

If you have received a copy of or found out about RESOURCE NOTES in an indirect way and would like to be included in future mailings, please send the following:

NAME, TITLE, MAILING ADDRESS and a list of the two or three subject areas that you are most interested in or that most directly relate to your job. Send this information to Phil Dittbener, BLM, RS-140, P.O. Box 25047, Denver, CO. 80225-0047 or [phil\\_dittbener@blm.gov](mailto:phil_dittbener@blm.gov) or FAX 303-236-3508.

If you would like to prepare a RESOURCE NOTE for distribution, or you have an idea and author in mind for a good RESOURCE NOTE, please contact Phil Dittbener at 303-236-1833, FAX 303-236-3508 or [phil\\_dittbener@blm.gov](mailto:phil_dittbener@blm.gov) with the topic and the name of writer, including an address, phone number, and e-mail address.

Thank you for your interest in RESOURCE NOTES.



National Science & Technology Center



Bureau of Land Management