

## Guidance on Fenofibrate

This guidance represents the Food and Drug Administration's (FDA's) current thinking on this topic. It does not create or confer any rights for or on any person and does not operate to bind FDA or the public. You can use an alternative approach if the approach satisfies the requirements of the applicable statutes and regulations. If you want to discuss an alternative approach, contact the Office of Generic Drugs.

**Active ingredient:** Fenofibrate

**Form/Route:** Tablets/Oral

**Recommended studies:** 2 studies

1. Type of study: Fasting  
Design: Single-dose, two-way crossover *in-vivo*  
Strength: 145 mg  
Subjects: Normal healthy males and females, general population  
Additional Comments:

---

2. Type of study: Fed  
Design: Single-dose, two-way crossover *in-vivo*  
Strength: 145 mg  
Subjects: Normal healthy males and females, general population  
Additional comments:

---

**Analytes to measure:** Due to the difficulties with the fenofibrate assay, only the metabolite, fenofibric acid, should be measured.

**Bioequivalence based on (90% CI):** Fenofibric Acid

**Waiver request of in-vivo testing:** 48 mg based on (i) acceptable bioequivalence studies on the 145 mg strength, (ii) proportional similarity of the formulations 48 mg and 145 mg strengths, and (iii) acceptable in vitro dissolution testing of 48 mg and 145 mg strengths.

**Dissolution test method and sampling times:**

Please note that a **Dissolution Methods Database** is available to the public at the OGD website at <http://www.fda.gov/cder/ogd/index.htm>. Please find the dissolution information for this product at this website. Please conduct comparative dissolution testing on 12 dosage units each of all strengths of the test and reference products. Specifications will be determined upon review of the application.