

Neuregulin 1 transcripts are differentially expressed in schizophrenia and regulated by 5' SNPs associated with the disease

Amanda J. Law^{*†}, Barbara K. Lipska[‡], Cynthia Shannon Weickert[‡], Thomas M. Hyde[‡], Richard E. Straub[‡], Ryota Hashimoto[‡], Paul J. Harrison^{*}, Joel E. Kleinman[‡], and Daniel R. Weinberger[‡]

^{*}Department of Psychiatry, University of Oxford, Warneford Hospital, Oxford OX3 7JX, United Kingdom; and [‡]Clinical Brain Disorders Branch, Intramural Research Program, National Institute of Mental Health, National Institutes of Health, Bethesda, MD 20892-1385

Communicated by Gerald D. Fischbach, Columbia University College of Physicians and Surgeons, New York, NY, March 13, 2006 (received for review September 16, 2005)

Genetic variation in neuregulin 1 (*NRG1*) is associated with schizophrenia. The disease-associated SNPs are noncoding, and their functional implications remain unknown. We hypothesized that differential expression of the *NRG1* gene explains its association to the disease. We examined four of the disease-associated SNPs that make up the original risk haplotype in the 5' upstream region of the gene for their effects on mRNA abundance of *NRG1* types I–IV in human postmortem hippocampus. Diagnostic comparisons revealed a 34% increase in type I mRNA in schizophrenia and an interaction of diagnosis and genotype (SNP8NRG221132) on this transcript. Of potentially greater interest, a single SNP within the risk haplotype (SNP8NRG243177) and a 22-kb block of this core haplotype are associated with mRNA expression for the novel type IV isoform in patients and controls. Bioinformatic promoter analyses indicate that both SNPs lead to a gain/loss of putative binding sites for three transcription factors, serum response factor, myelin transcription factor-1, and High Mobility Group Box Protein-1. These data implicate variation in isoform expression as a molecular mechanism for the genetic association of *NRG1* with schizophrenia.

genetics | mRNA | human postmortem brain | hippocampus | ErbB

Schizophrenia is a complex, heritable psychiatric disorder. Recently, several putative schizophrenia susceptibility genes have been identified (1). Genomewide linkage studies and metaanalyses of linkage scans have highlighted chromosome 8p as a susceptibility locus (2–8). Extensive fine-mapping of the 8p locus, haplotype-association analysis, and linkage disequilibrium (LD) tests subsequently implicated neuregulin 1 (*NRG1*) (6), a gene with pleiotropic roles in neurodevelopment and plasticity (9).

The *NRG1* gene spans 1.2 Mb (6) and gives rise to many structurally and functionally distinct isoforms, through alternative promoter usage. These isoforms are divided into three classic groups (9): type I (previously known as acetylcholine receptor inducing activity, heregulin, or neu differentiation factor), type II (glia growth factor) and type III (cysteine-rich domain containing), which are based on distinct amino termini. All isoforms have a bioactive EGF-like domain that is responsible for activation of ErbB receptor tyrosine kinases (ErbB2–ErbB4). Additional *NRG1* 5' exons have recently been identified, giving rise putatively to novel *NRG1* types IV–VI in the human brain (10). No biological information is available presently for these novel isoforms.

In the original report of association with schizophrenia in an Icelandic population, Stefansson and colleagues (6) identified a “core at-risk haplotype” consisting of five SNPs (SNP8NRG221132, SNP8NRG221533, SNP8NRG241930, SNP8NRG243177, and SNP8NRG433E1006) and two microsatellites covering the 5' end of the *NRG1* gene and extending into the second intron (hereafter referred to as the “deCODE haplotype”). Separate follow-up studies in Scottish, Irish, mixed United Kingdom, and Dutch populations confirmed the genetic association between schizophrenia and *NRG1* by using markers within the

same core haplotype (11–14) or with overlapping markers in the 5' region (15, 16). Studies in four Asian populations also showed a strong association between schizophrenia and *NRG1* polymorphisms at the 5' (17–20) and 3' end of the gene (19). Together these results, not withstanding two negative studies (21, 22), provide strong evidence that *NRG1* is a schizophrenia-susceptibility gene. Additional support for *NRG1*'s role in schizophrenia comes from the phenotype of *NRG1* and ErbB4 mutant mice (6, 23–25), which exhibit behaviors similar to those of established rodent models of schizophrenia (26).

Exactly how genetic variation in *NRG1* impacts on disease susceptibility remains uncertain because the SNPs associated with schizophrenia are all noncoding, being either intronic, synonymous exonic substitutions, or upstream of the transcription start site. It is possible that an as-yet unknown (rare) coding mutation(s) exists, but it is more probable that the noncoding SNPs themselves, or other SNPs with which they are in LD, are functionally associated with the disease. One plausible explanation is that the *NRG1* SNPs are regulatory and affect disease susceptibility by altering expression (via altering transcriptional activity, alternative splicing, or stability of the RNA) and, thereby, the amount or distribution of the protein and ultimately its function. Support for this hypothesis comes from data demonstrating increased *NRG1* type I mRNA in the prefrontal cortex in schizophrenia (27) and from altered gene expression and splicing being associated with polymorphic variation in other brain diseases (1, 28, 29).

The aim of this study was to address whether disease-associated polymorphic loci in the 5' upstream region of *NRG1* modulate *NRG1* mRNA expression and contribute to its association with schizophrenia. We performed a series of hypothesis-driven experiments aimed at confirming previously published positive and negative expression data in schizophrenia (i.e., elevated type I, *NRG1*, and no change in type II or III (27)) and to test specifically the hypothesis that disease-associated SNPs in the original risk haplotype influence expression of the novel type IV isoform based on their physical proximity to its 5' regulatory region.

To test for differential expression of *NRG1* in schizophrenia, we examined mRNA abundance for *NRG1* types I–IV in the human hippocampus, a region prominently implicated in the pathogenesis of schizophrenia (30) and in the neurobiology of *NRG1* (31). Examination of the effects of genetic variation on *NRG1* expression included 4 SNPs from the deCODE core haplotype (6), with each SNP being tested individually for associations with *NRG1* mRNA levels in patients and controls. LD between SNPs was examined, and the deCODE at-risk haplotype region was tested for association

Conflict of interest statement: No conflicts declared.

Abbreviations: htSNP, haplotype-tagging SNP; LD, linkage disequilibrium; *NRG1*, neuregulin 1; PMI, postmortem interval; SRF, serum response factor.

[†]To whom correspondence should be addressed. E-mail: amanda.law@psych.ox.ac.uk.

© 2006 by The National Academy of Sciences of the USA




Fig. 1. *NRG1* types I–IV mRNA expression in the hippocampus of schizophrenic patients and control individuals. Quantitative RT-PCR analysis of *NRG1* types I–IV normalized to PBGD. (A–C) $n = 53$ normal control subjects (NC) and 38 patients with schizophrenia (SZ). (D) $n = 46$ normal controls subjects and 28 patients with schizophrenia. Box represents the proportion of the distribution falling between the 25th and 75th percentiles. Bars outside the box represent the SD. Bar inside represents the mean. Significant differences were found between controls and patients for type I *NRG1* expression. **, significant differences ($P < 0.01$)

with *NRG1* mRNA abundance. Our results support our primary hypotheses and indicate that the region of the gene implicated by the core at-risk haplotype impacts on specific *NRG1* isoforms and interacts with their expression in schizophrenia.

Results

NRG1 Isoform mRNA Expression in Schizophrenia. Normalized hippocampal *NRG1* type I mRNA expression levels were increased by 34% in schizophrenic patients compared with control subjects (Fig. 1; $F(1, 85) = 8.65$; $P = 0.004$). Similar significant findings were observed in the less well matched full cohort (84 controls vs. 44 schizophrenics; data not shown). No significant differences were observed between groups for the other *NRG1* isoforms {type II [$F(1, 85) = 0.10$; $P = 0.74$]; type III, [$F(1, 85) = 0.36$; $P = 0.54$]; type IV, [$F(1, 68) = 1.6$; $P = 0.20$]; 17 individuals were not available for the type IV study because of a shortage of RNA}. Expression ratios for all *NRG1* isoforms were calculated to investigate relative expression abnormalities, given previous reports of altered isoform ratios in the dorsolateral prefrontal cortex in schizophrenia (27). Type I *NRG1* mRNA expression was increased relative to all other isoforms in schizophrenia: type I/II [$F(1, 85) = 5.20$; $P = 0.02$], I/III [$F(1, 85) = 5.0$; $P = 0.02$], and I/IV [$F(1, 68) = 10.11$; $P = 0.002$]. No changes in any other isoform ratio were seen. No significant differences were observed between diagnostic groups for any of the housekeeping genes *PBGD*, [$F(1, 85) = 0.27$; $P = 0.6$]; *TBP*, [$F(1, 85) = 0.16$; $P = 0.21$], and *GUSB*, [$F(1, 68) = 0.001$; $P = 0.98$]. Covariation for pH, postmortem interval (PMI), and age was used in all analyses (see *Supporting Text*, which is published as supporting information on the PNAS web site).

Effects of 5' SNPs on *NRG1* Expression. The effect of SNPs on *NRG1* type I–III isoform expression were examined in the whole cohort ($n = 84$ controls and $n = 44$ schizophrenics). For type IV analysis, 24 individuals were not available for analysis because of a shortage of RNA (leaving $n = 74$ controls and $n = 30$ schizophrenics). None of the SNPs examined showed any effect on expression of type II or type III isoforms.




Fig. 2. Association between SNP8NRG122132 and type I *NRG1* mRNA in normal controls (NC) and schizophrenics (SZ). A significant interaction of genotype and diagnosis was observed on normalized type I mRNA expression ($P < 0.003$). Post hoc t tests revealed a main effect of genotype on type I *NRG1* expression in controls, with A allele carriers having increased levels compared with homozygous G individuals ($P < 0.003$). Schizophrenic patients homozygous for the G allele had higher levels of type I mRNA compared with controls with the same genotype ($P < 0.001$). Two individuals were excluded from analysis because of genotyping failure. Box represents the proportion of the distribution falling between the 25th and 75th percentiles. Bars outside the box represent the SD. The bar inside represents the mean.

Genetic Variation and Type I *NRG1* mRNA. A genotype \times diagnosis interaction [$F(3, 122) = 9.21$; $P = 0.003$] was found for SNP8NRG221132 and type I *NRG1* mRNA. There was no main effect of genotype. Post hoc comparisons showed that the genotype was significant in the controls alone (post hoc t test; $P = 0.003$; Fig. 2) with individuals carrying the rare A (2) allele expressing higher levels than individuals homozygous for G (1/1 genotype). The G allele constitutes the risk allele in the deCODE haplotype (6, 11). However, this pattern was not seen in schizophrenic patients who tended in the opposite direction (Fig. 2). Schizophrenic patients homozygous for the risk allele (G) had increased expression of type I *NRG1* mRNA compared with control subjects homozygous for the same allele (t test; $P = 0.001$). To confirm these findings, we genotyped SNP8NRG221132 in brain tissue from an earlier study of a separate cohort, in whom *NRG1* mRNA expression for types I–III had been measured by identical quantitative RT-PCR methods (controls $n = 13$, schizophrenics $n = 16$; $n = 22$ African Americans and $n = 7$ Caucasians) (27), and *NRG1* type I expression was found to be increased in the dorsolateral prefrontal cortex of these patients with schizophrenia. In this cohort we observed a main effect of SNP8NRG221132 genotype in the whole sample [$F(3, 25) = 15.17$; $P = 0.0005$] on prefrontal type I *NRG1* mRNA, with subjects carrying the A allele ($n = 4$) again showing higher type I *NRG1* levels compared with homozygous G cases ($n = 25$, data not shown). A genotype \times diagnosis interaction also was found in this cohort [$F(3, 25) = 16.26$; $P = 0.0005$], with schizophrenic patients homozygous for the G allele ($n = 14$) having greater expression of type I *NRG1* mRNA than control subjects ($n = 11$) with the same genotype ($P = 0.02$). No other SNPs examined in the study were associated with type I *NRG1* expression.

Genetic Variation and Type IV *NRG1* mRNA. Because $>10\%$ of individuals in the cohort were homozygous for the rare risk allele (T) at SNP8NRG243177, a complete analysis was conducted based on the three genotype groups (C/C, C/T, and T/T). We found a main effect of genotype for SNP8NRG243177 on type IV *NRG1* mRNA abundance in the whole sample [$F(5, 98) = 3.15$; $P = 0.04$; Fig. 3A]. The data suggest an allele dose effect with individuals heterozygous for the (T) risk allele (6, 11) having 21% more type IV *NRG1* mRNA than homozygous C/C individuals and individuals homozygous for the risk allele having 49% more type IV *NRG1*




Fig. 3. Association between SNP8NRG-243177 and type IV *NRG1* mRNA expression. (A) In the whole cohort, a main effect of genotype was observed (ANOVA; $P = 0.04$). An allele dose-dependent effect is suggested, with individuals homozygous for the risk allele having the highest levels of type IV *NRG1* mRNA ($P = 0.05$). (B) Data parsed by diagnosis. No genotype \times diagnosis interaction was observed.

mRNA expression than homozygous C/C individuals. However, standard post hoc contrasts between the three groups revealed only the two homozygote groups to be significantly different in terms of type IV abundance (post hoc t test; $P = 0.05$; Fig. 3A). This effect appeared more pronounced in the schizophrenia group alone (Fig. 3B), although no hint of a diagnosis \times genotype interaction was observed. Of note, normal control individuals homozygous for the T allele also showed the relatively greatest type IV *NRG1* mRNA expression (Fig. 3B).

Significant diagnosis \times genotype interactions were found between five of the six haplotype-tagging SNPs (htSNPs) and type IV mRNA abundance (rs4268090, rs4298458, rs4452759, rs4733263, and rs4476964) [range $F(3, 100) = 3.35\text{--}5.82$; $P = 0.033\text{--}0.018$]. A trend for a main effect of genotype was observed for the htSNP, rs4268090 [$F(3, 100) = 3.55$; $P = 0.06$]. Carrying the T (2) allele at this SNP was associated with higher levels of type IV mRNA compared with subjects homozygous for the C (1/1) allele.

Haplotype Analysis. Results of LD tests between pairs of all 10 5' SNPs in African American and Caucasian individuals can be found in Tables 1 and 2, which are published as supporting material on the PNAS web site. The four markers chosen from the deCODE haplotype were in significant, but moderate, LD in both groups (see *Supporting Text*). The frequencies for the four common haplotypes comprised of the four deCODE SNPs are shown in Table 3, which is published as supporting information on the PNAS web site). Hap 4 contains the specific alleles that form part of the deCODE haplotype.

To test whether this four SNP risk haplotype (hap4) was associated with *NRG1* mRNA levels, we used SNPHAP (www-gene.cimr.cam.ac.uk/clayton/software) to assign a diplotype (haplotype pair) to each individual. We then compared hap4 carriers

(diplotypes hap1/hap4, hap2/hap4, hap3/hap4, and hap4/hap4) to non-hap4 individuals (diplotypes hap1/hap1, hap1/hap2, hap1/hap3, hap2/hap2, hap3/hap3, and hap2/hap3), testing for an effect of carrying the risk haplotype on *NRG1* mRNA abundance. ANOVA revealed a main effect of hap4 on type IV mRNA abundance in the entire sample [$F(3, 89) = 3.38$; $P = 0.04$]. Hap4 carriers had 27% more type IV mRNA compared with non-hap4 individuals (Fig. 4A). This effect appeared more pronounced in the schizophrenic patients, where a 53% increase in type IV *NRG1* mRNA was seen in hap4 carriers compared to noncarriers; in controls, only a 14% increase was observed (Fig. 4B). However, no diagnosis by genotype interaction was found. Race was not included as a factor in the analysis because of the small number of African American individuals carrying hap4. Hap4 showed no effect on the expression of any of the other *NRG1* isoforms.

Promoter Analysis Based on Transcription Factor Binding Sites. As a further exploration of the functional relevance of disease-associated SNPs in the *NRG1* gene, we performed an analysis of putative transcription factor binding sites by using MATINSPECTOR software (Genomatix, Munich), a computational suite for promoter informatics. Two SNPs in the haplotype were indicated to be contained within transcriptional regulatory elements; notably, these elements were the two SNPs that, as described above, were found to impact upon expression of *NRG1* isoforms, SNP8NRG221132, which is associated with type I *NRG1*, and SNP8NRG243177, which is associated with type IV *NRG1*. SNP8NRG221132 is within a predicted transcription factor binding domain for serum response factor (SRF), with the risk allele (G) abolishing SRF binding. SNP8NRG243177 is also within a putative binding site for SRF and for myelin transcription factor 1. Carrying the risk allele (T) results in a predicted loss of binding to both of these transcription factors




Fig. 4. Association between diplotypes containing the deCODE risk haplotype (hap4) and type IV *NRG1* mRNA. Individuals were divided according to diplotype into two groups, non-hap4 carriers (haplotypes 1/1, 1/2, 1/3, 2/2, 3/3, and 2/3) and hap4 carriers (haplotypes 1/4, 2/4, 3/4, and 4/4). A main effect of diplotype was observed on normalized type IV *NRG1* mRNA levels (ANOVA; $P = 0.04$). (A) Individuals carrying the hap4 risk haplotype had increased levels compared with individuals not carrying hap4. (B) Effect of diplotype on type IV *NRG1* mRNA levels in controls and patients. Eleven individuals were not included in the diplotype analysis because of either the failure of genotyping at one or more of the SNPs or low probability ($<93\%$) of diplotype assignment according to SNPHAP.

and the acquisition of the transcription factor binding site for High Mobility Group Box Protein-1. None of the other 8 SNPs genotyped in the study mapped to transcription factor binding domains.

Analysis of Negative SNP Controls. The two negative control SNPs (rs10954867 and rs7005288) showed no association with any *NRG1* isoform in either controls or schizophrenic patients (all $P > 0.2$).

Regional Distribution of Hippocampal *NRG1* mRNA in Schizophrenia. Because no information is available regarding the distribution of *NRG1* in the human hippocampus, and this data was not provided from the quantitative RT-PCR experiments, we examined *NRG1* mRNA in the hippocampus in schizophrenia by using *in situ* hybridization with a “*pan*” *NRG1* probe (*Supporting Text*; see also Fig. 5, which is published as supporting information on the PNAS web site). No differences were seen in the distribution of *pan* *NRG1* mRNA between subfields or its overall abundance in schizophrenia.

Discussion

We have investigated the expression of *NRG1* type I–IV mRNA in the human hippocampus and examined the effects of schizophrenia and disease-associated polymorphisms in the 5′ upstream region on expression of these transcripts. We hypothesized that the genetic association of *NRG1* with schizophrenia is mediated by altered expression of the gene based on the location and noncoding nature of the disease-associated polymorphisms and the fact that extensive sequencing of *NRG1* has failed to identify pathogenic coding mutations (6). We report three principal findings: (i) up-regulation of type I expression in the hippocampus in schizophrenia, (ii) association of type I expression with a single SNP residing in the original deCODE risk haplotype, and (iii) association of type IV expression with a single SNP and a four-marker haplotype representing the 5′ upstream region of the original at-risk haplotype associated with schizophrenia. We provide evidence of association between disease linked-variation in *NRG1* and altered *NRG1* isoform expression in the brain, and we propose that altered transcript regulation is a potential molecular mechanism behind the genetic association of *NRG1* with schizophrenia.

Our finding of increased type I mRNA *NRG1* expression in the hippocampus in schizophrenia replicates the finding in the dorso-lateral prefrontal cortex of a smaller and separate brain series (27). These findings suggest that enhanced type I expression is robust and found in two separate brain regions in schizophrenia. In addition, we also replicate the finding that type II and type III isoform expression is unaltered in schizophrenia, suggesting that these isoforms may not be directly relevant to the pathophysiology of the disease. However, we did observe increases in the relative abundance of type I to type II–IV, suggesting that the contribution of these latter isoforms to *NRG1* signaling in the hippocampus may be indirectly compromised in patients with schizophrenia. At present, it is unclear whether type I up-regulation in schizophrenia is primary or secondary to other abnormalities in *NRG1* isoform regulation or to other molecular changes associated with the disease.

When the four individual SNPs representing the 5′ region of the deCODE at-risk haplotype were tested for association with type I *NRG1* mRNA, no main effects of genotype were seen for any of the SNPs. A diagnosis × genotype interaction was observed at a single SNP, SNP8NRG221132, and post hoc tests showed an effect of genotype only in control subjects on type I mRNA abundance. In a second independent cohort of brains in which increased type I mRNA expression previously had been reported in the schizophrenia samples, we found a main effect of a genotype at this SNP in the entire sample and again a genotype by diagnosis interaction. This main effect of the genotype was not seen in the first cohort; however, we note that the main effect in the entire sample is driven primarily by the controls. The observations that the main effect of

the genotype was primarily in the controls, and that the four-marker risk haplotype had no effect on type I *NRG1* expression, raise the possibility that SNP8NRG221132 influences type I expression independent of its contribution to risk for schizophrenia. Additional support for the functional relevance of SNP8NRG221132 comes from the bioinformatic promoter analysis that predicts the risk allele (G) leads to a loss of binding for SRF. SRF is a transcription factor that regulates the expression of genes encoding cytoskeletal proteins, such as cofilin and actin (32), both of which have been linked directly to *NRG1*'s role in actin dynamics (33). The loss of SRF binding in controls homozygous for the risk allele, therefore, may be related to lower levels of type I mRNA transcription, as reported here. The direct functional consequences of this SNP for type I *NRG1* transcriptional control remain difficult to predict because the SNP resides 1 Mb upstream from the transcriptional start site of type I. However, SNP8NRG221132 could conceivably reside in a regulatory element of the gene, as is seen in other key developmental genes where genomic regions harboring cis-regulatory elements can be located as far as 1 Mb from the transcription unit (34).

The interrelationship between SNP8NRG221132, type I *NRG1* expression, and schizophrenia is somewhat more difficult to interpret, because in contrast to the effect seen in controls, we did not see a similar genotype effect in the patients. This issue is discussed in *Supporting Text, Discussion: Genetic Association and Type I Expression in Schizophrenia*.

In contrast to the type I finding, which is not manifestly related to genetic variation in *NRG1* associated with schizophrenia, the association between both SNP8NRG243177 and the four-marker at-risk haplotype with expression of a novel isoform of *NRG1*, type IV, suggests that we may have identified a genetic mechanism and a molecular phenotype underlying the involvement of *NRG1* in susceptibility for schizophrenia. The risk allele of SNP8NRG243177 and the deCODE haplotype predicted higher levels of type IV *NRG1* expression in our entire sample. Analysis of the three genotype groups for SNP8NRG243177 revealed that individuals homozygous for the risk allele had the highest levels of type IV expression, with evidence of an allele dose-dependant effect. This observation appeared more pronounced in the patients, although trends in the same direction were found in the normal controls and no diagnosis × genotype interaction was observed. SNP8NRG243177 is the most 3′ of the SNPs in the four-marker deCODE haplotype and is located ≈1.2 kb upstream of the transcriptional start of type IV. Because none of the other single SNPs in this haplotype were associated with type IV *NRG1* expression, our results suggest that SNP8NRG243177 is a functional polymorphic variant that regulates type IV *NRG1* mRNA levels or is in strong LD with a nearby functional mutation. Additional support for the functional relevance of SNP8NRG243177 for gene regulation comes from the bioinformatic prediction that this SNP determines a putative transcription factor binding domain for SRF, myelin transcription factor 1, and High Mobility Group Box Protein-1. Of note, SRF and myelin transcription factor 1 play critical roles in neuronal migration, synaptic plasticity, and oligodendrocyte proliferation and survival, respectively, providing a striking molecular convergence with current hypotheses regarding the neurobiology of schizophrenia and the potential role of *NRG1* (35). However, we do not know which, if any, of these changes in transcription factor binding sites might mediate the association between SNP8NRG243177 and type IV expression and schizophrenia. Of potential interest is High Mobility Group Box Protein-1, an abundant chromatin-binding protein, which acts as an architectural facilitator in transcription (36). In our sample, acquisition of two High Mobility Group Box Protein-1-binding motifs (i.e., homozygosity for the risk allele) was associated with significantly elevated type IV *NRG1* expression, whereas acquisition of one (i.e., heterozygosity for the risk allele) was not. This observation suggests (i) that this binding site may potentiate

Quantitative Real-Time RT-PCR. *NRG1* mRNA expression levels were measured by quantitative RT-PCR by using an ABI Prism 7900 sequence detection system with a 384-well format (Applied Biosystems) as described (see *Supporting Text*). Our primary data analysis is based on normalization of *NRG1* mRNAs to an endogenous control gene, because it accounts for variability in the initial concentration and quality of total RNA and in the conversion efficiency of the reverse transcription reaction (42). Optimal normalization of a target mRNA to an endogenous control gene requires that the two transcripts have similar expression levels and that the control gene expression levels do not differ between the comparison groups. Based on this finding, PBGD was considered the most reliable for normalization here. Similar results were, however, obtained with normalization to TBP or GUSB (data not shown).

NRG1 Genotype Determination. DNA was extracted from cerebellar tissue by using a standard protocol supplied by PUREGENE (Gentra Systems). We genotyped four SNPs from the deCODE core haplotype (SNP8NRG221132, SNP8NRG221533, SNP8NRG241930, and SNP8NRG243177) and selected six additional SNPs (rs10096573, rs4268090, rs4298458, rs4452759, rs4733263, and rs4476964; Table 5 and Fig. 6, which are published as supporting material on the PNAS web site) from HAPMAP (www.hapmap.org) based on designation of these as htSNPs by using HAPLOVIEW (www.broad.mit.edu/mpg/haploview). The additional markers define the common haplotypes in an LD block containing a 22-kb region upstream of the first exon in type IV *NRG1*, which includes the four most 5' SNPs of the deCODE core risk haplotype. These six htSNPs were chosen to maximize genetic coverage because they are highly informative tags for the common haplotypes in this region of the gene.

Two SNPs at the 3' end of *NRG1* were selected from the dbSNP database as negative control genotypes (rs10954867 and rs7005288). These SNPs previously have not been associated with schizophrenia, are not in known regulatory domains, and were included in the analysis as a control for random statistical effects.

Genotyping was performed by using the Taqman 5' exonuclease allelic discrimination assay (details available upon request). Genotype reproducibility was routinely assessed by re-genotyping all samples for selected SNPs and was generally >99%. LD between 5' SNPs was determined by using the program LDMAX/GOLD (43). The program SNP-HAP written by David Clayton (version 1.0) was used to calculate haplotype frequencies and to assign diplotypes to individuals.

Statistical Analyses. Correlations of mRNA levels with demographic variables were performed for all subjects by using Spearman's correlations. Correlations of mRNA levels with neuroleptic medication (lifetime neuroleptic exposure, daily dose, and final neuroleptic dose) were investigated in the schizophrenic cohort. Primary planned comparisons between diagnostic groups were made by using univariate ANCOVA for each mRNA with diagnosis as the independent variable and age, pH, and PMI as covariates. Effects of genetic variation on *NRG1* mRNA expression were examined by using ANOVA with genotype and diagnosis as independent factors. Primary comparisons examined the effects of four SNPs and the core haplotype on type I–IV expression in patients and controls. Secondary, post hoc analyses included examination of the 6 htSNPs, where warranted. Where there was a significant genotype \times diagnosis interaction, individual group post hoc tests were examined as part of the standard ANOVA readout. Analysis of the effects of race was restricted to African American and Caucasian individuals because of the small sample size in other ethnic groups. The genotype groups did not differ on any of the demographic variables, and no correlations were seen between *NRG1* isoform expression, age, pH, or PMI in the different genotype groups; therefore straight ANOVAs are reported for the effects of genotype, but the statistical results were not changed when covariates were included. To increase power for statistical analyses of SNPs with minor allele frequencies <10%, we grouped individuals heterozygous and homozygous for the rare allele. Examination of all three genotype groups was conducted when the minor allele frequency was >10%. All experiments were conducted blind to diagnosis.

- Harrison, P. J. & Weinberger, D. R. (2005) *Mol. Psychiatry* **10**, 40–68.
- Pulver, A. E., Lasseter, V. K., Kasch, L., Wolyniec, P., Nestadt, G., Blouin, J. L., Kimberland, M., Babb, R., Vourlis, S., Chen, H., et al. (1995) *Am. J. Med. Genet.* **60**, 252–260.
- Kendler, K. S., MacLean, C. J., O'Neill, F. A., Burke, J., Murphy, B., Duke, F., Shinkwin, R., Easter, S. M., Webb, B. T., Zhang, J., et al. (1996) *Am. J. Psychiatry* **153**, 1534–1540.
- Kendler, K. S., Myers, J. M., O'Neill, F. A., Martin, R., Murphy, B., MacLean, C. J., Walsh, D., & Straub, R. E. (2000) *Am. J. Psychiatry* **157**, 402–408.
- Blouin, J. L., Dombroski, B. A., Nath, S. K., Lasseter, V. K., Wolyniec, P. S., Nestadt, G., Thorngvist, M., Ullrich, G., McGrath, J., Kasch, L., et al. (1998) *Nat. Genet.* **20**, 70–73.
- Stefansson, H., Sigurdsson, E., Steinthorsdottir, V., Bjornsdottir, S., Sigmundsson, T., Ghosh, S., Brynjolfsson, J., Gunnarsdottir, S., Ivarsson, O., Chou, T. T., et al. (2002) *Am. J. Hum. Genet.* **71**, 877–892.
- Badner, J. A. & Gershon, E. S. (2002) *Mol. Psychiatry* **7**, 405–411.
- Lewis, C. M., Levinson, D. F., Wise, L. H., DeLisi, L. E., Straub, R. E., Hovatta, I., Williams, N. M., Schwab, S. G., Pulver, A. E., Faraone, S. V., et al. (2003) *Am. J. Hum. Genet.* **73**, 34–48.
- Falls, D. L. (2003) *J. Neurocytol.* **32**, 619–647.
- Steinthorsdottir, V., Stefansson, H., Ghosh, S., Birgisdottir, B., Bjornsdottir, S., Fasquel, A. C., Olafsson, O., Stefansson, K. & Gulcher, J. R. (2004) *Gene* **342**, 97–105.
- Stefansson, H., Sarginson, J., Kong, A., Yates, P., Steinthorsdottir, V., Gudfinnsson, E., Gunnarsdottir, S., Walker, N., Petursson, H., Crombie, C., et al. (2003) *Am. J. Hum. Genet.* **72**, 83–87.
- Williams, N. M., Preece, A., Spurlock, G., Norton, N., Williams, H. J., Zammit, S., O'Donovan, M. C. & Owen, M. J. (2003) *Mol. Psychiatry* **8**, 485–487.
- Bakker, S. C., Hoogendoorn, M. L., Seltén, J. P., Verdúijn, W., Pearson, P. L., Sinke, R. J. & Kahn, R. S. (2004) *Mol. Psychiatry* **9**, 1061–1063.
- Hall, D., Gogos, J. A., Karayiorgou, M. (2004) *Genes Brain Behav.* **3**, 240–248.
- Corvin, A. P., Morris, D. W., McGhee, K., Schwaiger, S., Scully, P., Quinn, J., Meagher, D., Clair, D. S., Waddington, J. L. & Gill, M. (2004) *Mol. Psychiatry* **9**, 208–213.
- Petryshen, T. L., Middleton, F. A., Kirby, A., Aldinger, K. A., Purcell, S., Tahl, A. R., Morley, C. P., McGann, L., Gentile, K. L., Rockwell, G. N., et al. (2005) *Mol. Psychiatry* **10**, 366–374.
- Yang, J. Z., Si, T. M., Ruan, Y., Ling, Y. S., Han, Y. H., Wang, X. L., Zhou, M., Zhang, H. Y., Kong, Q. M. & Liu, C. (2003) *Mol. Psychiatry* **8**, 706–709.
- Tang, J. X., Chen, W. Y., He, G., Zhou, J., Gu, N. F., Feng, G. Y. & He, L. (2004) *Mol. Psychiatry* **9**, 11–20.
- Li, T., Stefansson, H., Gudfinnsson, E., Cai, G., Liu, X., Murray, R. M., Steinthorsdottir, V., Januel, D., Gudnadottir, V. G., Petursson, H., Ingason, A., et al. (2004) *Mol. Psychiatry* **9**, 698–704.
- Zhao, X., Shi, Y., Tang, J., Tang, R., Yu, L., Gu, N., Feng, G., Zhu, S., Liu, H., Xing, Y., et al. (2004) *J. Med. Genet.* **41**, 31–34.
- Iwata, N., Suzuki, T., Ikeda, M., Kitajima, T., Yamanouchi, Y., Inada, T. & Ozaki, N. (2004) *Mol. Psychiatry* **9**, 126–127.
- Thiselton, D. L., Webb, B. T., Neale, B. M., Ribble, R. C., O'Neill, F. A., Walsh, D., Riley, B. P. & Kendler, K. S. (2004) *Mol. Psychiatry* **9**, 777–783.
- Gerlaci, R., Pisacane, P. & Erickson, S. (2000) *Behav. Brain Res.* **109**, 219–227.
- Bao, J., Wolpowitz, D., Role, L. W. & Talmage, D. A. (2003) *J. Cell Biol.* **161**, 1133–1141.
- Rimer, M., Barrett, D. W., Maldonado, M. A., Vock, V. M. & Gonzalez-Lima, F. (2005) *NeuroReport* **16**, 271–275.
- Lipska, B. K. (2004) *J. Psychiatry Neurosci.* **29**, 282–286.
- Hashimoto, R., Straub, R. E., Weickert, C. S., Hyde, T. M., Kleinman, J. E. & Weinberger, D. R. (2004) *Mol. Psychiatry* **9**, 299–307.
- Farrer, M., Chan, P., Chen, R., Tan, L., Lincoln, S., Hernandez, D., Forno, L., Gwinn-Hardy, K., Petrucelli, L., Hussey, J., et al. (2001) *Ann. Neurol.* **50**, 293–300.
- Bertram, L., Hiltunen, M., Parkinson, M., Ingelsson, M., Lange, C., Ramasamy, K., Mullin, K., Menon, R., Sampson, A. J., Hsiao, M. Y., et al. (2005) *N. Engl. J. Med.* **352**, 884–894.
- Harrison, P. J. (2004) *Psychopharmacology* **174**, 151–162.
- Law, A. J., Shannon, W. C., Hyde, T. M., Kleinman, J. E. & Harrison, P. J. (2004) *Neuroscience* **127**, 125–136.
- Alberti, S., Krause, S. M., Kretz, O., Philippart, U., Lemberger, T., Casanova, E., Wiebel, F. F., Schwarz, H., Frotscher, M., Schutz, G., et al. (2005) *Proc. Natl. Acad. Sci. USA* **102**, 6148–6153.
- Nagata-Ohashi, K., Ohta, Y., Goto, K., Chiba, S., Mori, R., Nishita, M., Ohashi, K., Kousaka, K., Iwamatsu, A., Niwa, R., et al. (2004) *J. Cell Biol.* **165**, 465–471.
- Kleinjan, D. A. & van Heyningen, V. (2005) *Am. J. Hum. Genet.* **76**, 8–32.
- Corfas, G., Roy, K. & Buxbaum, J. D. (2004) *Nat. Neurosci.* **7**, 575–580.
- Thomas, J. O. & Travers, A. A. (2001) *Trends Biochem. Sci.* **26**, 167–174.
- Cardon, L. R. & Abecasis, G. R. (2003) *Trends Genet.* **19**, 135–140.
- Bejerano, G., Pheasant, M., Makunin, I., Stephen, S., Kent, W. J., Mattick, J. S. & Haussler, D. (2004) *Science* **304**, 1321–1325.
- Hong, C. J., Huo, S. J., Liao, D. L., Lee, K., Wu, J. Y. & Tsai, S. J. (2004) *Neurosci. Lett.* **366**, 158–161.
- Harrison, P. J. (1999) *Brain* **122**, 593–624.
- Gu, Z., Jiang, Q., Fu, A. K., Ip, N. Y. & Yan, Z. (2005) *J. Neurosci.* **25**, 4974–4984.
- Burbach, G. J., Dehn, D., Del Turco, D. & Deller, T. (2003) *J. Neurosci. Methods* **131**, 83–91.
- Abecasis, G. R. & Cookson, W. O. (2000) *Bioinformatics* **16**, 182–183.