

$\phi(1680)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

$\phi(1680)$ MASS

e^+e^- PRODUCTION

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1680±20 OUR ESTIMATE				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1709±20±43		¹ AUBERT	08S BABR	10.6 $e^+e^- \rightarrow$ hadrons
1623±20	948	² AKHMETSHIN	03 CMD2	1.05–1.38 $e^+e^- \rightarrow K_L^0 K_S^0$
~ 1500		³ ACHASOV	98H RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$, $\omega\pi^+\pi^-$, K^+K^-
~ 1900		⁴ ACHASOV	98H RVUE	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1700±20		⁵ CLEGG	94 RVUE	$e^+e^- \rightarrow K^+K^-$, $K_S^0 K\pi$
1657±27	367	BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1655±17		⁶ BISELLO	88B DM2	$e^+e^- \rightarrow K^+K^-$
1680±10		⁷ BUON	82 DM1	$e^+e^- \rightarrow$ hadrons
1677±12		⁸ MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K\pi$

PHOTOPRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1753± 3	⁹ LINK	02K FOCS	20–160 $\gamma p \rightarrow K^+K^-p$
1726±22	⁹ BUSENITZ	89 TPS	$\gamma p \rightarrow K^+K^-X$
1760±20	⁹ ATKINSON	85C OMEG	20–70 $\gamma p \rightarrow K\bar{K}X$
1690±10	⁹ ASTON	81F OMEG	25–70 $\gamma p \rightarrow K^+K^-X$

$p\bar{p}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1700±8	¹⁰ AMSLER	06 CBAR	0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$

¹ From the simultaneous fit to the $K\bar{K}^*(892) + c.c.$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.

² From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known.

³ Using data from IVANOV 81, BARKOV 87, BISELLO 88B, DOLINSKY 91, and ANTONELLI 92.

⁴ Using the data from BISELLO 91C.

⁵ Using BISELLO 88B and MANE 82 data.

⁶ From global fit including ρ , ω , ϕ and $\rho(1700)$ assume mass 1570 MeV and width 510 MeV for ρ radial excitation.

⁷ From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_S^0 K_L^0$, $K_S^0 K^\pm \pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.

⁸ Fit to one channel only, neglecting interference with ω , $\rho(1700)$.

⁹ We list here a state decaying into K^+K^- possibly different from $\phi(1680)$.

¹⁰ Could also be $\rho(1700)$.

$\phi(1680)$ WIDTH

e^+e^- PRODUCTION

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

150±50 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

• • • We do not use the following data for averages, fits, limits, etc. • • •

322±77±160		¹¹ AUBERT	08S	BABR	10.6 $e^+e^- \rightarrow$ hadrons
139±60	948	¹² AKHMETSHIN	03	CMD2	1.05–1.38 $e^+e^- \rightarrow K_L^0 K_S^0$
300±60		¹³ CLEGG	94	RVUE	$e^+e^- \rightarrow K^+ K^-, K_S^0 K \pi$
146±55	367	BISELLO	91C	DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
207±45		¹⁴ BISELLO	88B	DM2	$e^+e^- \rightarrow K^+ K^-$
185±22		¹⁵ BUON	82	DM1	$e^+e^- \rightarrow$ hadrons
102±36		¹⁶ MANE	82	DM1	$e^+e^- \rightarrow K_S^0 K \pi$

PHOTOPRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

122±63	¹⁷ LINK	02K	FOCS	20–160 $\gamma p \rightarrow K^+ K^- p$
121±47	¹⁷ BUSENITZ	89	TPS	$\gamma p \rightarrow K^+ K^- X$
80±40	¹⁷ ATKINSON	85C	OMEG	20–70 $\gamma p \rightarrow K \bar{K} X$
100±40	¹⁷ ASTON	81F	OMEG	25–70 $\gamma p \rightarrow K^+ K^- X$

$p\bar{p}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

143±24	¹⁸ AMSLER	06	CBAR	0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$
--------	----------------------	----	------	--

¹¹ From the simultaneous fit to the $K \bar{K}^*(892) +$ c.c. and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.

¹² From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known.

¹³ Using BISELLO 88B and MANE 82 data.

¹⁴ From global fit including ρ , ω , ϕ and $\rho(1700)$

¹⁵ From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_S^0 K_L^0$, $K_S^0 K^\pm \pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.

¹⁶ Fit to one channel only, neglecting interference with ω , $\rho(1700)$.

¹⁷ We list here a state decaying into $K^+ K^-$ possibly different from $\phi(1680)$.

¹⁸ Could also be $\rho(1700)$.

$\phi(1680)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \bar{K}^*(892) +$ c.c.	dominant
Γ_2 $K_S^0 K \pi$	seen
Γ_3 $K \bar{K}$	seen
Γ_4 $K_L^0 K_S^0$	

Γ_5	$e^+ e^-$	seen
Γ_6	$\omega \pi \pi$	not seen
Γ_7	$\phi \eta$	
Γ_8	$K^+ K^- \pi^0$	

$\phi(1680) \Gamma(i)\Gamma(e^+ e^-)/\Gamma^2(\text{total})$

This combination of a branching ratio into channel (*i*) and branching ratio into $e^+ e^-$ is directly measured and obtained from the cross section at the peak. We list only data that have not been used to determine the branching ratio into (*i*) or $e^+ e^-$.

$$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \qquad \Gamma_4/\Gamma \times \Gamma_5/\Gamma$$

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.131 ± 0.059	948	¹⁹ AKHMETSHIN 03	CMD2	$1.05\text{--}1.38 e^+ e^- \rightarrow K_L^0 K_S^0$
-------------------	-----	-----------------------------	------	---

¹⁹ From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known. Recalculated by us.

$$\Gamma(K \bar{K}^*(892) + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \qquad \Gamma_1/\Gamma \times \Gamma_5/\Gamma$$

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.15 \pm 0.16 \pm 0.01$		²⁰ AUBERT 08S	BABR	$10.6 e^+ e^- \rightarrow K \bar{K}^*(892) \gamma +$
--------------------------	--	--------------------------	------	--

3.29 ± 1.57	367	²¹ BISELLO 91C	DM2	$1.35\text{--}2.40 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$
-----------------	-----	---------------------------	-----	---

²⁰ From the simultaneous fit to the $K \bar{K}^*(892) + \text{c.c.}$ and $\phi \eta$ data from AUBERT 08S using the results of AUBERT 07AK.

²¹ Recalculated by us with the published value of $B(K \bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+ e^-)$.

$$\Gamma(\phi \eta)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \qquad \Gamma_7/\Gamma \times \Gamma_5/\Gamma$$

<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.43 \pm 0.10 \pm 0.09$	²² AUBERT 08S	BABR	$10.6 e^+ e^- \rightarrow \phi \eta \gamma$
--------------------------	--------------------------	------	---

²² From the simultaneous fit to the $K \bar{K}^*(892) + \text{c.c.}$ and $\phi \eta$ data from AUBERT 08S using the results of AUBERT 07AK.

$\phi(1680)$ BRANCHING RATIOS

$$\Gamma(K \bar{K}^*(892) + \text{c.c.})/\Gamma(K_S^0 K \pi) \qquad \Gamma_1/\Gamma_2$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

dominant	MANE 82	DM1	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$
----------	---------	-----	---

$$\Gamma(K \bar{K})/\Gamma(K \bar{K}^*(892) + \text{c.c.}) \qquad \Gamma_3/\Gamma_1$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.07 ± 0.01	BUON 82	DM1	$e^+ e^-$
-----------------	---------	-----	-----------

$\Gamma(\omega\pi\pi)/\Gamma(K\bar{K}^*(892)+c.c.)$	Γ_6/Γ_1
VALUE	DOCUMENT ID TECN COMMENT
<0.10	BUON 82 DM1 e^+e^-

$\Gamma(\phi\eta)/\Gamma(K\bar{K}^*(892)+c.c.)$	Γ_7/Γ_1
VALUE	DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

≈ 0.37 ²³ AUBERT 08S BABR 10.6 $e^+e^- \rightarrow$ hadrons

²³ From the fit including data from AUBERT 07AK.

$\phi(1680)$ REFERENCES

AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)
AKHMETSHIN	03	PL B551 27	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
Also		PAN 65 1222	E.V. Anashkin, V.M. Aulchenko, R.R. Akhmetshin	
		Translated from YAF 65 1255.		
LINK	02K	PL B545 50	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BUSENITZ	89	PR D40 1	J.K. Busenitz <i>et al.</i>	(ILL, FNAL)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from ZETFP 46 132.		
ATKINSON	85C	ZPHY C27 233	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
BUON	82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)
MANE	82	PL 112B 178	F. Mane <i>et al.</i>	(LALO)
ASTON	81F	PL 104B 231	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
MANE	81	PL 99B 261	F. Mane <i>et al.</i>	(ORSAY)

OTHER RELATED PAPERS

ACHASOV	07A	PR D76 072012	M.N. Achasov <i>et al.</i>	(SND Collab.)
ACHASOV	06D	JETP 103 720	N.N. Achasov <i>et al.</i>	(SND Collab.)
		Translated from ZETF 130 831.		
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
CLOSE	02	PR D65 092003	F.E. Close, A. Donnachie, Yu.S. Kalashnikova	
LINK	02K	PL B545 50	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ACHASOV	97F	PAN 60 2029	N.N. Achasov, A.A. Kozhevnikov	(NOVM)
		Translated from YAF 60 2212.		
ATKINSON	86C	ZPHY C30 541	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	84	NP B231 15	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	84B	NP B231 1	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	83C	NP B229 269	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)
MANE	81	PL 99B 261	F. Mane <i>et al.</i>	(ORSAY)
ASTON	80F	NP B174 269	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)