

$\pi_2(1670)$

$$I^G(J^{PC}) = 1^-(2^{-+})$$

$\pi_2(1670)$ MASS

| <u>VALUE (MeV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u> |
|---|-------------|---|-------------|------------|---|
| 1672.4 ± 3.2 OUR AVERAGE | | Error includes scale factor of 1.4. See the ideogram below. | | | |
| 1749 ± 10 ± 100 | 145k | LU | 05 | B852 | 18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$ |
| 1676 ± 3 ± 8 | | 1 CHUNG | 02 | B852 | 18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$ |
| 1685 ± 10 ± 30 | | 2 BARBERIS | 01 | | 450 $p p \rightarrow p_f 3\pi^0 p_s$ |
| 1687 ± 9 ± 15 | | AMELIN | 99 | VES | 37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$ |
| 1669 ± 4 | | BARBERIS | 98B | | 450 $p p \rightarrow p_f \rho \pi p_s$ |
| 1670 ± 4 | | BARBERIS | 98B | | 450 $p p \rightarrow p_f f_2(1270) \pi p_s$ |
| 1730 ± 20 | | 3 AMELIN | 95B | VES | 36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$ |
| 1690 ± 14 | | 4 BERDNIKOV | 94 | VES | 37 $\pi^- A \rightarrow K^+ K^- \pi^- A$ |
| 1710 ± 20 | 700 | ANTIPOV | 87 | SIGM - | 50 $\pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$ |
| 1676 ± 6 | | 4 EVANGELIS... | 81 | OMEG - | 12 $\pi^- p \rightarrow 3\pi p$ |
| 1657 ± 14 | | 4,5 DAUM | 80D | SPEC - | 63-94 $\pi p \rightarrow 3\pi X$ |
| 1662 ± 10 | 2000 | 4 BALTAY | 77 | HBC + | 15 $\pi^+ p \rightarrow p 3\pi$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| 1742 ± 31 ± 49 | | ANTREASYAN | 90 | CBAL | $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$ |
| 1624 ± 21 | | 1 BELLINI | 85 | SPEC | 40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$ |
| 1622 ± 35 | | 6 BELLINI | 85 | SPEC | 40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$ |
| 1693 ± 28 | | 7 BELLINI | 85 | SPEC | 40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$ |
| 1710 ± 20 | | 8 DAUM | 81B | SPEC - | 63,94 $\pi^- p$ |
| 1660 ± 10 | | 4 ASCOLI | 73 | HBC - | 5-25 $\pi^- p \rightarrow p \pi_2$ |

¹ From $f_2(1270)\pi$ decay.

² From a fit to the invariant mass distribution.

³ From a fit to $J^{PC} = 2^{-+} f_2(1270)\pi, f_0(1370)\pi$ waves.

⁴ From a fit to $J^P = 2^- S$ -wave $f_2(1270)\pi$ partial wave.


⁵ Clear phase rotation seen in $2^- S, 2^- P, 2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.

⁶ From $\rho\pi$ decay.

⁷ From $\sigma\pi$ decay.

⁸ From a two-resonance fit to four $2^- 0^+$ waves. This should not be averaged with all the single resonance fits.

WEIGHTED AVERAGE
 1672.4±3.2 (Error scaled by 1.4)




$\pi_2(1670)$ WIDTH

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | CHG | COMMENT | |
|---|------|---|------|------|--|--|
| 259± 9 OUR AVERAGE | | Error includes scale factor of 1.3. See the ideogram below. | | | | |
| 408± 60± 250 | 145k | LU | 05 | B852 | 18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$ | |
| 254± 3± 31 | | ⁹ CHUNG | 02 | B852 | 18.3 $\pi^- p \rightarrow$ $\pi^+ \pi^- \pi^- p$ | |
| 265± 30± 40 | | ¹⁰ BARBERIS | 01 | | 450 $pp \rightarrow p_f 3\pi^0 p_S$ | |
| 168± 43± 53 | | AMELIN | 99 | VES | 37 $\pi^- A \rightarrow$ $\omega \pi^- \pi^0 A^*$ | |
| 268± 15 | | BARBERIS | 98B | | 450 $pp \rightarrow p_f \rho \pi p_S$ | |
| 256± 15 | | BARBERIS | 98B | | 450 $pp \rightarrow$ $p_f f_2(1270) \pi p_S$ | |
| 310± 20 | | ¹¹ AMELIN | 95B | VES | 36 $\pi^- A \rightarrow$ $\pi^+ \pi^- \pi^- A$ | |
| 190± 50 | | ¹² BERDNIKOV | 94 | VES | 37 $\pi^- A \rightarrow$ $K^+ K^- \pi^- A$ | |
| 170± 80 | 700 | ANTIPOV | 87 | SIGM | - | 50 $\pi^- Cu \rightarrow$ $\mu^+ \mu^- \pi^- Cu$ |
| 260± 20 | | ¹² EVANGELIS... | 81 | OMEG | - | 12 $\pi^- p \rightarrow 3\pi p$ |
| 219± 20 | | ^{12,13} DAUM | 80D | SPEC | - | 63-94 $\pi p \rightarrow 3\pi X$ |
| 285± 60 | 2000 | ¹² BALTAY | 77 | HBC | + | 15 $\pi^+ p \rightarrow p 3\pi$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | | |
| 236± 49± 36 | | ANTREASYAN | 90 | CBAL | | $e^+ e^- \rightarrow$ $e^+ e^- \pi^0 \pi^0 \pi^0$ |
| 304± 22 | | ⁹ BELLINI | 85 | SPEC | | 40 $\pi^- A \rightarrow$ $\pi^- \pi^+ \pi^- A$ |
| 404± 108 | | ¹⁴ BELLINI | 85 | SPEC | | 40 $\pi^- A \rightarrow$ $\pi^- \pi^+ \pi^- A$ |

| | | | | |
|----------|-----------------------|-----|------|--|
| 330 ± 90 | ¹⁵ BELLINI | 85 | SPEC | 40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$ |
| 312 ± 50 | ¹⁶ DAUM | 81B | SPEC | — 63,94 $\pi^- p$ |
| 270 ± 60 | ¹² ASCOLI | 73 | HBC | — 5–25 $\pi^- p \rightarrow p \pi_2$ |

- ⁹ From $f_2(1270)\pi$ decay.
- ¹⁰ From a fit to the invariant mass distribution.
- ¹¹ From a fit to $J^{PC} = 2^- + f_2(1270)\pi, f_0(1370)\pi$ waves.
- ¹² From a fit to $J^P = 2^- f_2(1270)\pi$ partial wave.
- ¹³ Clear phase rotation seen in $2^- S, 2^- P, 2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.
- ¹⁴ From $\rho\pi$ decay.
- ¹⁵ From $\sigma\pi$ decay.
- ¹⁶ From a two-resonance fit to four $2^- 0^+$ waves. This should not be averaged with all the single resonance fits.



$\pi_2(1670)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Confidence level |
|--------------------------------|--------------------------------|------------------|
| Γ_1 3π | (95.8 ± 1.4) % | |
| Γ_2 $\pi^+ \pi^- \pi^0$ | | |
| Γ_3 $\pi^0 \pi^0 \pi^0$ | | |
| Γ_4 $f_2(1270)\pi$ | (56.3 ± 3.2) % | |
| Γ_5 $\rho\pi$ | (31 ± 4) % | |
| Γ_6 $\sigma\pi$ | (10.9 ± 3.4) % | |
| Γ_7 $(\pi\pi)_S$ -wave | (8.7 ± 3.4) % | |

| | | | |
|---------------|---------------------------------|------------------------|-------|
| Γ_8 | $K\bar{K}^*(892) + \text{c.c.}$ | $(4.2 \pm 1.4) \%$ | |
| Γ_9 | $\omega\rho$ | $(2.7 \pm 1.1) \%$ | |
| Γ_{10} | $\gamma\gamma$ | $< 2.8 \times 10^{-7}$ | 90% |
| Γ_{11} | $\eta\pi$ | | |
| Γ_{12} | $\pi^\pm 2\pi^+ 2\pi^-$ | | |
| Γ_{13} | $\rho(1450)\pi$ | $< 3.6 \times 10^{-3}$ | 97.7% |
| Γ_{14} | $b_1(1235)\pi$ | $< 1.9 \times 10^{-3}$ | 97.7% |
| Γ_{15} | $\eta 3\pi$ | | |
| Γ_{16} | $f_1(1285)\pi$ | possibly seen | |
| Γ_{17} | $a_2(1320)\pi$ | not seen | |

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 6 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 1.9$ for 3 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | |
|-------|-------|-------|-------|
| x_5 | -53 | | |
| x_7 | -29 | -59 | |
| x_8 | -8 | -21 | -9 |
| | x_4 | x_5 | x_7 |

$\pi_2(1670)$ PARTIAL WIDTHS

| $\Gamma(\gamma\gamma)$ | | | | | | Γ_{10} |
|---|-----|------------------------|------|------|--|--|
| VALUE (keV) | CL% | DOCUMENT ID | TECN | CHG | COMMENT | |
| <0.072 | 90 | ¹⁷ ACCIARRI | 97T | L3 | $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$ | |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | | |
| <0.19 | 90 | ¹⁷ ALBRECHT | 97B | ARG | $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$ | |
| 1.41 $\pm 0.23 \pm 0.28$ | | ANTREASYAN 90 | CBAL | 0 | $e^+e^- \rightarrow e^+e^-\pi^0\pi^0\pi^0$ | |
| 0.8 $\pm 0.3 \pm 0.12$ | | ¹⁸ BEHREND | 90C | CELL | 0 | $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$ |
| 1.3 $\pm 0.3 \pm 0.2$ | | ¹⁹ BEHREND | 90C | CELL | 0 | $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$ |

¹⁷ Decaying into $f_2(1270)\pi$ and $\rho\pi$.

¹⁸ Constructive interference between $f_2(1270)\pi, \rho\pi$ and background.

¹⁹ Incoherent Ansatz.

$\pi_2(1670) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \qquad \Gamma_2\Gamma_{10}/\Gamma$

| VALUE (keV) | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------|-----|-----------------------------|------|--|
| <0.1 | 95 | ²⁰ SCHEGELSKY 06 | RVUE | $\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$ |

²⁰ From analysis of L3 data at 183–209 GeV.

$\pi_2(1670)$ BRANCHING RATIOS

$\Gamma(3\pi)/\Gamma_{\text{total}} \qquad \Gamma_1/\Gamma = (\Gamma_4+\Gamma_5+\Gamma_7)/\Gamma$

| VALUE | DOCUMENT ID |
|----------------------------|-------------|
| 0.958±0.014 OUR FIT | |

$\Gamma(\pi^0\pi^0\pi^0)/\Gamma(\pi^+\pi^-\pi^0) \qquad \Gamma_3/\Gamma_2$

| VALUE | DOCUMENT ID | COMMENT |
|-----------------------|---------------------------|--------------------------------------|
| 0.29±0.03±0.05 | ²¹ BARBERIS 01 | 450 $p p \rightarrow p_f 3\pi^0 p_s$ |

$\Gamma(\rho\pi)/0.565\Gamma(f_2(1270)\pi) \qquad \Gamma_5/0.565\Gamma_4$

(With $f_2(1270) \rightarrow \pi^+\pi^-$.)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------------------------------|------|---|
| 0.97±0.09 OUR AVERAGE | Error includes scale factor of 1.9. | | |
| 0.76±0.07±0.10 | CHUNG 02 | B852 | 18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$ |
| 1.01±0.05 | BARBERIS 98B | | 450 $p p \rightarrow p_f \pi^+\pi^-\pi^0 p_s$ |

$\Gamma(\sigma\pi)/\Gamma(f_2(1270)\pi) \qquad \Gamma_6/\Gamma_4$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|---------------------------|------|--|
| 0.19±0.06 OUR AVERAGE | | | |
| 0.17±0.02±0.07 | CHUNG 02 | B852 | 18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$ |
| 0.24±0.10 | ^{22,23} BAKER 99 | SPEC | 1.94 $\bar{p} p \rightarrow 4\pi^0$ |

$\frac{1}{2}\Gamma(\rho\pi)/\Gamma(\pi^\pm\pi^+\pi^-) \qquad \frac{1}{2}\Gamma_5/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$

| VALUE | DOCUMENT ID | TECN | CHG | COMMENT |
|--------------------------|------------------------|------|-----|-----------------|
| 0.29±0.04 OUR FIT | | | | |
| 0.29±0.05 | ²⁴ DAUM 81B | SPEC | | 63,94 $\pi^- p$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|------------|-----|---|--------------------------------|
| <0.3 | BARTSCH 68 | HBC | + | 8 $\pi^+ p \rightarrow 3\pi p$ |
|------|------------|-----|---|--------------------------------|

$0.565\Gamma(f_2(1270)\pi)/\Gamma(\pi^\pm\pi^+\pi^-) \qquad 0.565\Gamma_4/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$

(With $f_2(1270) \rightarrow \pi^+\pi^-$.)

| VALUE | DOCUMENT ID | TECN | CHG | COMMENT |
|---|-------------------------------------|------|-----|----------------------------------|
| 0.604±0.035 OUR FIT | | | | |
| 0.60 ±0.05 OUR AVERAGE | Error includes scale factor of 1.3. | | | |
| 0.61 ±0.04 | ²⁴ DAUM 81B | SPEC | | 63,94 $\pi^- p$ |
| 0.76 ^{+0.24} _{-0.34} | ARMENISE 69 | DBC | + | 5.1 $\pi^+ d \rightarrow d 3\pi$ |
| 0.35 ±0.20 | BALTAY 68 | HBC | + | 7–8.5 $\pi^+ p$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.59 | BARTSCH 68 | HBC | + | 8 $\pi^+ p \rightarrow 3\pi p$ |

$$0.624\Gamma((\pi\pi)_{S\text{-wave}})/\Gamma(\pi^\pm\pi^+\pi^-) \quad 0.624\Gamma_7/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$$

(With $(\pi\pi)_{S\text{-wave}} \rightarrow \pi^+\pi^-$.)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|----------|-----------------|
| 0.10±0.04 OUR FIT | | | |
| 0.10±0.05 | 24 DAUM | 81B SPEC | 63,94 $\pi^- p$ |

$$\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma(f_2(1270)\pi) \quad \Gamma_8/\Gamma_4$$

| VALUE | DOCUMENT ID | TECN | CHG | COMMENT |
|----------------------------|--------------|----------|-----|--|
| 0.075±0.025 OUR FIT | | | | |
| 0.075±0.025 | 25 ARMSTRONG | 82B OMEG | - | 16 $\pi^- p \rightarrow K^+ K^- \pi^- p$ |

$$\Gamma(\omega\rho)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|--------|---|
| 0.027±0.004±0.010 | 26 AMELIN | 99 VES | 37 $\pi^- A \rightarrow \omega\pi^-\pi^0 A^*$ |

$$\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-) \quad \Gamma_{11}/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$$

(All η decays.)

| VALUE | DOCUMENT ID | TECN | CHG | COMMENT |
|-----------------|-------------|--------|-----|-----------------|
| <0.09 | BALTAY | 68 HBC | + | 7-8.5 $\pi^+ p$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----------|--------|---|------------------------------------|
| <0.10 | CRENNELL | 70 HBC | - | 6 $\pi^- p \rightarrow f_2\pi^- N$ |
|-------|----------|--------|---|------------------------------------|

$$\Gamma(\pi^\pm 2\pi^+ 2\pi^-)/\Gamma(\pi^\pm\pi^+\pi^-) \quad \Gamma_{12}/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$$

| VALUE | DOCUMENT ID | TECN | CHG | COMMENT |
|-----------------|-------------|--------|-----|------------------------------------|
| <0.10 | CRENNELL | 70 HBC | - | 6 $\pi^- p \rightarrow f_2\pi^- N$ |
| <0.1 | BALTAY | 68 HBC | + | 7,8.5 $\pi^+ p$ |

$$\Gamma(\rho(1450)\pi)/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------|------|-------------|--------|---|
| <0.0036 | 97.7 | AMELIN | 99 VES | 37 $\pi^- A \rightarrow \omega\pi^-\pi^0 A^*$ |

$$\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------|------|-------------|--------|---|
| <0.0019 | 97.7 | AMELIN | 99 VES | 37 $\pi^- A \rightarrow \omega\pi^-\pi^0 A^*$ |

$$\Gamma(f_1(1285)\pi)/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma$$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------|------|-------------|---------|--|
| possibly seen | 69k | KUHN | 04 B852 | 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$ |

$$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma$$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------|------|-------------|---------|--|
| not seen | 69k | KUHN | 04 B852 | 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$ |

D-wave/S-wave RATIO FOR $\pi_2(1670) \rightarrow f_2(1270)\pi$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|---|
| -0.18±0.06 | ²² BAKER | 99 | SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.22±0.10 | ²⁴ DAUM | 81B | SPEC 63,94 $\pi^- p$ |

F-wave/P-wave RATIO FOR $\pi_2(1670) \rightarrow \rho\pi$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|------|---|
| -0.72±0.07±0.14 | CHUNG | 02 | B852 18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$ |

²¹ Using BARBERIS 98B.

²² Using preliminary CBAR data.

²³ With the $\sigma\pi$ in $L=2$ and the $f_2(1270)\pi$ in $L=0$.

²⁴ From a two-resonance fit to four $2^- 0^+$ waves.

²⁵ From a partial-wave analysis of $K^+ K^- \pi^-$ system.

²⁶ Normalized to the $B(\pi_2(1670) \rightarrow f_2\pi)$.

$\pi_2(1670)$ REFERENCES

| | | | | |
|--------------|-----|------------------------|-------------------------------------|-----------------------------------|
| SCHEGELSKY | 06 | EPJ A27 199 | V.A. Schegelsky <i>et al.</i> | |
| LU | 05 | PRL 94 032002 | M. Lu <i>et al.</i> | (BNL E852 Collab.) |
| KUHN | 04 | PL B595 109 | J. Kuhn <i>et al.</i> | (BNL E852 Collab.) |
| CHUNG | 02 | PR D65 072001 | S.U. Chung <i>et al.</i> | (BNL E852 Collab.) |
| BARBERIS | 01 | PL B507 14 | D. Barberis <i>et al.</i> | |
| AMELIN | 99 | PAN 62 445 | D.V. Amelin <i>et al.</i> | (VES Collab.) |
| | | Translated from YAF 62 | 487. | |
| BAKER | 99 | PL B449 114 | C.A. Baker <i>et al.</i> | |
| BARBERIS | 98B | PL B422 399 | D. Barberis <i>et al.</i> | (WA 102 Collab.) |
| ACCIARRI | 97T | PL B413 147 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ALBRECHT | 97B | ZPHY C74 469 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| AMELIN | 95B | PL B356 595 | D.V. Amelin <i>et al.</i> | (SERP, TBIL) |
| BERDNIKOV | 94 | PL B337 219 | E.B. Berdnikov <i>et al.</i> | (SERP, TBIL) |
| ANTREASIAN | 90 | ZPHY C48 561 | D. Antreasian <i>et al.</i> | (Crystal Ball Collab.) |
| BEHREND | 90C | ZPHY C46 583 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| ANTIPOV | 87 | EPL 4 403 | Y.M. Antipov <i>et al.</i> | (SERP, JINR, INRM+) |
| BELLINI | 85 | SJNP 41 781 | D. Bellini <i>et al.</i> | |
| | | Translated from YAF 41 | 1223. | |
| ARMSTRONG | 82B | NP B202 1 | T.A. Armstrong, B. Baccari | (AACH3, BARI, BONN+) |
| DAUM | 81B | NP B182 269 | C. Daum <i>et al.</i> | (AMST, CERN, CRAC, MPIM+) |
| EVANGELIS... | 81 | NP B178 197 | C. Evangelista <i>et al.</i> | (BARI, BONN, CERN+) |
| Also | | NP B186 594 | C. Evangelista | |
| DAUM | 80D | PL 89B 285 | C. Daum <i>et al.</i> | (AMST, CERN, CRAC, MPIM+) JP |
| BALTAY | 77 | PRL 39 591 | C. Baltay, C.V. Cautis, M. Kalelkar | (COLU) JP |
| ASCOLI | 73 | PR D7 669 | G. Ascoli | (ILL, TNTO, GENO, HAMB, MILA+) JP |
| CRENNELL | 70 | PRL 24 781 | D.J. Crennell <i>et al.</i> | (BNL) |
| ARMENISE | 69 | LNC 2 501 | N. Armenise <i>et al.</i> | (BARI, BGNA, FIRZ) |
| BALTAY | 68 | PRL 20 887 | C. Baltay <i>et al.</i> | (COLU, ROCH, RUTG, YALE) I |
| BARTSCH | 68 | NP B7 345 | J. Bartsch <i>et al.</i> | (AACH, BERL, CERN) JP |

OTHER RELATED PAPERS

| | | | | |
|-------------|-----|-------------------------|------------------------------|----------------------------|
| DZIERBA | 06 | PR D73 072001 | A.R. Dzierba <i>et al.</i> | (BNL E852 Collab.) |
| PAGE | 03 | PL B566 108 | P. Page, S. Capstick | |
| ZAIMIDOROGA | 99 | PAN 30 1 | O.A. Zaimidoriga | |
| | | Translated from SJPN 30 | 5. | |
| CHEN | 83B | PR D28 2304 | T.Y. Chen <i>et al.</i> | (ARIZ, FNAL, FLOR, NDAM+) |
| LEEDOM | 83 | PR D27 1426 | I.D. Leedom <i>et al.</i> | (PURD, TNTO) |
| BELLINI | 82B | NP B199 1 | G. Bellini <i>et al.</i> | (CERN, MILA, JINR+) |
| DAUM | 81B | NP B182 269 | C. Daum <i>et al.</i> | (AMST, CERN, CRAC, MPIM+) |
| PERNEGR | 78 | NP B134 436 | J. Pernegr <i>et al.</i> | (ETH, CERN, LOIC+) |
| FOCACCI | 66 | PRL 17 890 | M.N. Focacci <i>et al.</i> | (CERN) |
| LEVRAT | 66 | PL 22 714 | B. Levrat <i>et al.</i> | |
| VETLITSKY | 66 | PL 21 579 | I.A. Vetlitsky <i>et al.</i> | (ITEP) |
| FORINO | 65B | PL 19 68 | A. Forino <i>et al.</i> | (BGNA, BARI, FIRZ, ORSAY+) |