

SALT RUSH

Juncus lesueurii Boland

Plant Symbol = JULE

Contributed by: USDA NRCS National Plant Data Center



Alfred Brousseau
© Brother Eric Vogel, St. Mary's College
@ CalPhotos

Uses

Ethnobotanic: Coiled basketry prevails in Southern California, with the mottled yellowish brown of *Juncus* rush providing a natural colored and variegated background (Turnbaugh and Turnbaugh 1986). *Juncus* stems are used in the coiled baskets of Southern California basket weavers such as the Cahuilla, Luiseño, Chumash, Diegueño, Agua Caliente, Gabrieliño, Juaneño, Death Valley Shoshone, and Fernandeno (Barrows 1967; Murphy 1959). Chumash baskets, from southern California, are made with *Juncus* stems for the tan color and roots for the black color (Timbrook 1997). The foundation material is made of *Juncus lesueurii* or *Juncus balticus*, and the sewing material is made of *Juncus textilis*.

Rushes are cut off at ground level, or at the length desired. The rush, in its natural state, furnishes a

variety of colors; a deep red near the base, lightening in color upwards passing through several shades of light brown, and ending at the top in a brownish

yellow. *Juncus* stems can be bleached in the summer sun for several months to assure a light tan uniform color.

The Cahuilla, Diegeño, Luiseño, and Chumash dye the mature rushes black by steeping them for several hours in an infusion of either horned sea-blite (*Suaeda calceoliformis*) or bush seepweed (*Sueda moquinii*). This dye is very penetrating, and the color is durable, but has a fetid, disagreeable smell. *Juncus* species are also dyed yellow in an infusion of indigo bush (*Psoralea emoryi*) (Barrows 1967; Merrill 1970).

Other Uses: A wide range of mammal and avian species for food and habitat (Hoag and Zierke 1998) uses *Juncus* species. Waterfowl, songbirds and small mammals such as jack rabbits, cottontail, muskrat, porcupine, and gopher (Martin 1951) eats rush seeds. Rushes help improve habitat for amphibians and spawning areas for fish. Muskrats feed on the roots and rhizomes, and various wetland wading birds find shelter among the stems.

Rushes provide the following conservation uses: erosion control, sediment accretion and stabilization, nutrient uptake and transformation, wildlife food and cover, restoration and creation of wetland ecosystems, and wastewater treatment applications. The rhizomatous nature, nitrogen fixation capabilities, dense root system, and phenotypic plasticity to flooding and drought stress provide high soil and slope stabilization capabilities, particularly in areas with flooded soils or fluctuating hydrology. The rhizomes form a matrix for many beneficial bacteria, making this plant an excellent addition for wastewater treatment. Rushes tend to be resistant to grazing pressure and fairly unpalatable to cattle, so they tend to increase in species composition in stockwater ponds and troughs.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

Plant Materials <<http://plant-materials.nrcs.usda.gov/>>

Plant Fact Sheet/Guide Coordination Page <<http://plant-materials.nrcs.usda.gov/intranet/pfs.html>>

National Plant Data Center <<http://npdc.usda.gov>>

Description

General: Rush Family (Juncaceae). *Juncus lesueurii* is a grasslike, perennial, rhizomatous wetland plant. Salt rush has stems that are 3-9 dm tall, stout, smooth, erect or aching, with a distinctive twist to the stems. The stems arise from stout, creeping rhizomes. The terete leaf sheaths are bladeless, and shining or light brown in color. The inflorescence appears lateral with open branches, the lowest bract cylindrical, with many flowers. The capsule is oblong-ovoid, light brown, and three-angled. This species is sometimes confused with *Juncus balticus*, from which it may be distinguished by its longer and usually darker-colored segments.

Distribution

For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site.

Establishment

Adaptation: Salt rush grows in salt marshes or sand dunes along the California coast from Ventura County north to British Columbia. *Juncus lesueurii* can tolerate mild to moderate soil salinity and flooded soil conditions. Often these plants are found on drier or seasonally fluctuating wetland sites (for example, dune swales) and can tolerate periodic drought.

General: *Juncus* species may be planted from bare rootstock or seedlings, from container stalk or directly seeded into the soil. Bare rootstock or seedlings are preferred revegetation methods where there is moving water. *Juncus lesueurii* requires moderate summer watering (irrigation), generally 1 - 4 times per month depending on the absorption rate and water retention capacity of the soil. Salt rush plants may need to have their roots in moist or wet soils. These native plants are especially good for stabilizing or restoring disturbed or degraded areas for erosion and slope control. Salt rush may become invasive.

Live Plant Collections: J. Chris Hoag and Mike Zierke (USDA, NRCS, Plant Materials Center, Aberdeen, Idaho) provided the following information on *Juncus balticus*. Due to their taxonomic and habitat similarity, it is likely that *Juncus lesueurii* establishes in a similar manner. Planting plugs is the surest way to establish a new stand of this species. Plug spacing of 25-30 cm will fill in within one growing season. Fluctuating the water level during the establishment period may speed spread of the

rush. Water levels can be managed to enhance spread and control weeds.

- Clip leaves and stems to 15 to 25 cm (6 to 10 inches) before planting; this allows the plant to allocate more energy into root production. Transplants should be planted as soon as possible in moist (not flooded or anoxic) soils. Plants should be transported and stored in a cool location prior to planting. The roots should always remain moist or in water until planted.
- Soil should be kept saturated after planting. Plants can tolerate 2.5 - 8 cm of standing water as long as the level fluctuates over the growing season. Allow roots to become established before flooding soils if possible.
- Ideally, plants should be planted in late fall just after the first rains (usually late October to November). Survival is highest when plants are dormant and soils are moist.
- Fertilization is very helpful for plant growth and reproduction. Many more seeds are produced with moderate fertilization.

Seed Collections:

- The flowering period is late May to August, occasionally to September. Seed ripens in early August. Phenology will change by area, aspect, elevation, and specific site conditions.
- Seed may be collected by hand, using a pair of hand shears, or with a gas-powered handheld seed harvester.
- The tiny, black seeds are easily lost from the capsules when collecting by hand. Be careful to keep capsules upright before putting in collection bag. Use paper sacks when collecting seeds for this species.
- To clean the seed, run the collection through a hammer mill to break up debris and knock the seeds loose. Use a 1/20 inch screen on the top and a solid sheet on the bottom of the seed cleaner. Adjust the air flow to blow off the chaff. Shaking the hammer milled collection to settle seed to the bottom of the pan can speed up the cleaning process. The top portion of the chaff can then be discarded and the seed-rich mixture that is left in the bottom can be run through the seed cleaner.

Seed germination in greenhouse:

- Seeds need light, moisture and heat for germination. Soaking the seeds in water for 1 - 7 days will decrease the time the seed takes to sprout.
- To grow seeds, place on soil surface and press in lightly to assure good soil contact. Do not cover

the seed. Soil should be kept moist. Greenhouse should be kept hot (32-38°C).

- Seeds begin to germinate in approximately 1 week. Maintain soil moisture until plants are to be transplanted. Seedlings cannot withstand long periods without water while growing in the greenhouse.
- Plants are ready in 100 - 120 days to come out as plugs. By planting seeds in August, plugs are ready to plant in soil by November. These plants are very small; growing plants to a larger size will result in increased revegetation success.

Management

Hydrology is the most important factor in determining wetland type, revegetation success, and wetland function and value. Changes in water levels influence species composition, structure, and distribution of plant communities. Water management is absolutely critical during plant establishment, and remains crucial through the life of the wetland for proper community management (Hoag et al. 1995). *Juncus* species can tolerate periods of drought and total inundation. It is important to keep transplanted plugs moist, not flooded, until roots are established. Water levels can then be managed to either enhance or reduce spread as well as control terrestrial weeds.

Muskrats have evolved with wetland ecosystems and form a valuable component of healthy functioning wetland communities. Muskrats use emergent wetland vegetation such as *Juncus* species for hut construction and for food. Typically, an area of open water is created around the huts.

Juncus species tend to be fairly resilient to insect and disease problems. Aphids may feed on the stems, but rarely cause significant damage. If an insect or disease problem is encountered in the greenhouse, treatment options may be limited by cultural constraints if these plants are to be used by Indian basket weavers. *Juncus* culms are split with the mouth to process basketry materials; therefore, an unusually high degree of human exposure and risk occur with plants designated for ethnobotanic use. Rushes are perennial, rhizomatous plants. In most cases, they will out-compete other species within the wetland area of the site, eliminating the need for manual or chemical control of invasive species.

Traditional Resource Management: Management of *Juncus lesueurii* stands includes the following: ownership of prime basket rush sites, stimulation of new growth through harvesting stalks, periodic

burning, and not harvesting when soils are very mucky and likely to be compacted. According to one Northern Diegueño basket weaver, most weavers have favorite collecting areas where the basket rush is plentiful and having characteristics valued by basket weavers (long, flexible, tough stems, deep red color, access is available and relatively easy). Any *Juncus* stand will have plants that are immature, those which are mature but still in seed, and those which are starting to senesce. The stalks are cut above the rhizomes and roots, leaving plenty of buds to re-grow new shoots. As with other rhizomatous species, harvesting stimulates new growth and maintains the clone in a juvenile or immature growth phase, where productivity is highest. The only harvesting prohibition might be during times of heavy rain or flooding, when deep water and mud make many plants inaccessible.

Cultivars, Improved and Selected Materials (and area of origin)

Contact your local Natural Resources Conservation Service (formerly Soil Conservation Service) office for more information. Look in the phone book under "United States Government." The Natural Resources Conservation Service will be listed under the subheading "Department of Agriculture."

References

- Cooke, S.S. 1997. *A field guide to the common wetland plants of Western Washington and Northwestern Oregon*. Seattle Audubon Society and Washington Native Plant Society. 414 pp.
- Dahl, T.E. & C.E. Johnson 1991. *Status and trends of wetlands in the coterminous United States, mid-1970s to mid-1980s*. USDI, FWS, Washington, D.C. 28 pp.
- Gunther, E. 1945 rev. 1973. *Ethnobotany of western Washington*. University of Washington Publications in Anthropology, 10(1). University of Washington Press, Seattle, Washington.
- Hickman, J.C. (ed.) 1993. *The Jepson Manual. Higher plants of California*. University of California Press. 1399 pp.
- Hoag, J.C. n.d. *Wetland plant fact sheet - Baltic rush (Juncus balticus)*. Interagency Riparian/Wetland Project. USDA, NRCS, Plant Materials Center, Aberdeen, Idaho. 83210. 3 pp.
- Hoag, J.C. & M. Zierke (February) 1998. *A reference guide for the collection and use of ten*

common wetland plants of the Great Basin and Intermountain West. Riparian/Wetland Project Information Series No. 13. USDA, NRCS, Plant Materials Center, Aberdeen, Idaho. 13 pp.

Hoag, J.C. & M.E. Sellers (April) 1995. *Use of greenhouse propagated wetland plants versus live transplants to vegetate constructed or created wetlands*. Riparian/Wetland Project Information Series No. 7. USDA, NRCS, Plant Materials Center, Aberdeen, Idaho. 6 pp.

Hurd, E.G., N.L. Shaw, & L.C. Smithman 1992. *Cyperaceae and Juncaceae - selected low-elevation species*. Proceedings of Symposium of Ecology, Management, and Restoration of Intermountain Annual Rangelands, Boise, Idaho. May 18-22, 1992. Pages 380-383.

Manning, M.E., S.R. Swanson, T. Svejcar, & J. Trent 1989. *Rooting characteristics of four Intermountain meadow community types*. Journal of Range Management 42(4):309-312.

Martin, A.C., H.S. Zim, & A.L. Nelson 1951. *American wildlife and plants: A guide to wildlife food habits*. Dover Publications, Inc., New York, New York. 500 pp.

Merrill, R.E. 1970. *Plants used in basketry by the California Indians*. Acoma Books, Ramona, California.

Moser, C.L. 1993. *Native American basketry of southern California*. Riverside Museum Press. 155 pp.

Murphy, E.V.A. 1959. *Indian uses of native plants*. Mendocino County Historical Society. 81 pp.

Strike, S.S. 1994. *Ethnobotany of the California Indians. Volume 2. Aboriginal uses of California's indigenous plants*. Koeltz Scientific Books, USA/Germany. 210 pp.

Timbrook, J. (June) 1997. *California Indian Basketweavers Association newsletter*.

Turnbaugh, S.P. & W.A. Turnbaugh 1986. *Indian baskets*. Schiffler Publishing, Ltd., West Chester, Pennsylvania. Pages 194-205.

Prepared By

Michelle Stevens

formerly USDA, NRCS, National Plant Data Center

Species Coordinator

M. Kat Anderson

USDA, NRCS, National Plant Data Center
c/o Environmental Horticulture Department,
University of California, Davis, California

Edited: 05dec00 jsp; 20may03 ahv; 060801 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Read about [Civil Rights at the Natural Resources Conservation Service](#).