



Summary and Recommendations

This Report addresses the substantial impact of daily dietary patterns on the health of Americans. Good health does not always come easily. It is the product of complex interactions among environmental, behavioral, social, and genetic factors. Some of these are, for practical purposes, beyond personal control. But there are many ways in which each of us can influence our chances for good health through the daily choices we make.

In recent years, scientific investigations have produced abundant information on the ways personal behavior affects health. This information can help us decide whether to smoke, when and how much to drink, how far to walk or climb stairs, whether to wear seat belts, and how or whether to engage in any other activity that might alter the risk of incurring disease or disability. For the two out of three adult Americans who do not smoke and do not drink excessively, one personal choice seems to influence long-term health prospects more than any other: what we eat.

Food sustains us, it can be a source of considerable pleasure, it is a reflection of our rich social fabric and cultural heritage, it adds valued dimensions to our lives. Yet what we eat may affect our risk for several of the leading causes of death for Americans, notably, coronary heart disease, stroke, atherosclerosis, diabetes, and some types of cancer. These disorders together now account for more than two-thirds of all deaths in the United States.

Undernutrition remains a problem in several parts of the world, as well as for certain Americans. But for most of us the more likely problem has become one of overeating—too many calories for our activity levels and an imbalance in the nutrients consumed along with them. Although much is still uncertain about how dietary patterns protect or injure human health, enough has been learned about the overall health impact of the dietary patterns now prevalent in our society to recommend significant changes in those patterns.

This first *Surgeon General's Report on Nutrition and Health* offers comprehensive documentation of the scientific basis for the recommended dietary changes. Through the extensive review contained in its chapters, the Report examines in detail current knowledge about the relationships among specific dietary practices and specific disease conditions and sum-

marizes the implications of this information for individual food choices, public health policy initiatives, and further research. **The Report's main conclusion is that overconsumption of certain dietary components is now a major concern for Americans. While many food factors are involved, chief among them is the disproportionate consumption of foods high in fats, often at the expense of foods high in complex carbohydrates and fiber that may be more conducive to health.** A list of the key recommendations based on the evidence presented in the Report is provided in Table 1.

Magnitude of the Problem

Diet has always had a vital influence on health. Until as recently as the 1940's, diseases such as rickets, pellagra, scurvy, beriberi, xerophthalmia, and goiter (caused by lack of adequate dietary vitamin D, niacin, vitamin C, thiamin, vitamin A, and iodine, respectively) were prevalent in this country and throughout the world. Today, thanks to an abundant food supply, fortification of some foods with critical trace nutrients, and better methods for determining and improving the nutrient content of foods, such "deficiency" diseases have been virtually eliminated in developed countries. For example, the introduction of iodized salt in the 1920's contributed greatly to eliminating iodine-deficiency goiter as a public health problem in the United States. Similarly, pellagra disappeared subsequent to the discovery of the dietary causes of this disease. Nutrient deficiencies are reported rarely in the United States, and the few cases of protein-energy malnutrition that are listed annually as causes of death generally occur as a secondary result of severe illness or injury, child neglect, the problems of the house-bound aged, premature birth, alcoholism, or some combination of these factors.

As the diseases of nutritional deficiency have diminished, they have been replaced by diseases of dietary excess and imbalance—problems that now rank among the leading causes of illness and death in the United States, touch the lives of most Americans, and generate substantial health care costs. Table 2, for example, lists the 10 leading causes of death in the United States in 1987.

In addition to the five of these causes that scientific studies have associated with diet (coronary heart disease, some types of cancer, stroke, diabetes mellitus, and atherosclerosis), another three—cirrhosis of the liver, accidents, and suicides—have been associated with excessive alcohol intake.

Table 1
Recommendations

Issues for Most People:

- *Fats and cholesterol:* Reduce consumption of fat (especially saturated fat) and cholesterol. Choose foods relatively low in these substances, such as vegetables, fruits, whole grain foods, fish, poultry, lean meats, and low-fat dairy products. Use food preparation methods that add little or no fat.
- *Energy and weight control:* Achieve and maintain a desirable body weight. To do so, choose a dietary pattern in which energy (caloric) intake is consistent with energy expenditure. To reduce energy intake, limit consumption of foods relatively high in calories, fats, and sugars, and minimize alcohol consumption. Increase energy expenditure through regular and sustained physical activity.
- *Complex carbohydrates and fiber:* Increase consumption of whole grain foods and cereal products, vegetables (including dried beans and peas), and fruits.
- *Sodium:* Reduce intake of sodium by choosing foods relatively low in sodium and limiting the amount of salt added in food preparation and at the table.
- *Alcohol:* To reduce the risk for chronic disease, take alcohol only in moderation (no more than two drinks a day), if at all. Avoid drinking any alcohol before or while driving, operating machinery, taking medications, or engaging in any other activity requiring judgment. Avoid drinking alcohol while pregnant.

Other Issues for Some People:

- *Fluoride:* Community water systems should contain fluoride at optimal levels for prevention of tooth decay. If such water is not available, use other appropriate sources of fluoride.
 - *Sugars:* Those who are particularly vulnerable to dental caries (cavities), especially children, should limit their consumption and frequency of use of foods high in sugars.
 - *Calcium:* Adolescent girls and adult women should increase consumption of foods high in calcium, including low-fat dairy products.
 - *Iron:* Children, adolescents, and women of childbearing age should be sure to consume foods that are good sources of iron, such as lean meats, fish, certain beans, and iron-enriched cereals and whole grain products. This issue is of special concern for low-income families.
-

Table 2
Estimated Total Deaths and Percent of Total Deaths for the
10 Leading Causes of Death: United States, 1987

Rank	Cause of Death	Number	Percent of Total Deaths
1 ^a	Heart diseases	759,400	35.7
	(Coronary heart disease)	(511,700)	(24.1)
	(Other heart disease)	(247,700)	(11.6)
2 ^a	Cancers	476,700	22.4
3 ^a	Strokes	148,700	7.0
4 ^b	Unintentional injuries	92,500	4.4
	(Motor vehicle)	(46,800)	(2.2)
	(All others)	(45,700)	(2.2)
5	Chronic obstructive lung diseases	78,000	3.7
6	Pneumonia and influenza	68,600	3.2
7 ^a	Diabetes mellitus	37,800	1.8
8 ^b	Suicide	29,600	1.4
9 ^b	Chronic liver disease and cirrhosis	26,000	1.2
10 ^a	Atherosclerosis	23,100	1.1
. . .	All causes	2,125,100	100.0

^aCauses of death in which diet plays a part.

^bCauses of death in which excessive alcohol consumption plays a part.

Source: National Center for Health Statistics, *Monthly Vital Statistics Report*, vol. 37, no. 1, April 25, 1988.

Although the precise proportion attributable to diet is uncertain, these eight conditions accounted for nearly 1.5 million of the 2.1 million total deaths in 1987. Dietary excesses or imbalances also contribute to other problems such as high blood pressure, obesity, dental diseases, osteoporosis, and gastrointestinal diseases. Together, these diet-related conditions inflict a substantial burden of illness on Americans. For example:

- **Coronary Heart Disease.** Despite the recent sharp decline in the death rate from this condition, coronary heart disease still accounts for the largest number of deaths in the United States. More than 1.25 million heart attacks occur each year (two-thirds of them in men), and more than 500,000 people die as a result. In 1985, illness and deaths from coronary heart disease cost Americans an estimated \$49 billion in direct health care expenditures and lost productivity.
- **Stroke.** Strokes occur in about 500,000 persons per year in the United States, resulting in nearly 150,000 deaths in 1987 and long-term disability for many individuals. Approximately 2 million living Americans suffer from stroke-related disabilities, at an estimated annual cost of more than \$11 billion.

- **High Blood Pressure.** High blood pressure (hypertension) is a major risk factor for both heart disease and stroke. Almost 58 million people in the United States have hypertension, including 39 million who are under age 65. The occurrence of hypertension increases with age and is higher for black Americans (of which 38 percent are hypertensive) than for white Americans (29 percent).
- **Cancer.** More than 475,000 persons died of cancer in the United States in 1987, making it the second leading cause of death in this country. During the same period, more than 900,000 new cases of cancer occurred. The costs of cancer for 1985 have been estimated to be \$22 billion for direct health care, \$9 billion in lost productivity due to treatment or disability, and \$41 billion in lost productivity due to premature mortality, for a total cost of \$72 billion.
- **Diabetes Mellitus.** Approximately 11 million Americans have diabetes, but almost half of them have not been diagnosed. In addition to the nearly 38,000 deaths in 1987 attributed directly to this condition, diabetes also contributes to an estimated 95,000 deaths per year from associated cardiovascular and kidney complications. In 1985, diabetes was estimated to cost \$13.8 billion per year, or about 3.6 percent of total health care expenses.
- **Obesity.** Obesity affects approximately 34 million adults ages 20 to 74 years in the United States, with the highest rates observed among the poor and minority groups. Obesity is a risk factor for coronary heart disease, high blood pressure, diabetes, and possibly some types of cancer as well as other chronic diseases.
- **Osteoporosis.** Approximately 15 to 20 million Americans are affected by osteoporosis, which contributes to some 1.3 million bone fractures per year in persons 45 years and older. One-third of women 65 years and older have vertebral fractures. On the basis of x-ray evidence, by age 90 one-third of women and one-sixth of men will have suffered hip fractures, leading to death in 12 to 20 percent of those cases and to long-term nursing care for many who survive. The total costs of osteoporosis to the U.S. economy were estimated to be \$7 to \$10 billion in 1983.
- **Dental Diseases.** Dental caries and periodontal disease continue to affect a large proportion of Americans and cause substantial pain, restriction of activity, and work loss. Although dental caries among children, as well as some forms of adult periodontal disease, appear to be declining, the overall prevalence of these conditions imposes a substantial burden on Americans. The costs of dental care were estimated at \$21.3 billion in 1985.

- *Diverticular Disease.* Because most persons with diverticular disease do not have symptoms, the true prevalence of this condition is unknown. Frequency increases with age, and up to 70 percent of people between the ages of 40 and 70 may be affected. In 1980, diverticulosis was accountable for some 200,000 hospitalizations.

In assessing the role that diet might play in prevention of these conditions, it must be understood that they are caused by a combination (and interaction) of multiple environmental, behavioral, social, and genetic factors. The exact proportion that can be attributed directly to diet is uncertain. Although some experts have suggested that dietary factors overall are responsible for perhaps a third or more of all cases of cancer, and similar estimates have been made for coronary heart disease, such suggestions are based on interpretations of research studies that cannot completely distinguish dietary from genetic, behavioral, or environmental causes.

We know, for example, that cigarette smoking exerts a powerful influence on the occurrence of both coronary heart disease and some types of cancer. We also know that some people are genetically predisposed to coronary heart disease, stroke, and diabetes and that the interaction of genetic predisposition with dietary patterns is an important determinant of individual risk. For these reasons, it is not yet possible to determine the proportion of chronic diseases that could be reduced by dietary changes. Nonetheless, it is now clear that diet contributes in substantial ways to the development of these diseases and that modification of diet can contribute to their prevention. The magnitude of the health and economic cost of diet-related disease suggests the importance of the dietary changes suggested. This Report reviews these issues in detail.

Nature of the Evidence

Whereas centuries of clinical observations and decades of basic and clinical research prove that dietary deficiencies of single, identifiable nutrients can cause disease, research on the relationship of dietary excesses and imbalances to chronic disease yields results that rarely provide such direct proof of causality. Instead, investigators must piece together various kinds of information from several kinds of sources. Nevertheless, the quantity of current animal, laboratory, clinical, and epidemiologic evidence that associates dietary excesses and imbalances with chronic disease is substantial and, when evaluated according to established principles, compelling.

Scientists must often draw inferences about the relationships between dietary factors and disease from laboratory animal studies or human meta-

bolic and population studies that approach the issues indirectly. Data sources for such human studies include clinical and laboratory measurements of physiologic indicators of nutritional status or risk factors, as well as dietary intake data estimated for populations or individuals. Epidemiologic studies using these data compare dietary intake and disease rates in different countries or in defined groups within the same country.

Interpretations of animal studies are limited by uncertainties about their applicability to people. Clinical, laboratory, and dietary intake studies can provide useful information, but each has limitations. Currently available clinical and laboratory measurements reveal only a small part of the complex physiological responses to diet, and they may reflect past rather than current nutritional status. Dietary surveys depend on accurate recall of the types and portion sizes of consumed foods as well as on the assumption that the food intake during any one period represents typical intake. Reported intake, however, is not always accurate, and intake reported for a given period may differ significantly from that typical of longer time periods. Dietary intake data provide useful indicators for populations, but even when an association or correlation between a dietary factor and a disease is observed, it is often difficult to prove that the dietary factor is an actual or sole cause of that disease.

This difference between association and causation is basic to understanding the scientific evidence that links diet to chronic disease. Uncertainties in the ability to determine causation have sometimes made it difficult to achieve consensus on appropriate public health nutrition policies. Established principles require evaluation of the supporting evidence for a given association between a dietary factor and a disease on the basis of its consistency, strength, specificity, and biological plausibility. The evidence showing that dietary intake of saturated fat raises blood cholesterol, which in turn increases the chance of coronary heart disease, illustrates this point. The similarity in results from laboratory, clinical, and epidemiologic research, the apparent relationship between dose and effect in these studies, the observations that the increase in blood cholesterol level is specific to saturated fatty acids but not to other types, and the biological plausibility of explanations for the observations, when taken together, provide considerable support for concluding that the association is causal, at least for some individuals.

For some of the other diseases reviewed in this Report, the available evidence is less complete and less consistent. Nevertheless, much evidence supports credible associations between a dietary pattern of excesses

and imbalances and several important chronic diseases. These associations, in turn, suggest that the overall health of Americans could be improved by a few specific but fundamental dietary changes.

Key Findings and Recommendations

Even though the results of various individual studies may be inconclusive, the preponderance of the evidence presented in the Report's comprehensive scientific review substantiates an association between dietary factors and rates of chronic diseases. In particular, the evidence suggests strongly that a dietary pattern that contains excessive intake of foods high in calories, fat (especially saturated fat), cholesterol, and sodium, but that is low in complex carbohydrates and fiber, is one that contributes significantly to the high rates of major chronic diseases among Americans. It also suggests that reversing such dietary patterns should lead to a reduced incidence of these chronic diseases.

This *Surgeon General's Report on Nutrition and Health* provides a comprehensive review of the most important scientific evidence in support of current Federal nutrition policy as stated in the *Dietary Guidelines for Americans*. These *Guidelines*, issued jointly by the Department of Agriculture and the Department of Health and Human Services, recommend:

- Eat a variety of foods.
- Maintain desirable weight.
- Avoid too much fat, saturated fat, and cholesterol.
- Eat foods with adequate starch and fiber.
- Avoid too much sugar.
- Avoid too much sodium.
- If you drink alcoholic beverages, do so in moderation.

Evidence presented in this Report expands the focus of these seven guidelines and provides considerable insight into priorities. Clearly emerging as the primary priority for dietary change is the recommendation to reduce intake of total fats, especially saturated fat, because of their relationship to development of several important chronic disease conditions. Because excess body weight is a risk factor for several chronic diseases, maintenance of desirable weight is also an important public health priority. Evidence further supports the recommendation to consume a dietary pattern that contains a variety of foods, provided that these foods are generally low in calories, fat, saturated fat, cholesterol, and sodium.

Taken together, the recommendations in this Report promote a dietary pattern that emphasizes consumption of vegetables, fruits, and whole grain products—foods that are rich in complex carbohydrates and fiber and relatively low in calories—and of fish, poultry prepared without skin, lean meats, and low-fat dairy products selected to minimize consumption of total fat, saturated fat, and cholesterol.

The evidence presented in this Report suggests that such overall dietary changes will lead to substantial improvements in the nutritional quality of the American diet. Consuming a higher proportion of calories from fruits, vegetables, and grains may lead to a modest reduction in protein intake for some people, but this reduction is unlikely to impair nutritional status. Average levels of protein consumption in the United States, 60 grams per day for women and 90 grams per day for men, are well above the National Research Council's recommendations of 44 and 56 grams per day, respectively.

The evidence also suggests that most Americans generally need not consume nutrient supplements. An estimated 40 percent of Americans consume supplemental vitamins, minerals, or other dietary components at an annual cost of more than \$2.7 billion. Although nutrient supplements are usually safe in amounts corresponding to the Recommended Dietary Allowances (and such Allowances are set to ensure that the nutrient needs of practically all the population are met), there are no known advantages to healthy people consuming excess amounts of any nutrient, and amounts greatly exceeding recommended levels can be harmful. For example, some nutrients such as selenium have a narrow range of safe level of intake. Toxicity has been reported for most minerals and trace elements, as well as some vitamins, indicating that excessive supplementation with these substances can be hazardous.

Finally, some recommendations for dietary change apply broadly to the general public whereas others apply only to specific population groups. These major findings and recommendations of *The Surgeon General's Report on Nutrition and Health* are noted below.

Issues for Most People

- **Fats and cholesterol:** Reduce consumption of fat (especially saturated fat) and cholesterol. Choose foods relatively low in these substances, such as vegetables, fruits, whole grain foods, fish, poultry, lean meats, and low-fat dairy products. Use food preparation methods that add little or no fat.

High intake of total dietary fat is associated with increased risk for obesity, some types of cancer, and possibly gallbladder disease. Epidemiologic, clinical, and animal studies provide strong and consistent evidence for the relationship between saturated fat intake, high blood cholesterol, and increased risk for coronary heart disease. Conversely, reducing blood cholesterol levels reduces the risk for death from coronary heart disease. Excessive saturated fat consumption is the major dietary contributor to total blood cholesterol levels. Dietary cholesterol raises blood cholesterol levels, but the effect is less pronounced than that of saturated fat. While polyunsaturated fatty acid consumption, and probably monounsaturated fatty acid consumption, lowers total blood cholesterol, the precise effects of specific fatty acids are not well defined.

Dietary fat contributes more than twice as many calories as equal quantities (by weight) of either protein or carbohydrate, and some studies indicate that diets high in total fat are associated with higher obesity rates. In addition, there is substantial, although not yet conclusive, epidemiologic and animal evidence in support of an association between dietary fat intake and increased risk for cancer, especially breast and colon cancer. Similarly, epidemiologic studies suggest an association between gallbladder disease, excess caloric intake, high dietary fat, and obesity. More precise conclusions about the role of dietary fat await the development of improved methods to distinguish among the contributions of the high-calorie, high-fat, and low-fiber components of current American dietary patterns.

At present, dietary fat accounts for about 37 percent of the total energy intake of Americans—well above the upper limit of 30 percent recommended by the American Heart Association and the American Cancer Society, and above the percent consumed by many societies, such as Mediterranean countries, Japan, and China, for example, where coronary heart disease rates are much lower than those observed in the United States. Consumption of saturated fat and cholesterol is also substantially higher among many Americans than levels recommended by several expert groups.

The major dietary sources of fat in the American diet are meat, poultry, fish, dairy products, and fats and oils. Animal products tend to be higher in both total and saturated fats than most plant sources. Although some plant fats such as coconut and palm kernel oils also contain high proportions of saturated fatty acids, these make minor contributions to total intake of saturated fats in the United States. Dietary cholesterol is found only in foods of animal origin, such as eggs, meat, poultry, fish, and dairy prod-

ucts. To help reduce consumption of total fat, especially saturated fat and cholesterol, food choices should emphasize intake of fruits, vegetables, and whole grain products and cereals. They should also emphasize consumption of fish, poultry prepared without skin, lean meats, and low-fat dairy products. Among vegetable fats, those that are more unsaturated are better choices.

- ***Energy and weight control:*** Achieve and maintain a desirable body weight. To do so, choose a dietary pattern in which energy (caloric) intake is consistent with energy expenditure. To reduce energy intake, limit consumption of foods relatively high in calories, fats, and sugars and minimize alcohol consumption. Increase energy expenditure through regular and sustained physical activity.

People are considered overweight if their body mass index, or BMI (a ratio of weight to height described in the Report), exceeds the 85th percentile for young American adults (approximately 120 percent of desirable body weight); they are considered severely overweight if their BMI exceeds the 95th percentile (approximately 140 percent of desirable body weight). Overweight individuals are at increased risk for diabetes mellitus, high blood pressure and stroke, coronary heart disease, some types of cancer, and gallbladder disease. Epidemiologic and animal studies have shown consistently that overall risk for death is increased with excess weight, with risk increasing as severity of obesity increases.

Type II (noninsulin-dependent) diabetes mellitus accounts for approximately 90 percent of all cases of diabetes and is strongly associated with obesity. Clinical studies indicate that weight loss can improve control of Type II diabetes.

Obesity increases the risk for high blood pressure, and consequently for stroke; it also increases blood cholesterol levels associated with coronary heart disease. In addition, it appears to be an independent risk factor for coronary heart disease. Weight reduction has been shown to reduce high blood pressure and high blood cholesterol. Most obese individuals who achieve a more desirable body weight improve their cholesterol profile, achieving a decrease in both total blood cholesterol and LDL (low density lipoprotein) cholesterol.

Some studies have found an association between overweight and increased risk for several cancers, especially cancer of the uterus and breast. In addition, overweight increases the risk for gallbladder disease.

More than a quarter of American adults are overweight. Black women age 45 and above have the highest prevalence, about 60 percent. Although evidence suggests a genetic component to the tendency of many people to become overweight, patterns of dietary caloric intake and energy expenditure play a key role. Sustained and long-term efforts to reduce body weight can best be achieved as a result of improving energy balance by reducing energy consumption and raising energy expenditure through physical activity and exercise.

Maintenance of desirable body weight throughout the lifespan requires a balance between energy (calorie) intake and expenditure. Weight control may be facilitated by decreasing energy intake, especially by choosing foods relatively low in calories, fats, and sugars, and by minimizing alcohol consumption. Energy expenditure can be enhanced through regular physical activities such as daily walks or by jogging, bicycling, or swimming at least three times a week for at least 20 minutes.

- ***Complex carbohydrates and fiber:* Increase consumption of whole grain foods and cereal products, vegetables (including dried beans and peas), and fruits.**

Dietary patterns emphasizing foods high in complex carbohydrates and fiber are associated with lower rates of diverticulosis and some types of cancer. The association shown in epidemiologic and animal studies between diets high in complex carbohydrates and reduced risk for coronary heart disease and diabetes mellitus is, however, difficult to interpret. The fact that such diets tend also to be lower in energy and fats, especially saturated fat and cholesterol, clearly contributes to this difficulty. Some evidence from clinical studies also suggests that water-soluble fibers from foods such as oat bran, beans, or certain fruits are associated with lower blood glucose and blood lipid levels. Consuming foods with dietary fiber is usually beneficial in the management of constipation and diverticular disease.

While inconclusive, some evidence also suggests that an overall increase in intake of foods high in fiber might decrease the risk for colon cancer. Among several unresolved issues is the role of the various types of fiber, which differ in their effects on water-holding capacity, viscosity, bacterial fermentation, and intestinal transit time.

Other food components associated with decreased cancer risk are commonly found in diets high in whole grain cereal products containing complex carbohydrates and fiber. In addition, some epidemiologic evidence

suggests that frequent consumption of vegetables and fruits, particularly dark green and deep yellow vegetables and cruciferous vegetables (such as cabbage and broccoli), may lower risk for cancers of the lung and bladder as well as some cancers of the alimentary tract. However, the specific components in these foods that may have protective effects have not yet been established. Current evidence suggests the prudence of increasing consumption of whole grain foods and cereals, vegetables (including dried beans and peas), and fruits.

- **Sodium:** Reduce intake of sodium by choosing foods relatively low in sodium and limiting the amount of salt added in food preparation and at the table.

Studies indicate a relationship between a high sodium intake and the occurrence of high blood pressure and stroke. Salt contains about 40 percent sodium by weight and is used widely in the preservation, processing, and preparation of foods. Although sodium is necessary for normal metabolic function, it is consumed in the United States at levels far beyond the 1.1 to 3.3 grams per day found to be as safe and adequate for adults by the National Research Council. Average current sodium intake for adults in the United States is in the range of 4 to 6 grams per day.

Blacks and persons with a family history of high blood pressure are at greater risk for this condition. While some people maintain normal blood pressure levels over a wide range of sodium intake, others appear to be “salt sensitive” and display increased blood pressure in response to high sodium intakes.

Although not all individuals are equally susceptible to the effects of sodium, several observations suggest that it would be prudent for most Americans to reduce sodium intake. These include the lack of a practical biological marker for individual sodium sensitivity, the benefit to persons whose blood pressures do rise with sodium intake, and the lack of harm from moderate sodium restriction.

Processed foods provide about a third or more of dietary sodium. Because about another third of the sodium consumed by Americans is added by the consumer, much can be done to reduce sodium consumption by using less salt at the table and substituting alternative flavoring such as herbs, spices, and lemon juice in the preparation of foods. In addition, choices can be made of foods modified to lower sodium content and less frequent choices could be made of foods to which sodium is added in processing and preservation.

- **Alcohol:** To reduce the risk for chronic disease, take alcohol only in moderation (no more than two drinks a day), if at all. Avoid drinking any alcohol before or while driving, operating machinery, taking medications, or engaging in any other activity requiring judgment. Avoid drinking alcohol while pregnant.

Alcohol is a drug that can produce addiction in susceptible individuals, birth defects in some children born to mothers who drink alcohol during pregnancy, impaired judgment, impaired ability to drive automobiles or operate machinery, and adverse reactions in people taking certain medications. In addition, alcohol abuse has been associated with disrupted family functioning, suicides, and homicides.

Excessive use of alcohol is also associated with liver disease, some types of cancer, high blood pressure, stroke, and disorders of the heart muscle. Extensive epidemiologic and clinical evidence has identified alcohol consumption as the principal cause of liver cirrhosis in the United States, at least in part as a result of the direct toxic effects of alcohol on the liver. Smoking and alcohol appear to act synergistically to increase the risk for cancers of the mouth, larynx, and esophagus. Less conclusive and somewhat conflicting evidence suggests a role of alcohol in other types of cancers such as those of the liver, rectum, breast, and pancreas.

Studies indicate a direct association between increased blood pressure and the consumption of alcohol at levels beyond about two drinks^a daily. Extremely excessive alcohol consumption is associated with cardiomyopathy. Alcohol consumption by the mother during pregnancy has also been associated with fetal malformations.

Although consumption of up to two drinks per day has not been associated with disease among healthy men and nonpregnant women, surveys suggest that at least 9 percent of the total population consumes two or more drinks per day and those in this group need to reduce their alcohol consumption. A threshold level of safety for alcohol intake during pregnancy has not been established. Thus, pregnant women and women who may become pregnant should avoid drinking alcohol.

^aOne drink is defined as a 12 ounce beer, a 5 ounce glass of wine, or 1½ fluid ounces (one jigger) of distilled spirits, each of which contains about 1 ounce of alcohol.

Other Issues for Some People

- **Fluoride:** Community water systems should contain fluoride at optimal levels for prevention of tooth decay. If such water is not available, use other appropriate sources of fluoride.

The most efficient means of making fluoride available to the general public to reduce dental disease is through drinking water. Numerous epidemiologic and clinical studies have attested to the efficacy, safety and cost-effectiveness of systemic fluoride in the prevention of tooth decay. Lifetime use of water containing an optimal fluoride concentration of approximately 1 part per million has been shown to reduce the prevalence of dental caries by more than 50 percent. Water fluoridation is considered one of the most successful public health efforts introduced in the United States.

For children living in areas with inadequate concentrations of fluoride in the water, supplementary fluoride sources should be used at dosages that depend on the fluoride content of the local water supply and the age of the child. The effectiveness of prenatal fluoride administration, however, is uncertain because clinical studies of its effects on subsequent caries incidence have been equivocal. Excessive fluoride should be avoided because it may cause mottling of developing teeth.

- **Sugars:** Those who are particularly vulnerable to dental caries (cavities), especially children, should limit their consumption and frequency of use of foods high in sugars.

Although genetic, behavioral, and other dietary factors also influence dental health, the major role of sugars in promotion of tooth decay is well established from animal, epidemiologic, clinical, and biochemical studies. Newly erupting teeth are generally more vulnerable to decay than mature teeth.

Research has shown that three conditions must exist for the formation of dental caries: the presence of fermentable carbohydrate, acid-producing bacteria, and a susceptible tooth. Caries-producing bacteria metabolize a range of sugars (glucose, fructose, maltose, lactose, and sucrose) to acids that demineralize teeth. The unique role of sucrose (common table sugar) in dental caries is related to its special ability to be converted by these bacteria into long, complex molecules that adhere firmly to teeth and form plaque.

The most important diet-related interventions are fluoridation of drinking water, or the use of other means of fluoride administration, and control of intake of sugars. While fluoride is the most important factor overall in dental caries prevention, reduction in the frequency of consumption and in the quantity of sugar-rich foods in the diet will also help reduce decay. Sticky sweet foods that adhere to the teeth are more cariogenic than those that wash off quickly. The longer cariogenic foods remain in the mouth, the more they are likely to increase the initiation and progression of tooth decay.

- **Calcium:** Adolescent girls and adult women should increase consumption of foods high in calcium, including low-fat dairy products.

Inadequate dietary calcium consumption in the first three to four decades of life may be associated with increased risk for osteoporosis in later life. Osteoporosis, a chronic disease characterized by progressive loss of bone mass with aging, occurs in both women and men, although postmenopausal women are twice as likely as men to have severe osteoporosis with consequent bone fractures. Evidence shows that chronically low calcium intake, especially during adolescence and early adulthood, may compromise development of peak bone mass. In postmenopausal women, the group at highest risk for osteoporosis, estrogen replacement therapy under medical supervision is the most effective means to reduce the rate of bone loss and risk for fractures. Maintenance of adequate levels of physical activity and cessation of cigarette smoking have also been associated with reduced osteoporosis risk.

Although the precise relationship of dietary calcium to osteoporosis has not been elucidated, it appears that higher intakes of dietary calcium could increase peak bone mass during adolescence and delay the onset of bone fractures later in life. Thus, increased consumption of foods rich in calcium may be especially beneficial for adolescents and young women. Food sources of calcium consistent with other dietary recommendations in this Report include low-fat dairy products, some canned fish, certain vegetables, and some calcium-enriched grain products.

- **Iron:** Children, adolescents, and women of childbearing age should be sure to consume foods that are good sources of iron, such as lean meats, fish, certain beans, and iron-enriched cereals and whole grain products. This issue is of special concern for low-income families.

Dietary iron deficiency is responsible for the most prevalent form of anemia in the United States. Iron deficiency hampers the body's ability to produce hemoglobin, a substance needed to carry oxygen in the blood. A

principal consequence of iron deficiency is reduced work capacity, although depressed immune function, changes in behavior, and impaired intellectual performance may also result. Because of the serious consequences of iron deficiency, continual monitoring of the iron status of individuals at high risk—particularly children from low-income families, adolescents, and women of childbearing age—is vital, as is treatment of those identified to be iron deficient.

Proper infant feeding—preferably breastfeeding, otherwise use of iron-fortified formula—is the most important safeguard against iron deficiency in infants. Among adolescents and adults, iron intake can be improved by increasing consumption of iron-rich foods such as lean meats, fish, certain kinds of beans, and iron-enriched cereals and whole grain products. Also, consuming foods that contain vitamin C increases the likelihood that iron will be absorbed efficiently.

Policy Implications

Dietary Guidance

General Public

Educating the public about the dietary choices most conducive to prevention and control of certain chronic diseases is essential. Educational efforts should begin in primary school and continue throughout the secondary grades, and should focus on the dietary principles outlined in this Report—the potential health benefits of eating a diet that is lower in fat (especially saturated fat) and rich in complex carbohydrates and fiber. The importance of adequate physical activity should also be stressed. Efforts should continue throughout each stage of life to promote the principles outlined in the *Dietary Guidelines for Americans*.

Special Populations

A disproportionate burden of diet-related disease is borne by subgroups in our population. Black Americans, for example, have higher rates of high blood pressure, strokes, diabetes, and other diseases associated with obesity (but lower rates of osteoporosis) than the general population. Some groups of Native Americans exhibit the highest rates of diabetes in the world. Pregnant and lactating women also have special nutritional needs. Particular effort should be made to identify and remove the barriers to optimal health and nutritional status in such high-risk groups, using methods that take into consideration their diverse cultural backgrounds.

Many older persons suffer from chronic diseases that can reduce functional independence; many take multiple medications that may adversely interact with nutrients. Sound public education directed toward this group—and professional education directed toward individuals who care for older Americans—should focus on dietary means to reduce risk factors for chronic disease, to promote functional independence, and to prevent adverse consequences of use of medications.

Health Professionals

Improved nutrition training of physicians and other health professionals is needed. Training should emphasize basic principles of nutrition, the role of diet in health promotion and disease prevention, nutrition assessment methodologies and their interpretation, therapeutic aspects of dietary intervention, behavioral aspects of dietary counseling, and the role of dietitians and nutritionists in dietary counseling of patients.

Programs and Services

Food Labels

Food labeling offers opportunities to inform people about the nutrient content of foods so as to facilitate dietary choices most conducive to health. Food manufacturers should be encouraged to make full use of nutrition labels. Labels of processed foods should state the content of calories, protein, carbohydrate, fats, cholesterol, sodium, and vitamins and minerals. To the extent permitted by analytical methods, manufacturers should disclose information where appropriate on the content of saturated and unsaturated fatty acids and total fiber in foods that normally contain them. Descriptive terms such as “low calorie” and “sodium reduced” in compliance with the Food and Drug Administration’s regulations for food labeling may also be helpful, and the expanded use of these terms should be encouraged.

Nutrition Services

Health care programs for individuals of all ages should include nutrition services such as, when appropriate, nutrition counseling for individuals or groups, interpretation and implementation of prescribed therapeutic diets tailored to individual food preferences and lifestyle, referral to appropriate community services and food assistance programs, monitoring of progress, and appropriate followup. These services should routinely incorporate assessment of nutritional status and needs based on established crite-

ria to identify individuals with nutritional risk factors who would profit from preventive measures and those with nutritional disorders who need remedial care.

Food Services

Lack of access to an appropriate diet should not be a health problem for any American. Wherever food is served to people or provided through food assistance programs, it should reflect the principles of good nutrition stated in this Report. Whether served in hospitals, schools, military installations, soup kitchens, day care centers, or nursing homes, or whether delivered to homes, food service programs offer important opportunities for improving health and providing dietary education. Such programs should pay special attention to the nutritional needs of older people, pregnant women, and children, especially those of low income or other special dietary needs. Because a large proportion of the population takes meals in restaurants and convenience food facilities, improvements in the overall nutritional balance of the meals served in such places can be expected to contribute to health benefits.

Food service programs should also take particular care to ensure that special diets lower in fat, especially saturated fat, are provided to people with elevated blood cholesterol, heart disease, or diabetes; that diets low in sodium are provided to individuals with high blood pressure; and that protein-restricted diets are made available to people with end-stage kidney disease.

Food Products

The public would benefit from increased availability of foods and food products low in calories, total fat, saturated fat, cholesterol, sodium, and sugars, but high in a variety of natural forms of fiber and, perhaps, certain minerals and vitamins. Food manufacturers can contribute to improving the quality of the American diet by increasing the availability of palatable, easily prepared food products that will help people to follow the dietary principles outlined here. Because the public is becoming increasingly conscious of the role of nutrition in health, development of such products should also benefit the food industry.

Research and Surveillance

Impressive evidence already links nutrition to chronic disease. However, much more information is needed to continue to identify changes in the

national diet that will lead to better health for the Nation. Gaps in our knowledge of nutrition suggest future research and surveillance needs. Examples are:

- The role of specific dietary factors in the etiology and prevention of chronic diseases.
- The childhood dietary pattern that will best prevent later development of chronic diseases.
- The effects of maternal nutrition on the health of the developing fetus.
- The nutrient and energy requirements of older adults.
- How nutrient requirements translate into healthful dietary patterns.
- The development of biochemical markers of dietary intake to monitor better the effects of dietary intervention.
- The identification of effective educational methods to translate dietary recommendations into appropriate food choices.
- The establishment of a nutrition surveillance system that will enhance the monitoring of population-specific and State-specific trends in the occurrence of nutrition-related risk factors and conditions.



Chapter 1

Introduction and Background

*Power of nutriment reaches to bone and to all the parts of bone, to sinew, to vein, to artery, to muscle, to membrane, to flesh, fat, blood, phlegm, marrow, brain, spinal marrow, the intestines, and all their parts; it reaches also to heat, breath, and moisture.
Hippocrates (460–377 B.C.)*

Introduction

It has long been understood that optimal health depends on adequate nutrition, yet knowledge of the ways in which specific dietary factors affect the risk for disease is incomplete. Dietary deficiencies can be manifested in various ways. A deficient intake of energy or nutrients can lead to protein-energy malnutrition or to classic deficiency diseases such as rickets, pellagra, or iron deficiency anemia. Protein-energy malnutrition and diseases due to deficiencies of various nutrients are prominent causes of premature death and disability in developing countries but, with a few exceptions, appear to have been eliminated in the United States.

When nutrient deficiencies are reported in this country, they are most often observed to be associated with poverty, the additional nutrient requirements of pregnancy or infancy (IOM 1985), the abuse or neglect of children or older persons, or some combination of these factors. They also are observed to result from the restricted food intake that sometimes accompanies aging, alcohol or drug abuse, unusually severe and prolonged injury or illness (including prolonged hospitalization), excessive dieting, or restrictive dietary practices. Thus, pregnant women, young infants, children, older persons, alcohol and drug abusers, and chronically ill and disabled individuals are at greatest risk for malnutrition due to dietary deficiencies, especially if their income is low. Whatever its root cause, inadequate nutrition retards normal growth, lowers resistance to infectious disease, impairs maternal and child health, and may adversely affect the ability to function at peak physical and mental capacity. These issues are discussed in detail in the relevant chapters of this Report.

As problems of nutritional deficiency have diminished in the United States, they have been replaced by problems of dietary imbalance and excess. These imbalances and excesses have contributed to the increased prevalence and severity of chronic diseases that are major causes of death and disability among Americans. Table 1-1 lists the 10 leading causes of death in this country. Among them, five—coronary heart disease and generalized atherosclerosis, stroke, some types of cancer, and diabetes—have been associated with dietary excesses or imbalances, and another three—cirrhosis of the liver, accidents, and suicides—are often the result of excessive alcohol intake. Together, these conditions account for as much as 70 percent of annual deaths among Americans (Collins 1986; NCHS 1986). Dietary excesses or imbalances also have been associated with high blood pressure, obesity, dental diseases, osteoporosis, and, perhaps, kidney and gastrointestinal diseases. Such conditions also contribute to much illness, disability, and death in the United States, and thus to substantial human and economic costs to society.

Table 1-1
Estimated Total Deaths and Percent of Total Deaths for the
10 Leading Causes of Death: United States, 1987

Rank	Cause of Death	Number	Percent of Total Deaths
1 ^a	Heart diseases (Coronary heart disease) (Other heart disease)	759,400 (511,700) (247,700)	35.7 (24.1) (11.6)
2 ^a	Cancers	476,700	22.4
3 ^a	Strokes	148,700	7.0
4 ^b	Unintentional injuries (Motor vehicle) (All others)	92,500 (46,800) (45,700)	4.4 (2.2) (2.2)
5	Chronic obstructive lung diseases	78,000	3.7
6	Pneumonia and influenza	68,600	3.2
7 ^a	Diabetes mellitus	37,800	1.8
8 ^b	Suicide	29,600	1.4
9 ^b	Chronic liver disease and cirrhosis	26,000	1.2
10 ^a	Atherosclerosis	23,100	1.1
. . .	All causes	2,125,100	100.0

^aCauses of death in which diet plays a part.

^bCauses of death in which excessive alcohol consumption plays a part.

Source: National Center for Health Statistics 1988.

Much about the ways in which excessive intake of energy and nutrients might affect health remain to be elucidated. Yet despite uncertainties, much has been learned about diet-disease relationships. This first *Surgeon General's Report on Nutrition and Health* examines the current state of knowledge of associations among dietary patterns, nutrients, and certain disease conditions. Its purpose is to review the available research evidence that relates diet to health to establish a basis for policies that promote dietary means to improve health.

Development and Organization of the Report

This *Surgeon General's Report on Nutrition and Health* has been developed in response to the increasing interest of the public, health professionals, and policy leaders in the role of diet in health promotion and prevention of chronic disease. In preparing the Report, the Public Health Service (PHS) reviewed past and current research related to diet and disease as a basis for examination of the implications for public policies on nutrition education, services, and research.

The Report reviews current knowledge of the influence of dietary factors on specific aspects of health. This first chapter introduces the major themes of the Report in their historical context; it also reviews and synthesizes basic information about essential nutrients in the human diet, the levels of intake required for human health, and American dietary patterns, and it explains the criteria used to examine the various kinds of research studies that are reviewed throughout the Report. Chapters 2 through 14 describe the scientific research that has examined associations between specific dietary factors and selected disease conditions in the United States (Coronary Heart Disease, High Blood Pressure, Cancer, Diabetes, Obesity, Skeletal Diseases, Dental Diseases, Kidney Diseases, Gastrointestinal Diseases, Infections and Immunity, Anemia, Neurologic Disorders, and Behavior). Chapters 15 (Maternal and Child Nutrition) and 16 (Aging) review the special nutritional challenges at especially vulnerable stages of the human life cycle. The interactions between nutrients and alcohol and between nutrients and drugs, and the effects of these substances on human nutritional status, are reviewed in chapters 17 and 18, respectively. The Report closes with a final chapter on dietary fads and frauds.

Most of the chapters follow a common format. Each begins with a brief introduction and a section entitled *Historical Perspective* that is designed to establish a historical context for the area under review. A section on

Significance for Public Health contains information currently available on the incidence, prevalence, and cost to the Nation of each of the conditions under review. Many of the chapters contain a further introductory *Scientific Background* section that summarizes technical information needed to understand the research issues reviewed in the Report.

The major part of the chapters is devoted to a review of *Key Scientific Issues* that summarizes current knowledge about possible associations between dietary factors and disease that are most relevant to public policy. Each chapter closes with a section entitled *Implications for Public Health Policy* that summarizes the significance of the research evidence for dietary guidance and education, nutrition programs and services, and nutrition research and surveillance. Finally, *Literature Cited* provides an extensive list of references to support the scientific findings in each chapter.

Historical Perspective

Throughout history, human societies have observed relationships between the consumption of certain foods and the preservation of good health or the prevention or treatment of diseases. Although the word “diet” occurs frequently in writings attributed to Hippocrates and to Galen, the term “nutrition” did not appear as an English word until the mid-1400’s and was used infrequently until the second half of the 19th century (Todhunter 1973). The modern concept of nutrition—that human life depends on a steady intake of a variety of specific dietary substances in defined amounts—is less than 200 years old.

Development of Nutritional Science

Conditions related to nutritional deficiency, such as beriberi, rickets, or scurvy, were described in very early writings, but the identification of the specific dietary factors required to prevent or treat these conditions began to occur only in the late 18th century and did not approach completion for another 150 years. Some of the major events in this gradual development of nutrition as a science from the earliest records to 1950 are listed in Table 1-2. The chapters of this Report review the great expansion of nutrition research and knowledge that has occurred since then.

The earliest efforts to establish the scientific basis of nutrition are usually attributed to the French chemist Lavoisier, who demonstrated in 1789 that the oxygen breathed in air was consumed in the body to produce carbon dioxide and water, and that this central metabolic process was measurable, variable, and related to both the level of physical activity and the amount of

Table 1-2
Selected Events in the History of Nutritional Science to 1950

c. 1500 B.C.	<i>Papyrus Ebers</i> contains prescription believed to refer to diabetes.
c. 400 B.C.	Hippocrates wrote of relationship of diet to health.
c. 300 B.C.	Beriberi described in ancient Chinese texts.
c. 200 A.D.	Arataeus gave the name diabetes to the condition of "too much passing of urine."
1250	Joinville described scurvy among troops of Louis IX at the siege of Cairo.
1614	Sanctorius published studies relating body weight to food intake.
1650	Glisson described rickets in <i>De Rachitide</i> .
1730	Casal described pellagra, calling it "mal de la rosa."
1747	Lind proved that citrus fruits cure scurvy in first controlled human dietary experiment. Menghini established presence of iron in blood.
1752	Reaumur published experiments on digestion in birds.
1780	Spallanzani produced evidence that digestion was the chemical action of gastric juices.
1789	Lavoisier and Seguin make first measurements relating oxygen consumption to human energy metabolism. Cod liver oil used as treatment for rickets.
1796	Lemon juice officially introduced in British Navy to prevent scurvy.
1807	Davy isolated sodium, potassium, calcium, magnesium, sulphur, and boron.
1810–23	Chevreul studied chemistry of animal fats.
1810	Wollaston isolated cystic oxide (later named cystine) from urine—first amino acid discovered.
1816	Magendie identified dietary nitrogen requirements in dogs.
1827	Prout classified food constituents as saccharine, oily, and albuminous (sugar, fat, and protein).
1833	Beaumont reported observations and experiments on digestion in his patient St. Martin.
1838	Mulder introduced the term "protein."
1839	Boussingault conducts first nitrogen balance studies in animals.
1840	Liebig published <i>Animal Chemistry</i> , stating basic principles of metabolism.