

TABLE 8

NUMBER OF REPORTED SHIGELLA ISOLATES BY REGION
FOR 1982-1996

REGION	YEAR																TOTAL
	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996		
NEW ENGLAND	1226	543	354	861	729	645	651	546	646	1757	555	520	414	586	373	10406	
MID-ATLANTIC	1226	2050	1094	1607	1263	2331	1179	1310	1595	1263	1071	1215	1562	1758	2240	22764	
EAST NORTH CENTRAL	1484	1881	1863	1494	1550	3816	3597	2673	2096	1643	2154	2816	1970	2105	1190	32332	
WEST NORTH CENTRAL	402	530	495	387	330	671	1151	1359	630	403	589	828	1368	1391	699	11233	
SOUTH ATLANTIC	1084	990	997	683	686	1425	2562	2568	1894	1565	1727	2860	4790	2181	1912	27924	
EAST SOUTH CENTRAL	131	155	114	188	60	219	683	482	799	915	305	768	1023	802	334	6978	
WEST SOUTH CENTRAL	1776	1619	1263	1218	1611	1671	2535	1408	1832	1409	1453	2943	1771	1944	1225	25678	
MOUNTAIN	1356	1525	1150	1223	1137	1174	1692	1161	1655	2248	1174	1348	1436	2713	1665	22657	
PACIFIC	4852	5664	5385	4215	4906	4812	8286	4465	6159	5302	5895	6213	4449	5850	4433	80886	
TOTAL	13537	14957	12715	11876	12272	16764	22336	15972	17306	16505	14923	19511	18783	19330	14071	240858	

TABLE 8A

NUMBER OF REPORTED *S. SONNEI* ISOLATES BY REGION
FOR 1982-1996

REGION	YEAR																TOTAL
	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996		
NEW ENGLAND	1050	383	214	713	596	491	481	358	448	1573	435	428	302	456	264	8192	
MID-ATLANTIC	941	1629	823	1128	772	2030	883	1015	1321	1020	880	985	1325	1527	2048	18327	
EAST NORTH CENTRAL	1099	1470	1458	1110	1123	3279	2685	2061	1443	1176	1671	2440	1707	1773	896	25391	
WEST NORTH CENTRAL	294	414	354	246	147	503	870	1084	446	255	478	698	1166	1281	575	8811	
SOUTH ATLANTIC	750	705	704	450	462	1099	2155	2268	1535	1210	1451	2280	2695	1966	1772	21502	
EAST SOUTH CENTRAL	109	124	103	179	52	136	523	319	513	587	152	341	649	511	232	4530	
WEST SOUTH CENTRAL	1101	1071	767	817	1161	1293	2103	1022	1415	1072	1268	2765	1595	1721	1085	20256	
MOUNTAIN	654	654	575	554	571	470	921	522	1124	1601	722	824	920	2180	1115	13407	
PACIFIC	2236	2825	2792	1878	2421	1981	4407	1593	2871	2240	3049	3578	2087	3396	2275	39629	
TOTAL	8234	9275	7790	7075	7305	11282	15028	10242	11116	10734	10106	14339	12446	14811	10262	160045	

TABLE 8B

NUMBER OF REPORTED S. FLEXNERI ISOLATES BY REGION
FOR 1982-1996

REGION	YEAR																TOTAL
	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1996	
NEW ENGLAND	139	142	114	128	93	109	127	150	122	94	107	92	106	115	94	1732	
MID-ATLANTIC	271	335	256	377	341	278	257	269	254	227	177	211	213	206	179	3851	
EAST NORTH CENTRAL	342	345	344	326	314	361	422	439	465	339	330	287	238	289	267	5108	
WEST NORTH CENTRAL	82	90	103	108	168	134	156	168	125	115	77	67	121	71	105	1690	
SOUTH ATLANTIC	121	93	102	90	116	154	143	101	156	124	137	173	343	196	122	2171	
EAST SOUTH CENTRAL	19	30	10	8	6	9	8	9	10	8	2	4	18	7	7	155	
WEST SOUTH CENTRAL	618	491	435	346	380	328	376	333	367	280	170	136	141	188	113	4702	
MOUNTAIN	617	774	509	548	496	556	503	472	381	491	371	382	353	464	441	7358	
PACIFIC	1964	1924	1879	1602	1653	1831	2346	1741	2151	2034	1879	1709	1568	1483	1376	27140	
TOTAL	4173	4224	3752	3533	3567	3760	4338	3682	4031	3712	3250	3061	3101	3019	2704	53907	

TABLE 8C

NUMBER OF REPORTED S. BOYDII ISOLATES BY REGION
FOR 1982-1996

REGION	YEAR																TOTAL
	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996		
NEW ENGLAND	9	7	10	8	6	14	7	16	17	12	4	.	5	11	6	132	
MID-ATLANTIC	6	8	8	16	16	16	21	17	10	10	6	11	15	12	5	177	
EAST NORTH CENTRAL	17	42	38	35	37	28	64	42	29	20	16	30	19	31	19	467	
WEST NORTH CENTRAL	4	6	4	1	1	2	6	5	5	7	3	.	7	6	3	60	
SOUTH ATLANTIC	10	11	2	3	6	7	5	10	8	8	5	10	27	7	10	129	
EAST SOUTH CENTRAL	1	1	1	.	1	2	2	1	1	.	1	.	.	.	2	13	
WEST SOUTH CENTRAL	49	47	35	33	54	36	39	37	35	44	11	33	24	27	19	523	
MOUNTAIN	46	48	34	66	43	59	52	48	47	30	14	14	18	32	91	642	
PACIFIC	152	246	254	196	209	192	276	290	246	181	164	123	92	103	120	2844	
TOTAL	294	416	386	358	373	356	472	466	398	312	224	221	207	229	275	4987	

TABLE 8D

NUMBER OF REPORTED *S. DYSENTERIAE* ISOLATES BY REGION
FOR 1982-1996

REGION	YEAR																TOTAL
	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996		
NEW ENGLAND	9	2	6	5	4	4	12	9	8	6	6	.	.	3	6	80	
MID-ATLANTIC	6	7	7	16	11	7	18	9	9	6	8	2	6	6	6	124	
EAST NORTH CENTRAL	16	22	20	20	7	27	21	23	15	11	21	4	6	11	8	232	
WEST NORTH CENTRAL	3	6	2	3	3	8	8	7	3	1	6	1	2	1	2	56	
SOUTH ATLANTIC	3	15	4	1	2	4	4	2	5	6	4	4	5	2	8	69	
EAST SOUTH CENTRAL	.	.	.	1	1	1	2	.	2	.	1	1	1	.	.	10	
WEST SOUTH CENTRAL	8	8	26	20	13	12	17	15	14	10	2	9	10	5	7	176	
MOUNTAIN	14	20	19	14	9	10	8	25	7	12	5	12	9	6	18	188	
PACIFIC	72	105	95	129	68	159	121	126	118	93	73	72	55	56	48	1390	
TOTAL	131	185	179	209	118	232	211	216	181	145	126	105	94	90	103	2325	

TABLE 8E

NUMBER OF REPORTED SHIGELLA ISOLATES WITH UNKNOWN SEROGROUP
BY REGION FOR 1982-1996

REGION	YEAR																TOTAL
	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996		
NEW ENGLAND	19	9	10	7	30	27	24	13	51	72	3	.	1	1	3	270	
MID-ATLANTIC	2	71	.	70	123	.	.	.	1	.	.	6	3	7	2	285	
EAST NORTH CENTRAL	10	2	3	3	69	121	405	108	144	97	116	55	.	1	.	1134	
WEST NORTH CENTRAL	19	14	32	29	11	24	111	95	51	25	25	62	72	32	14	616	
SOUTH ATLANTIC	200	166	185	139	100	161	255	187	190	217	130	393	1720	10	.	4053	
EAST SOUTH CENTRAL	2	71	148	153	273	320	149	422	355	284	93	2270	
WEST SOUTH CENTRAL	.	2	.	2	3	2	.	1	1	3	2	.	1	3	1	21	
MOUNTAIN	25	29	13	41	18	79	208	94	96	114	62	116	136	31	.	1062	
PACIFIC	428	564	365	410	555	649	1136	715	773	754	730	731	647	812	614	9883	
TOTAL	705	857	608	701	909	1134	2287	1366	1580	1602	1217	1785	2935	1181	727	19594	