

TABLE 2

Laboratory confirmed *Shigella* isolates reported to the CDC by Serotype in 2003

Rank	Serotype	Reported	Percent
1	<i>S. sonnei</i>	9263	80.2
2	<i>S. flexneri</i> unspecified	813	7.0
3	<i>S. flexneri</i> 2 unspecified	186	1.6
4	<i>S. flexneri</i> 3 unspecified	112	1.0
5	<i>S. flexneri</i> 1 unspecified	100	0.9
6	<i>S. flexneri</i> 2a	92	0.8
7	<i>S. flexneri</i> 3a	79	0.7
8	<i>S. boydii</i> unspecified	68	0.6
9	<i>S. flexneri</i> 4 unspecified	61	0.5
10	<i>S. flexneri</i> 6	58	0.5
11	<i>S. flexneri</i> 4a	49	0.4
12	<i>S. flexneri</i> 1b	31	0.3
13	<i>S. boydii</i> 2	29	0.3
14	<i>S. dysenteriae</i> unspecified	20	0.2
15	<i>S. flexneri</i> 2b	17	0.2
16	<i>S. flexneri</i> variant y	15	0.1
17	<i>S. flexneri</i> 3b	12	0.1
18	<i>S. boydii</i> 1	10	0.1
19	<i>S. dysenteriae</i> 2	10	0.1
20	<i>S. flexneri</i> 5 unspecified	10	0.1
21	<i>S. boydii</i> 4	9	0.1
22	<i>S. flexneri</i> 1a	6	0.1
23	<i>S. flexneri</i> 4b	6	0.1
24	<i>S. flexneri</i> variant x	6	0.1
25	<i>S. dysenteriae</i> 1	5	0.0
26	<i>S. flexneri</i> 88-893	5	0.0
27	<i>S. boydii</i> 10	2	0.0
28	<i>S. boydii</i> 8	2	0.0
29	<i>S. dysenteriae</i> 3	2	0.0
30	<i>S. dysenteriae</i> 4	2	0.0
31	<i>S. boydii</i> 12	1	0.0
32	<i>S. boydii</i> 14	1	0.0
33	<i>S. boydii</i> 15	1	0.0
34	<i>S. boydii</i> 20	1	0.0
35	<i>S. boydii</i> 5	1	0.0
36	<i>S. dysenteriae</i> 14	1	0.0
37	<i>S. dysenteriae</i> 9	1	0.0
38	<i>S. flexneri</i> 4c	1	0.0
39	<i>S. flexneri</i> 5a	1	0.0
	Sub Total	11089	96.0
	Unknown	463	4.0
	Sub Total	463	4.0
	Total	11552	100.0