

Table 3-334. Estimates of lost-time injuries by farm type for white workers in the Northeast region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	595.8	4.4	595.8	4.4
Field crops	1724.8	12.9	2320.6	17.3
Veg, fruit, nuts	741.7	5.5	3062.3	22.9
Nursery	1371.5	10.2	4433.8	33.1
Beef, hogs, sheep	1758.1	13.1	6191.9	46.2
Poultry	150.6	1.1	6342.5	47.3
Dairy	5896.8	44.0	12239.3	91.3
Other farms	1161.9	8.7	13401.2	100.0

Table 3-335. Estimates of lost-time injuries by status of the injured worker for white workers in the Northeast region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	10040.7	74.9	10040.7	74.9
Hired	3246.9	24.2	13287.6	99.2
Other	113.6	0.8	13401.2	100.0

Table 3-336. Estimates of lost-time injuries by gender for white workers in the Northeast region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	9961.2	74.3	9961.2	74.3
Female	3440.0	25.7	13401.2	100.0

Table 3-337. Estimates of lost-time injuries by age group for white workers in the Northeast region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	1283.8	9.6	1283.8	9.6
20-29	3704.6	27.6	4988.4	37.2
30-39	1324.5	9.9	6312.9	47.1
40-49	2581.5	19.3	8894.4	66.4
50-59	2229.3	16.6	11123.7	83.0
60-69	1329.0	9.9	12452.7	92.9
>=70	948.5	7.1	13401.2	100.0

Table 3-338. Estimates of lost-time injuries by month for white workers in the Northeast region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	153.5	1.1	153.5	1.1
February	667.6	5.0	821.1	6.1
March	397.6	3.0	1218.7	9.1
April	1381.4	10.3	2600.1	19.4
May	1414.8	10.6	4014.9	30.0
June	746.1	5.6	4761.0	35.5
July	2967.1	22.1	7728.1	57.7
August	1034.9	7.7	8763.0	65.4
September	1347.1	10.1	10110.1	75.4
October	977.7	7.3	11087.8	82.7
November	1231.5	9.2	12319.3	91.9
December	1081.9	8.1	13401.2	100.0

Table 3-339. Estimates of lost-time injuries by severity of the injury for white workers in the Northeast region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	12933.0	96.5	12933.0	96.5
Permanent	468.2	3.5	13401.2	100.0

Table 3-340. Estimates of lost-time injuries by whether professional medical attention was sought for white workers in the Northeast region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	468.2	3.5	468.2	3.5
Yes	10644.0	79.4	11112.2	82.9
No	2289.0	17.1	13401.2	100.0

Table 3-341. Estimates of lost-time injuries by body part injured for white workers in the Northeast region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	705.7	5.3	705.7	5.3
Eye	1002.4	7.5	1708.1	12.7
Chest/trunk	666.0	5.0	2374.1	17.7
Back	840.4	6.3	3214.5	24.0
Arm/shoulder	957.3	7.1	4171.8	31.1
Finger	2842.7	21.2	7014.5	52.3
Hand/wrist	3272.6	24.4	10287.1	76.8
Leg/knee/hip	1379.9	10.3	11667.0	87.1
Foot	1197.1	8.9	12864.1	96.0
Mult. parts	178.5	1.3	13042.6	97.3
Other	358.6	2.7	13401.2	100.0

Table 3-242. Estimates of lost-time injuries by nature of injury for white workers in the Northeast region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	203.9	1.5	203.9	1.5
Bruise	2165.2	16.2	2369.1	17.7
Cut	2869.5	21.4	5238.6	39.1
Crush	489.1	3.6	5727.7	42.7
Fracture	2432.3	18.1	8160.0	60.9
Puncture	389.7	2.9	8549.7	63.8
Sprain/strain	3735.1	27.9	12284.8	91.7
Mult. injury	113.6	0.8	12398.4	92.5
Other	1002.8	7.5	13401.2	100.0

Table 3-243. Estimates of lost-time injuries by type of event for white workers in the Northeast region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	525.6	3.9	525.6	3.9
Caught between	1815.1	13.5	2340.7	17.5
Caught under	309.2	2.3	2649.9	19.8
Struck by/against	1204.5	9.0	3854.4	28.8
Struck by falling	620.3	4.6	4474.7	33.4
Struck by flying	838.4	6.3	5313.1	39.6
Contact sharp obj.	2428.2	18.1	7741.3	57.8
Fall, same level	1781.0	13.3	9522.3	71.1
Fall, elevation	1084.5	8.1	10606.8	79.1
Overexertion	1486.0	11.1	12092.8	90.2
Other	1308.4	9.8	13401.2	100.0

Table 3-344. Estimates of lost-time injuries by activity when worker was injured for white workers in the Northeast region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	1982.0	14.8	1982.0	14.8
Machine maint.	2237.2	16.7	4219.2	31.5
Field work	2492.5	18.6	6711.7	50.1
Crop handling	2253.3	16.8	8965.0	66.9
Livestock handling	3306.3	24.7	12271.3	91.6
Other	1129.9	8.4	13401.2	100.0

Table 3-345. Estimates of lost-time injuries by source of the injury for white workers in the Northeast region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	316.9	2.4	316.9	2.4
Machinery	2589.4	19.3	2906.3	21.7
Livestock	2656.2	19.8	5562.5	41.5
Hand tool	427.9	3.2	5990.4	44.7
Power tool	1465.2	10.9	7455.6	55.6
Plant/tree	326.6	2.4	7782.2	58.1
Working surface	1413.9	10.6	9196.1	68.6
Truck/auto	471.1	3.5	9667.2	72.1
Other vehicle	69.3	0.5	9736.5	72.7
Other	3664.7	27.3	13401.2	100.0

Table 3-346. Estimates of lost-time injuries by farm type for Hispanic workers in the Northeast region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Field crops	152.3	15.1	152.3	15.1
Veg, fruit, nuts	756.9	74.9	909.2	90.0
Nursery	36.1	3.6	945.3	93.6
Beef, hogs, sheep	64.9	6.4	1010.2	100.0

Table 3-347. Estimates of lost-time injuries by status of the injured worker for Hispanic workers in the Northeast region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	735.6	72.8	735.6	72.8
Hired	274.6	27.2	1010.2	100.0

Table 3-348. Estimates of lost-time injuries by gender for Hispanic workers in the Northeast region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	992.5	98.2	992.5	98.2
Female	17.7	1.8	1010.2	100.0

Table 3-349. Estimates of lost-time injuries by age group for Hispanic workers in the Northeast region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
30-39	409.2	40.5	409.2	40.5
50-59	17.7	1.8	426.9	42.3
>=70	583.3	57.7	1010.2	100.0

Table 3-350. Estimates of lost-time injuries by month for Hispanic workers in the Northeast region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	0.0	0.0	0.0	0.0
February	0.0	0.0	0.0	0.0
March	0.0	0.0	0.0	0.0
April	64.9	6.4	64.9	6.4
May	86.8	8.6	151.7	15.0
June	0.0	0.0	151.7	15.0
July	0.0	0.0	151.7	15.0
August	0.0	0.0	151.7	15.0
September	170.7	16.9	322.4	31.9
October	670.1	66.3	992.5	98.2
November	0.0	0.0	992.5	98.2
December	17.7	1.8	1010.2	100.0

Table 3-351. Estimates of lost-time injuries by severity of the injury for Hispanic workers in the Northeast region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	1010.2	100.0	1010.2	100.0

Table 3-352. Estimates of lost-time injuries by whether professional medical attention was sought for Hispanic workers in the Northeast region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Yes	1010.2	100.0	1010.2	100.0

Table 3-353. Estimates of lost-time injuries by body part injured for Hispanic workers in the Northeast region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	64.9	6.4	64.9	6.4
Eye	36.1	3.6	101.0	10.0
Finger	134.6	13.3	235.6	23.3
Hand/wrist	583.3	57.7	818.9	81.1
Leg/knee/hip	86.8	8.6	905.7	89.7
Toe	17.7	1.8	923.4	91.4
Foot	86.8	8.6	1010.2	100.0

Table 3-354. Estimates of lost-time injuries by nature of injury for Hispanic workers in the Northeast region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Bruise	64.9	6.4	64.9	6.4
Cut	36.1	3.6	101.0	10.0
Crush	134.6	13.3	235.6	23.3
Fracture	86.8	8.6	322.4	31.9
Puncture	86.8	8.6	409.2	40.5
Other	601.0	59.5	1010.2	100.0

Table 3-355. Estimates of lost-time injuries by type of event for Hispanic workers in the Northeast region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	86.8	8.6	86.8	8.6
Caught between	134.6	13.3	221.4	21.9
Struck by/against	36.1	3.6	257.5	25.5
Struck by falling	64.9	6.4	322.4	31.9
Contact sharp obj.	86.8	8.6	409.2	40.5
Other	601.0	59.5	1010.2	100.0

Table 3-356. Estimates of lost-time injuries by activity when worker was injured for Hispanic workers in the Northeast region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	64.9	6.4	64.9	6.4
Machine maint.	134.6	13.3	199.5	19.7
Field work	209.7	20.8	409.2	40.5
Crop handling	583.3	57.7	992.5	98.2
Livestock handling	17.7	1.8	1010.2	100.0

Table 3-357. Estimates of lost-time injuries by source of the injury for Hispanic workers in the Northeast region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Machinery	221.4	21.9	221.4	21.9
Plant/tree	36.1	3.6	257.5	25.5
Working surface	64.9	6.4	322.4	31.9
Other	687.8	68.1	1010.2	100.0

Table 3-358. Estimates of lost-time injuries by farm type for white workers in the South region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	2778.4	7.7	2778.4	7.7
Field crops	5546.8	15.3	8325.2	23.0
Veg, fruit, nuts	2286.1	6.3	10611.3	29.3
Nursery	1216.7	3.4	11828.0	32.6
Beef, hogs, sheep	19062.7	52.6	30890.7	85.2
Poultry	570.0	1.6	31460.7	86.8
Dairy	3695.6	10.2	35156.3	97.0
Other farms	1097.0	3.0	36253.3	100.0

Table 3-359. Estimates of lost-time injuries by status of the injured worker for white workers in the South region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	27270.3	75.2	27270.3	75.2
Hired	8983.0	24.8	36253.3	100.0

Table 3-360. Estimates of lost-time injuries by gender for white workers in the South region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	32191.8	88.8	32191.8	88.8
Female	4061.5	11.2	36253.3	100.0

Table 3-361. Estimates of lost-time injuries by age group for white workers in the South region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	1056.7	2.9	1056.7	2.9
20-29	3595.6	9.9	4652.3	12.8
30-39	8393.4	23.2	13045.7	36.0
40-49	6880.2	19.0	19925.9	55.0
50-59	6386.8	17.6	26312.7	72.6
60-69	5757.8	15.9	32070.5	88.5
>=70	4182.8	11.5	36253.3	100.0

Table 3-362. Estimates of lost-time injuries by month for white workers in the South region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	2872.7	7.9	2872.7	7.9
February	3975.7	11.0	6848.4	18.9
March	1583.9	4.4	8432.3	23.3
April	2942.8	8.1	11375.1	31.4
May	2772.9	7.6	14148.0	39.0
June	4935.4	13.6	19083.4	52.6
July	1340.7	3.7	20424.1	56.3
August	3605.9	9.9	24030.0	66.3
September	3022.0	8.3	27052.0	74.6
October	1690.8	4.7	28742.8	79.3
November	3760.2	10.4	32503.0	89.7
December	3750.3	10.3	36253.3	100.0

Table 3-363. Estimates of lost-time injuries by severity of the injury for white workers in the South region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	31769.3	87.6	31769.3	87.6
Permanent	4111.4	11.3	35880.7	99.0
Fatality	372.6	1.0	36253.3	100.0

Table 3-364. Estimates of lost-time injuries by whether professional medical attention was sought for white workers in the South region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	4484.0	12.4	4484.0	12.4
Yes	26398.0	72.8	30882.0	85.2
No	5371.3	14.8	36253.3	100.0

Table 3-365. Estimates of lost-time injuries by body part injured for white workers in the South region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	1953.8	5.4	1953.8	5.4
Eye	1056.5	2.9	3010.3	8.3
Chest/trunk	5422.3	15.0	8432.6	23.3
Back	3351.6	9.2	11784.2	32.5
Arm/shoulder	2527.8	7.0	14312.0	39.5
Finger	3477.3	9.6	17789.3	49.1
Hand/wrist	3779.1	10.4	21568.4	59.5
Leg/knee/hip	7340.5	20.2	28908.9	79.7
Toe	964.8	2.7	29873.7	82.4
Foot	4087.3	11.3	3961.0	93.7
Mult. parts	2167.5	6.0	36128.5	99.7
Other	124.8	0.3	36253.3	100.0

Table 3-366. Estimates of lost-time injuries by nature of injury for white workers in the South region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	398.6	1.1	398.6	1.1
Bruise	5128.0	14.1	5526.6	15.2
Burn	511.1	1.4	6037.7	16.7
Cut	6023.3	16.6	12061.0	33.3
Crush	4365.8	12.0	16426.8	45.3
Fracture	9986.9	27.5	26413.7	72.9
Puncture	440.5	1.2	26854.2	74.1
Sprain/strain	4302.3	11.9	31156.5	85.9
Mult. injury	2671.2	7.4	33827.7	93.3
Other	2425.6	6.7	36253.3	100.0

Table 3-367. Estimates of lost-time injuries by type of event for white workers in the South region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	1275.0	3.5	1275.0	3.5
Caught between	2729.6	7.5	4004.6	11.0
Caught under	950.6	2.6	4955.2	13.7
Struck by/against	10778.2	29.7	15733.4	43.4
Struck by falling	2824.2	7.8	18557.6	51.2
Struck by flying	905.3	2.5	19462.9	53.7
Contact sharp obj.	4254.6	11.7	23717.5	65.4
Fall, same level	5048.5	13.9	28766.0	79.3
Fall, elevation	3877.1	10.7	32643.1	90.0
Overexertion	1399.1	3.9	34042.2	93.9
Other	2211.1	6.1	36253.3	100.0

Table 3-368. Estimates of lost-time injuries by activity when worker was injured for white workers in the South region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	6486.3	17.9	6486.3	17.9
Machine maint.	4024.1	11.1	10510.4	29.0
Field work	5731.6	15.8	16242.0	44.8
Crop handling	6103.6	16.8	22345.6	61.6
Livestock handling	13489.9	37.2	35835.5	98.8
Other	417.8	1.2	36253.3	100.0

Table 3-369. Estimates of lost-time injuries by source of the injury for family workers in the Northeast region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	4009.0	11.1	4009.0	11.1
Machinery	4098.7	11.3	8107.7	22.4
Livestock	9663.0	26.7	17770.7	49.0
Hand tool	2798.8	7.7	20569.5	56.7
Power tool	805.9	2.2	21375.4	59.0
Pesticide/chem.	886.1	2.4	22261.5	61.4
Plant/tree	1768.6	4.9	24030.1	66.3
Working surface	3300.4	9.1	27330.5	75.4
Truck/auto	1178.2	3.2	28508.7	78.6
Other vehicle	703.9	1.9	29212.6	80.6
Other	7040.7	19.4	36253.3	100.0

Table 3-370. Estimates of lost-time injuries by farm type for Hispanic workers in the South region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Field crops	1517.1	26.8	1517.1	26.8
Veg, fruit, nuts	2354.2	41.6	3871.3	68.5
Nursery	744.0	13.2	4615.3	81.6
Beef, hogs, sheep	277.5	4.9	4892.8	86.5
Dairy	367.7	6.5	5260.5	93.1
Other farms	392.7	6.9	5653.2	100.0

Table 3-371. Estimates of lost-time injuries by status of the injured worker for Hispanic workers in the South region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Hired	5375.7	95.1	5375.7	95.1
Other	277.5	4.9	5653.2	100.0

Table 3-372. Estimates of lost-time injuries by gender for Hispanic workers in the South region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	5355.6	94.7	5355.6	94.7
Female	297.6	5.3	5653.2	100.0

Table 3-373. Estimates of lost-time injuries by age group for Hispanic workers in the South region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	337.5	6.0	337.5	6.0
20-29	3467.8	61.3	3805.3	67.3
30-39	1446.8	25.6	5252.1	92.9
40-49	148.8	2.6	5400.9	95.5
50-59	252.3	4.5	5653.2	100.0

Table 3-374. Estimates of lost-time injuries by month for Hispanic workers in the South region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	0.0	0.0	0.0	0.0
February	0.0	0.0	0.0	0.0
March	277.5	4.9	277.5	4.9
April	148.8	2.6	426.3	7.5
May	0.0	0.0	426.3	7.5
June	516.5	9.1	942.8	16.7
July	905.3	16.0	1848.1	32.7
August	1076.1	19.0	2924.2	51.7
September	148.8	2.6	3073.0	54.4
October	2580.2	45.6	5653.2	100.0
November	0.0	0.0	5653.2	100.0
December	0.0	0.0	5653.2	100.0

Table 3-375. Estimates of lost-time injuries by severity of the injury for Hispanic workers in the South region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	5653.2	100.0	5653.2	100.0

Table 3-376. Estimates of lost-time injuries by whether professional medical attention was sought for Hispanic workers in the South region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Yes	5653.2	100.0	5653.2	100.0

Table 3-377. Estimates of lost-time injuries by body part injured for Hispanic workers in the South region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	252.3	4.5	252.3	4.5
Eye	166.7	2.9	419.0	7.4
Back	1637.6	29.0	2056.6	36.4
Finger	148.8	2.6	2205.4	39.0
Hand/wrist	879.0	15.5	3084.4	54.6
Leg/knee/hip	252.3	4.5	3336.7	59.0
Foot	1830.2	32.4	5166.9	91.4
Mult. parts	486.3	8.6	5653.2	100.0

Table 3-378. Estimates of lost-time injuries by nature of injury for Hispanic workers in the South region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Bruise	277.5	4.9	277.5	4.9
Cut	2107.5	37.3	2385.0	42.2
Fracture	516.5	9.1	2901.5	51.3
Puncture	337.5	6.0	3239.0	57.3
Sprain/strain	1612.4	28.5	4851.4	85.8
Mult. injury	148.8	2.6	5000.2	88.4
Other	653.0	11.6	5653.2	100.0

Table 3-379. Estimates of lost-time injuries by type of event for Hispanic workers in the South region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught between	367.7	6.5	367.7	6.5
Caught under	337.5	6.0	705.2	12.5
Struck by/against	670.2	11.9	1375.4	24.3
Struck by falling	1377.3	24.4	2752.7	48.7
Contact sharp obj.	337.5	6.0	3090.2	54.7
Fall, same level	297.6	5.3	3387.8	59.9
Overexertion	1949.9	34.5	5337.7	94.4
Other	315.5	5.6	5653.2	100.0

Table 3-380. Estimates of lost-time injuries by activity when worker was injured for Hispanic workers in the South region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	1273.8	22.5	1273.8	22.5
Field work	2895.1	51.2	4168.9	73.7
Crop handling	446.4	7.9	4615.3	81.6
Livestock handling	1037.9	18.4	5653.2	100.0

Table 3-381. Estimates of lost-time injuries by source of the injury for Hispanic workers in the South region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Machinery	3011.3	53.3	3011.3	53.3
Livestock	670.2	11.9	3681.5	65.1
Plant/tree	297.6	5.3	3979.1	70.4
Working surface	549.9	9.7	4529.0	80.1
Other vehicle	367.7	6.5	4896.7	86.6
Other	756.5	13.4	5653.2	100.0

Table 3-382. Estimates of lost-time injuries by farm type for black and other race workers in the South region of the U.S., 1993.

<u>Farm Type</u>	<u>Frequency</u>	<u>Percent</u>	<u>Cumulative Frequency</u>	<u>Cumulative Percent</u>
Cash grain	306.8	15.0	306.8	15.0
Field crops	124.8	6.1	431.6	21.1
Veg, fruit, nuts	266.1	13.0	697.7	34.2
Nursery	260.9	12.8	958.6	46.9
Beef, hogs, sheep	398.6	19.5	1357.2	66.4
Poultry	385.7	18.9	1742.9	85.3
Other farms	300.0	14.7	2042.9	100.0

Table 3-383. Estimates of lost-time injuries by status of the injured worker for black and other race workers in the South region of the U.S., 1993.

<u>Status of Injured Worker</u>	<u>Frequency</u>	<u>Percent</u>	<u>Cumulative Frequency</u>	<u>Cumulative Percent</u>
Family	385.7	18.9	385.7	18.9
Hired	1657.2	81.1	2042.9	100.0

Table 3-384. Estimates of lost-time injuries by gender for black and other race workers in the South region of the U.S., 1993.

<u>Gender</u>	<u>Frequency</u>	<u>Percent</u>	<u>Cumulative Frequency</u>	<u>Cumulative Percent</u>
Male	2042.9	100.0	2042.9	100.0

Table 3-385. Estimates of lost-time injuries by age group for black and other race workers in the South region of the U.S., 1993.

<u>Age Group (Years)</u>	<u>Frequency</u>	<u>Percent</u>	<u>Cumulative Frequency</u>	<u>Cumulative Percent</u>
20-29	664.7	32.5	664.7	32.5
30-39	567.7	27.8	1232.4	60.3
50-59	124.8	6.1	1357.2	66.4
60-69	385.7	18.9	1742.9	85.3
>=70	300.0	14.7	2042.9	100.0

Table 3-386. Estimates of lost-time injuries by month for black and other race workers in the South region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	0.0	0.0	0.0	0.0
February	0.0	0.0	0.0	0.0
March	266.1	13.0	266.1	13.0
April	0.0	0.0	266.1	13.0
May	153.4	7.5	419.5	20.5
June	0.0	0.0	419.5	20.5
July	153.4	7.5	572.9	28.0
August	0.0	0.0	572.9	28.0
September	523.4	25.6	1096.3	53.7
October	0.0	0.0	1096.3	53.7
November	0.0	0.0	1096.3	53.7
December	946.6	46.3	2042.9	100.0

Table 3-387. Estimates of lost-time injuries by severity of the injury for black and other race workers in the South region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	2042.9	100.0	2042.9	100.0

Table 3-388. Estimates of lost-time injuries by whether professional medical attention was sought for black and other race workers in the South region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Yes	1396.3	68.3	1396.3	68.3
No	646.6	31.7	2042.9	100.0

Table 3-389. Estimates of lost-time injuries by body part injured for black and other race workers in the South region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Eye	398.6	19.5	398.6	19.5
Back	572.9	28.0	971.5	47.6
Finger	124.8	6.1	1096.3	53.7
Hand/wrist	260.9	12.8	1357.2	66.4
Leg/knee/hip	300.0	14.7	1657.2	81.1
Foot	385.7	18.9	2042.9	100.0

Table 3-390. Estimates of lost-time injuries by nature of injury for black and other race workers in the South region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Bruise	800.0	39.2	800.0	39.2
Cut	523.4	25.6	1323.4	64.8
Puncture	300.0	14.7	1623.4	79.5
Sprain/strain	419.5	20.5	2042.9	100.0

Table 3-391. Estimates of lost-time injuries by type of event for black and other race workers in the South region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught between	385.7	18.9	385.7	18.9
Struck by/against	552.0	27.0	937.7	45.9
Struck by falling	385.7	18.9	1323.4	64.8
Contact sharp obj.	300.0	14.7	1623.4	79.5
Fall, same level	266.1	13.0	1889.5	92.5
Overexertion	153.4	7.5	2042.9	100.0

Table 3-392. Estimates of lost-time injuries by activity when worker was injured for black and other race workers in the South region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	685.7	33.6	685.7	33.6
Machine maint.	390.9	19.1	1076.6	52.7
Crop handling	567.7	27.8	1644.3	80.5
Other	398.6	19.5	2042.9	100.0

Table 3-393. Estimates of lost-time injuries by source of the injury for black and other race workers in the South region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	385.7	18.9	385.7	18.9
Machinery	453.4	22.2	839.1	41.1
Hand tool	390.9	19.1	1230.0	60.2
Plant/tree	398.6	19.5	1628.6	79.7
Working surface	260.9	12.8	1889.5	92.5
Truck/auto	153.4	7.5	2042.9	100.0

Table 3-394. Estimates of lost-time injuries by farm type for white workers in the Midwest region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	23294.4	29.1	23294.4	29.1
Field crops	2731.4	3.4	26025.8	32.5
Veg, fruit, nuts	1145.6	1.4	27171.4	33.9
Nursery	1749.5	2.2	28920.9	36.1
Beef, hogs, sheep	30282.3	37.8	59203.2	73.8
Poultry	80.0	0.1	59283.2	73.9
Dairy	20263.6	25.3	79546.8	99.2
Other farms	640.0	0.8	80186.8	100.0

Table 3-395. Estimates of lost-time injuries by status of the injured worker for white workers in the Midwest region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	69374.0	86.5	69374.0	86.5
Hired	10368.2	12.9	79742.2	99.4
Other	444.6	0.6	80186.8	100.0

Table 3-396. Estimates of lost-time injuries by gender for white workers in the Midwest region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	71706.1	89.4	71706.1	89.4
Female	8480.7	10.6	80186.8	100.0

Table 3-397. Estimates of lost-time injuries by age group for white workers in the Midwest region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	7085.7	8.8	7085.7	8.8
20-29	9280.8	11.6	16366.5	20.4
30-39	20843.1	26.0	37209.6	46.4
40-49	13057.2	16.3	50266.8	62.7
50-59	15728.7	19.6	65995.5	82.3
60-69	10778.2	13.4	76773.7	95.7
>=70	3413.1	4.3	80186.8	100.0

Table 3-398. Estimates of lost-time injuries by month for white workers in the Midwest region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	3823.4	4.8	3823.4	4.8
February	1545.2	1.9	5368.6	6.7
March	1815.0	2.3	7183.6	9.0
April	7139.9	8.9	14323.5	17.9
May	7051.1	8.8	21374.6	26.7
June	11873.2	14.8	33247.8	41.5
July	11087.2	13.8	44335.0	55.3
August	8250.0	10.3	52585.0	65.6
September	9920.4	12.4	62505.4	77.9
October	6387.5	8.0	68892.9	85.9
November	6437.8	8.0	75330.7	93.9
December	4856.1	6.1	80186.8	100.0

Table 3-399. Estimates of lost-time injuries by severity of the injury for white workers in the Midwest region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	77316.4	96.4	77316.4	96.4
Permanent	1992.1	2.5	79308.5	98.9
Fatality	878.3	1.1	80186.8	100.0

Table 3-400. Estimates of lost-time injuries by whether professional medical attention was sought for white workers in the Midwest region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	2870.4	3.6	2870.4	3.6
Yes	65539.2	81.7	68409.6	85.3
No	11777.2	14.7	80186.8	100.0

Table 3-401. Estimates of lost-time injuries by body part injured for white workers in the Midwest region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	4016.9	5.0	4016.9	5.0
Eye	3609.4	4.5	7626.3	9.5
Chest/trunk	5015.0	6.3	12641.3	15.8
Back	9197.5	11.5	21838.8	27.2
Arm/shoulder	6953.3	8.7	28792.1	35.9
Finger	9493.2	11.8	38285.3	47.7
Hand/wrist	9595.5	12.0	47880.8	59.7
Leg/knee/hip	14649.1	18.3	62529.9	78.0
Toe	455.8	0.6	62985.7	78.5
Foot	8237.5	10.3	71223.2	88.8
Mult. parts	7947.6	9.9	79170.8	98.7
Other	1016.0	1.3	80186.8	100.0

Table 3-402. Estimates of lost-time injuries by nature of injury for white workers in the Midwest region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	361.4	0.5	361.4	0.5
Bruise	11127.6	13.9	11489.0	14.3
Burn	314.3	0.4	11803.3	14.7
Cut	15301.0	19.1	27104.3	33.8
Crush	4534.3	5.7	31638.6	39.5
Fracture	12058.5	15.0	43697.1	54.5
Poisoning	33.3	0.0	43730.4	54.5
Puncture	4940.3	6.2	48670.7	60.7
Sprain/strain	19591.8	24.4	68262.5	85.1
Mult. injury	7320.1	9.1	75582.6	94.3
Other	4604.2	5.7	80186.8	100.0

Table 3-403. Estimates of lost-time injuries by type of event for white workers in the Midwest region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	1996.3	2.5	1996.3	2.5
Caught between	8383.9	10.5	10380.2	12.9
Caught under	4749.0	5.9	15129.2	18.9
Struck by/against	17777.5	22.2	32906.7	41.0
Struck by falling	3729.9	4.7	36636.6	45.7
Struck by flying	2280.4	2.8	38917.0	48.5
Contact sharp obj.	8218.1	10.2	47135.1	58.8
Fall, same level	11117.6	13.9	58252.7	72.6
Fall, elevation	8944.1	11.2	67196.8	83.8
Overexertion	6024.6	7.5	73221.4	91.3
Other	6965.4	8.7	80186.8	100.0

Table 3-404. Estimates of lost-time injuries by activity when worker was injured for white workers in the Midwest region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	13082.6	16.3	13082.6	16.3
Machine maint.	12020.9	15.0	25103.5	31.3
Field work	10889.1	13.6	35992.6	44.9
Crop handling	14715.2	18.4	50707.8	63.2
Livestock handling	24497.2	30.6	75205.0	93.8
Other	4981.8	6.2	80186.8	100.0

Table 3-405. Estimates of lost-time injuries by source of the injury for white workers in the Midwest region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	4931.0	6.1	4931.0	6.1
Machinery	15880.7	19.8	20811.7	26.0
Livestock	15470.2	19.3	36281.9	45.2
Hand tool	9570.5	11.9	45852.4	57.2
Power tool	2646.4	3.3	48498.8	60.5
Pesticide/chem.	337.6	0.4	48836.4	60.9
Plant/tree	1274.9	1.6	50111.3	62.5
Working surface	7367.7	9.2	57479.0	71.7
Truck/auto	3321.6	4.1	60800.6	75.8
Other vehicle	2666.2	3.3	63466.8	79.1
Liquids	493.3	0.6	63960.1	79.8
Other	16226.7	20.2	80186.8	100.0

Table 3-406. Estimates of lost-time injuries by farm type for Hispanic workers in the Midwest region of the U.S., 1993.

<u>Farm Type</u>	<u>Frequency</u>	<u>Percent</u>	<u>Cumulative Frequency</u>	<u>Cumulative Percent</u>
Cash grain	222.3	11.3	222.3	11.3
Field crops	578.3	29.3	800.6	40.6
Veg, fruit, nuts	613.6	31.1	1414.2	71.7
Beef, hogs, sheep	558.2	28.3	1972.4	100.0

Table 3-407. Estimates of lost-time injuries by status of the injured worker for Hispanic workers in the Midwest region of the U.S., 1993.

<u>Status of Injured Worker</u>	<u>Frequency</u>	<u>Percent</u>	<u>Cumulative Frequency</u>	<u>Cumulative Percent</u>
Family	616.9	31.3	616.9	31.3
Hired	1355.5	68.7	1972.4	100.0

Table 3-408. Estimates of lost-time injuries by gender for Hispanic workers in the Midwest region of the U.S., 1993.

<u>Gender</u>	<u>Frequency</u>	<u>Percent</u>	<u>Cumulative Frequency</u>	<u>Cumulative Percent</u>
Male	1972.4	100.0	1972.4	100.0

Table 3-408. Estimates of lost-time injuries by age group for Hispanic workers in the Midwest region of the U.S., 1993.

<u>Age Group (Years)</u>	<u>Frequency</u>	<u>Percent</u>	<u>Cumulative Frequency</u>	<u>Cumulative Percent</u>
10-19	279.1	14.2	279.1	14.2
20-29	120.7	6.1	399.8	20.3
30-39	845.2	42.9	1245.0	63.1
40-49	389.6	19.8	1634.6	82.9
60-69	58.7	3.0	1693.3	85.8
>=70	279.1	14.2	1972.4	100.0

Table 3-410. Estimates of lost-time injuries by month for Hispanic workers in the Midwest region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	772.0	39.1	772.0	39.1
February	281.0	14.2	1053.0	53.4
March	0.0	0.0	1053.0	53.4
April	0.0	0.0	1053.0	53.4
May	0.0	0.0	1053.0	53.4
June	0.0	0.0	1053.0	53.4
July	250.7	12.7	1303.7	66.1
August	0.0	0.0	1303.7	66.1
September	279.1	14.2	1582.8	80.2
October	0.0	0.0	1582.8	80.2
November	389.6	19.8	1972.4	100.0
December	0.0	0.0	1972.4	100.0

Table 3-411. Estimates of lost-time injuries by severity of the injury for Hispanic workers in the Midwest region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	1972.4	100.0	1972.4	100.0

Table 3-412. Estimates of lost-time injuries by whether professional medical attention was sought for Hispanic workers in the Midwest region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Yes	1693.3	85.8	1693.3	85.8
No	279.1	14.2	1972.4	100.0

Table 3-413. Estimates of lost-time injuries by body part injured for Hispanic workers in the Midwest region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	188.7	9.6	188.7	9.6
Chest/trunk	120.7	6.1	309.4	15.7
Back	279.1	14.2	588.5	29.8
Finger	222.3	11.3	810.8	41.1
Hand/wrist	772.0	39.1	1582.8	80.2
Leg/knee/hip	389.6	19.8	1972.4	100.0

Table 3-414. Estimates of lost-time injuries by nature of injury for Hispanic workers in the Midwest region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Bruise	789.4	40.0	789.4	40.0
Cut	622.9	31.6	1412.3	71.6
Crush	222.3	11.3	1634.6	82.9
Sprain/strain	337.8	17.1	1972.4	100.0

Table 3-415. Estimates of lost-time injuries by type of event for Hispanic workers in the Midwest of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught between	611.9	31.0	611.9	31.0
Caught under	279.1	14.2	891.0	45.2
Struck by/against	130.0	6.6	1021.0	51.8
Contact sharp obj.	492.9	25.0	1513.9	76.8
Fall, elevation	179.4	9.1	1693.3	85.8
Overexertion	279.1	14.2	1972.4	100.0

Table 3-416. Estimates of lost-time injuries by activity when worker was injured for Hispanic workers in the Midwest region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	222.3	11.3	222.3	11.3
Field work	510.3	25.9	732.6	37.1
Crop handling	492.9	25.0	1225.5	62.1
Livestock handling	746.9	37.9	1972.4	100.0

Table 3-417. Estimates of lost-time injuries by source of the injury for Hispanic workers in the Midwest region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	389.6	19.8	389.6	19.8
Livestock	409.1	20.7	798.7	40.5
Hand tool	492.9	25.0	1291.6	65.5
Plant/tree	120.7	6.1	1412.3	71.6
Working surface	58.7	3.0	1471.0	74.6
Other	501.4	25.4	1972.4	100.0

Table 3-418. Estimates of lost-time injuries by farm type for white workers in the West region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	1598.4	7.0	1598.4	7.0
Field crops	2028.2	8.9	3626.6	16.0
Veg, fruit, nuts	5998.4	26.4	9625.0	42.4
Nursery	2500.0	11.0	12125.0	53.4
Beef, hogs, sheep	8692.9	38.3	20817.9	91.7
Dairy	1288.0	5.7	22105.9	97.4
Other farms	596.8	2.6	22702.7	100.0

Table 3-419. Estimates of lost-time injuries by status of the injured worker for white workers in the West region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	16038.7	70.6	16038.7	70.6
Hired	6295.5	27.7	22334.2	98.4
Other	368.5	1.6	22702.7	100.0

Table 3-420. Estimates of lost-time injuries by gender for white workers in the West region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	21151.7	93.2	21151.7	93.2
Female	1551.0	6.8	22702.7	100.0

Table 3-421. Estimates of lost-time injuries by age group for white workers in the West region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	2445.9	10.8	2445.9	10.8
20-29	3998.6	17.6	6444.5	28.4
30-39	4741.5	20.9	11186.0	49.3
40-49	4651.2	20.5	15837.2	69.8
50-59	3274.5	14.4	19111.7	84.2
60-69	2136.9	9.4	21248.6	93.6
>=70	1454.1	6.4	22702.7	100.0

Table 3-422. Estimates of lost-time injuries by month for white workers in the West region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	2125.0	9.4	2125.0	9.4
February	607.0	2.7	2732.0	12.0
March	2183.3	9.6	4915.3	21.7
April	667.5	2.9	5582.8	24.6
May	3114.0	13.7	8696.8	38.3
June	2060.3	9.1	10757.1	47.4
July	2898.7	12.8	13655.8	60.2
August	2570.3	11.3	16226.1	71.5
September	1678.8	7.4	17904.9	78.9
October	961.0	4.2	18865.9	83.2
November	916.0	4.0	19781.9	87.2
December	2900.0	12.8	22681.9	100.0

(Frequency Missing = 20.8)

Table 3-423. Estimates of lost-time injuries by severity of the injury for white workers in the West region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	20736.0	91.3	20736.0	91.3
Permanent	1966.7	8.7	22702.7	100.0

Table 3-424. Estimates of lost-time injuries by whether professional medical attention was sought for white workers in the West region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	1966.7	8.7	1966.7	8.7
Yes	17535.5	77.2	19502.2	85.9
No	3200.5	14.1	22702.7	100.0

Table 3-425. Estimates of lost-time injuries by body part injured for white workers in the West region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	1091.5	4.8	1091.5	4.8
Eye	3318.6	14.6	4410.1	19.4
Chest/trunk	1348.8	5.9	5758.9	25.4
Back	3059.8	13.5	8818.7	38.8
Arm/shoulder	1350.3	5.9	10169.0	44.8
Finger	3844.7	16.9	14013.7	61.7
Hand/wrist	1389.8	6.1	15403.5	67.8
Leg/knee/hip	4254.2	18.7	19657.7	86.6
Toe	198.2	0.9	19855.9	87.5
Foot	826.6	3.6	20682.5	91.1
Mult. parts	1193.7	5.3	21876.2	96.4
Other	826.5	3.6	22702.7	100.0

Table 3-426. Estimates of lost-time injuries by nature of injury for white workers in the West region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	190.3	0.8	190.3	0.8
Bruise	2383.3	10.5	2573.6	11.3
Burn	422.0	1.9	2995.6	13.2
Cut	5562.1	24.5	8557.7	37.7
Crush	2379.1	10.5	10936.8	48.2
Fracture	3163.1	13.9	14099.9	62.1
Poisoning	356.5	1.6	14456.4	63.7
Puncture	2148.5	9.5	16604.9	73.1
Sprain/strain	3614.2	15.9	20219.1	89.1
Mult. injury	1459.6	6.4	21678.7	95.5
Other	1024.0	4.5	22702.7	100.0

Table 3-427. Estimates of lost-time injuries by type of event for white workers in the West region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	655.4	2.9	655.4	2.9
Caught between	3361.0	14.8	4016.4	17.7
Caught under	663.6	2.9	4680.0	20.6
Struck by/against	4878.8	21.5	9558.8	42.1
Struck by falling	450.9	2.0	10009.7	44.1
Struck by flying	2496.2	11.0	12505.9	55.1
Contact sharp obj.	1349.3	5.9	13855.2	61.0
Fall, same level	1989.2	8.8	15844.4	69.8
Fall, elevation	2811.8	12.4	18656.2	82.2
Overexertion	2356.0	10.4	21012.2	92.6
Other	1690.5	7.4	22702.7	100.0

Table 3-428. Estimates of lost-time injuries by activity when worker was injured for white workers in the West region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	1849.4	8.1	1849.4	8.1
Machine maint.	5577	24.6	7426.4	32.7
Field work	484.3	2.1	7910.7	34.8
Crop handling	1022.3	4.5	8933	39.3
Livestock handling	6971.7	30.7	15904.7	70.1
Other	6798	29.9	22702.7	100.0

Table 3-429. Estimates of lost-time injuries by source of the injury for white workers in the West region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Machinery	3578.5	15.8	3578.5	15.8
Livestock	5008.8	22.1	8587.3	37.8
Hand tool	3343.5	14.7	11930.8	52.6
Power tool	219.0	1.0	12149.8	53.5
Pesticide/chem.	115.8	0.5	12265.6	54.0
Plant/tree	576.1	2.5	12841.7	56.6
Working surface	1457.3	6.4	14299.0	63.0
Truck/auto	244.4	1.1	14543.4	64.1
Liquids	734.4	3.2	15277.8	67.3
Other	7424.9	32.7	22702.7	100.0

Table 3-430. Estimates of lost-time injuries by farm type for Hispanic workers in the West region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	115.8	0.3	115.8	0.3
Field crops	2090.6	6.0	2206.4	6.3
Veg, fruit, nuts	18350.9	52.3	20557.3	58.6
Nursery	6581.2	18.7	27138.5	77.3
Beef, hogs, sheep	1298.8	3.7	28437.3	81.0
Dairy	5026.8	14.3	33464.1	95.3
Other farms	1642.2	4.7	35106.3	100.0

Table 3-431. Estimates of lost-time injuries by status of the injured worker for Hispanic workers in the West region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	1074.3	3.1	1074.3	3.1
Hired	33868.0	96.5	34942.3	99.5
Other	164.0	0.5	35106.3	100.0

Table 3-432. Estimates of lost-time injuries by gender for Hispanic workers in the West region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	34186.7	97.4	34186.7	97.4
Female	919.6	2.6	35106.3	100.0

Table 3-433. Estimates of lost-time injuries by age group for Hispanic workers in the West region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	194.4	0.6	194.4	0.6
20-29	12777.4	36.4	12971.8	37.0
30-39	11033.9	31.4	24005.7	68.4
40-49	8163.9	23.3	32169.6	91.6
50-59	2326.6	6.6	34496.2	98.3
>=70	610.1	1.7	35106.3	100.0

Table 3-434. Estimates of lost-time injuries by month for Hispanic workers in the West region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	1762.5	5.0	1762.5	5.0
February	419.1	1.2	2181.6	6.2
March	1930.1	5.5	4111.7	11.7
April	1459.2	4.2	5570.9	15.9
May	1224.1	3.5	6795.0	19.4
June	5583.7	15.9	12378.7	35.3
July	5929.7	16.9	18308.4	52.2
August	7089.8	20.2	25398.2	72.3
September	2251.9	6.4	27650.1	78.8
October	2560.7	7.3	30210.8	86.1
November	4583.1	13.1	34793.9	99.1
December	312.4	0.9	35106.3	100.0

Table 3-435. Estimates of lost-time injuries by severity of the injury for Hispanic workers in the West region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	34648.1	98.7	34648.1	98.7
Permanent	458.2	1.3	35106.3	100.0

Table 3-436. Estimates of lost-time injuries by whether professional medical attention was sought for Hispanic workers in the West region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	458.2	1.3	458.2	1.3
Yes	29455.4	83.9	29913.6	85.2
No	5192.7	14.8	35106.3	100.0

Table 3-437. Estimates of lost-time injuries by body part injured for Hispanic workers in the West region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	1932.2	5.5	1932.2	5.5
Eye	3809.4	10.9	5741.6	16.4
Chest/trunk	4383.2	12.5	10124.8	28.8
Back	10584.9	30.2	20709.7	59.0
Arm/shoulder	3619.5	10.3	24329.2	69.3
Finger	3736.6	10.6	28065.8	79.9
Hand/wrist	2032.6	5.8	30098.4	85.7
Leg/knee/hip	1861.7	5.3	31960.1	91.0
Toe	194.4	0.6	32154.5	91.6
Foot	954.4	2.7	33108.9	94.3
Mult. parts	46.3	0.1	33155.2	94.4
Other	1951.1	5.6	35106.3	100.0

Table 3-438. Estimates of lost-time injuries by nature of injury for Hispanic workers in the West region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Bruise	5002.6	14.2	5002.6	14.2
Burn	1034.5	2.9	6037.1	17.2
Cut	4388.4	12.5	10425.5	29.7
Crush	809.8	2.3	11235.3	32.0
Fracture	1577.3	4.5	12812.6	36.5
Puncture	1149.0	3.3	13961.6	39.8
Sprain/strain	16939.2	48.3	30900.8	88.0
Other	4205.5	12.0	35106.3	100.0

Table 3-439. Estimates of lost-time injuries by type of event for Hispanic workers in the West region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	610.1	1.7	610.1	1.7
Caught between	1228.7	3.5	1838.8	5.2
Caught under	1080.8	3.1	2919.6	8.3
Struck by/against	1234.5	3.5	4154.1	11.8
Struck by falling	1457.0	4.2	5611.1	16.0
Struck by flying	671.8	1.9	6282.9	17.9
Contact sharp obj.	5144.4	14.7	11427.3	32.6
Fall, same level	2931.9	8.4	14359.2	40.9
Fall, elevation	1478.3	4.2	15837.5	45.1
Overexertion	11980.2	34.1	27817.7	79.2
Other	7288.6	20.8	35106.3	100.0

Table 3-440. Estimates of lost-time injuries by activity when worker was injured for Hispanic workers in the West region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	5519.0	15.7	5519.0	15.7
Machine maint.	1649.5	4.7	7168.5	20.4
Field work	13639.5	38.9	20808.0	59.3
Crop handling	4253.9	12.1	25061.9	71.4
Livestock handling	1295.0	3.7	26356.9	75.1
Other	8749.4	24.9	35106.3	100.0

Table 3-441. Estimates of lost-time injuries by source of the injury for Hispanic workers in the West region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	583.9	1.7	583.9	1.7
Machinery	4084.5	11.6	4668.4	13.3
Livestock	2177.6	6.2	6846.0	19.5
Hand tool	5326.9	15.2	12172.9	34.7
Power tool	2267.1	6.5	14440.0	41.1
Plant/tree	3287.9	9.4	17727.9	50.5
Working surface	2583.3	7.4	20311.2	57.9
Truck/auto	772.2	2.2	21083.4	60.1
Other vehicle	1034.3	2.9	22117.7	63.0
Other	12988.6	37.0	35106.3	100.0

Table 3-442. Estimates of lost-time injuries by farm type for black and other race workers in the West region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Field crops	734.9	38.7	734.9	38.7
Veg, fruit, nuts	114.5	6.0	849.4	44.8
Beef, hogs, sheep	151.9	8.0	1001.3	52.8
Poultry	895.8	47.2	1897.1	100.0

Table 3-443. Estimates of lost-time injuries by status of the injured worker for black and other race workers in the West region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	752.8	39.7	752.8	39.7
Hired	1144.3	60.3	1897.1	100.0

Table 3-444. Estimates of lost-time injuries by gender for black and other race workers in the Westt region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	1334.6	70.3	1334.6	70.3
Female	562.5	29.7	1897.1	100.0

Table 3-445. Estimates of lost-time injuries by age group for black and other race workers in the West region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	190.3	10.0	190.3	10.0
20-29	315.9	16.7	506.2	26.7
30-39	190.3	10.0	696.5	36.7
40-49	638.1	33.6	1334.6	70.3
60-69	562.5	29.7	1897.1	100.0

Table 3-446. Estimates of lost-time injuries by month for black and other race workers in the West region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	151.9	8.0	151.9	8.0
February	0.0	0.0	151.9	8.0
March	0.0	0.0	151.9	8.0
April	0.0	0.0	151.9	8.0
May	0.0	0.0	151.9	8.0
June	0.0	0.0	151.9	8.0
July	304.8	16.1	456.7	24.1
August	0.0	0.0	456.7	24.1
September	544.6	28.7	1001.3	52.8
October	0.0	0.0	1001.3	52.8
November	0.0	0.0	1001.3	52.8
December	895.8	47.2	1897.1	100.0

Table 3-447. Estimates of lost-time injuries by severity of the injury for black and other race workers in the West region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	1706.8	90.0	1706.8	90.0
Permanent	190.3	10.0	1897.1	100.0

Table 3-448. Estimates of lost-time injuries by whether professional medical attention was sought for black and other race workers in the West region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	190.3	10.0	190.3	10.0
Yes	1390.9	73.3	1581.2	83.3
No	315.9	16.7	1897.1	100.0

Table 3-449. Estimates of lost-time injuries by body part injured for black and other race workers in the West region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Eye	114.5	6.0	114.5	6.0
Chest/trunk	151.9	8.0	266.4	14.0
Back	354.3	18.7	620.7	32.7
Finger	333.3	17.6	954.0	50.3
Hand/wrist	943.1	49.7	1897.1	100.0

Table 3-450. Estimates of lost-time injuries by nature of injury for black and other race workers in the West region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Crush	523.6	27.6	523.6	27.6
Fracture	752.8	39.7	1276.4	67.3
Sprain/strain	506.2	26.7	1782.6	94.0
Other	114.5	6.0	1897.1	100.0

Table 3-451. Estimates of lost-time injuries by type of event for black and other race workers in the West region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	610.1	1.7	610.1	1.7
Caught between	1228.7	3.5	1838.8	5.2
Caught under	1080.8	3.1	2919.6	8.3
Struck by/against	1234.5	3.5	4154.1	11.8
Struck by falling	1457.0	4.2	5611.1	16.0
Struck by flying	671.8	1.9	6282.9	17.9
Contact sharp obj.	5144.4	14.7	11427.3	32.6
Fall, same level	2931.9	8.4	14359.2	40.9
Fall, elevation	1478.3	4.2	15837.5	45.1
Overexertion	11980.2	34.1	27817.7	79.2
Other	7288.6	20.8	35106.3	100.0

Table 3-452. Estimates of lost-time injuries by activity when worker was injured for black and other race workers in the West region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	190.3	10.0	190.3	10.0
Machine maint.	687.6	36.2	877.9	46.3
Field work	562.5	29.7	1440.4	75.9
Crop handling	190.3	10.0	1630.7	86.0
Livestock handling	151.9	8.0	1782.6	94.0
Other	114.5	6.0	1897.1	100.0

Table 3-453. Estimates of lost-time injuries by source of the injury for black and other race workers in the West region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Machinery	523.6	27.6	523.6	27.6
Livestock	151.9	8.0	675.5	35.6
Hand tool	544.6	28.7	1220.1	64.3
Pesticide/chem.	114.5	6.0	1334.6	70.3
Plant/tree	562.5	29.7	1897.1	100.0