

**Analysis of Bottomland Hardwood Areas and Assessment of Wildlife
Habitat Quality at the Potential Future Waters Bluff Reservoir Site**

Final Report to

Texas Water Development Board for the fulfillment of
interagency agreement No. 97-483-226

Texas Parks & Wildlife Department
Austin, Texas

April, 1998

Part 1

Analysis of Bottomland Areas and Classification of Land Cover Types

Principal Investigators:

Changxiang Liu, Ph.D., Christine Coats, Jason Singhurst, and
A. Kim Ludeke, Ph.D.

GIS Lab, Texas Parks & Wildlife Department

April, 1998

Introduction

Bottomland hardwood areas are considered among the most important habitats for wildlife. In these areas, forests, swamps, ponds, and meandering channels provide not only various habitats needed by a variety of fish and wildlife species and migrating birds, but also contain resources necessary for their survival. These areas are also rich in plant and animal species diversity. Bottomland hardwood forests play an important role in controlling soil erosion, maintaining water quality, recharging groundwater, and preventing flood damage. On the other hand, the bottomland hardwood areas are primary locations for economic development. Hardwood trees, fertile soils, water resources, and unique landscape attract many kinds of development activities, such as agriculture, forestry, tourism, industry, and water development. As a consequence, bottomland hardwood forests have been disappearing rapidly since the turn of this century in Texas as well as throughout the nation.

As human population increases, demand for water supply in urban areas, by agriculture, and industries rises dramatically. In Texas, periodic droughts have made the water supply very critical for the economic development and sustainability of the state, hence the driving force for more water development projects. Unavoidably, there are always losses associated with every water resources development project. One of those losses is the loss of bottomland hardwood areas and wildlife habitats. Federal and state laws mandate that an assessment of negative impacts and mitigation be made before any projects can take place.

This part of the project is to study the direct impact of the proposed Waters Bluff reservoir on bottomland hardwood areas at the proposed reservoir site and provides the Texas Water Development Board (TWDB) with essential data on the types and acreage of bottomland forests subject to inundation.

The land cover types identified in this study are based solely on remotely sensed data (TM imagery and aerial photography) and field survey at the study site. No soil or hydrological data were collected and used in the determination of the land cover types as would be required in a federal delineation of forested wetlands in the bottomland hardwood areas. Some portions of the bottomland hardwood forest, as defined in this study, and other land cover types identified here may occupy the same areas as federally delineated bottomland hardwood forested wetlands. However, the land cover units presented in this report do not substitute for a federal jurisdictional wetland delineation to determine wetland areas for which mitigation may be required.

Study Area

The proposed Waters Bluff reservoir site is located between Smith, Upshur, and Wood counties along the Sabine River in northeastern Texas (Map 1). Most of the area falls

in the pineywood region of eastern Texas, even though post oak savanna appears occasionally in the western part of the study area.

Shortleaf pine (*Pinus echinata* Mill.), loblolly pine (*Pinus taeda* L.), and other oak and hardwood species dominate forest communities in the uplands and on hill slopes. The bottomland areas and floodplains are mainly made up of bottomland hardwood species typical to east and southeast Texas. Other common tree species in the study area include eastern red cedar (*Juniperus virginiana* L.), post oak (*Q. stellata* Wangenh.), black hickory (*Carya texana* Buckl.), sweetgum (*Liquidambar styraciflua* L.), water oak (*Q. nigra* L.), southern red oak (*Q. falcata* Michx.), blackgum (*Nyssa sylvatica* Marsh.), blackjack oak (*Q. marilandica* Muenchh.), willow oak (*Q. phellos* L.), basket oak (*Q. michauxii* Nutt.), overcup oak (*Q. lyrata* Walt.), river birch (*Betula nigra* L.), red maple (*Acer rubrum* L.), green ash (*Fraxinus pennsylvanica* Marsh.), and American elm (*Ulmus americana* L.). Almost all the vegetation types in the area have been subjected to various human disturbances in the past (i.e. logging). Most of the area was logged or selectively logged at least once.

METHODS

We used one LandSat Thematic Mapper (TM) satellite imagery scene (path 25 and row 37), which was obtained from the Corps of Engineers, as the primary source for the land cover classification. The TM scene was originally acquired by the Earth Resources Observation Satellite Company (EROS) in September 1994. It contains 6-band data (the thermal band 6 was removed) in Universal Transverse Mercator (UTM) projection (zone 15 and datum NAD27) with a spatial resolution of 25 meters. We also purchased recent National Aerial Photograph Program (NAPP) color infrared (CIR) aerial photographs of the study area for photointerpretation and groundtruthing. Reservoir normal elevation lines (300 ft. above sea level) were drawn and bottomland hardwood areas were delineated on USGS 7.5' quadrangle paper maps after a field reconnaissance. These maps were scanned and lines were vectorized to create vector coverages. The road network is based on digital county road maps by Texas Department of Transportation (TXDOT). Texas Water Development Board provided dam locations, reservoir pool elevation and layout drawings. Data processing, classification, and analysis were performed by using ERDAS Imagine (ERDAS Inc. 1997) and ARC/INFO (ESRI 1996).

The 6-band TM image was first subjected to a principal component analysis (PCA) then a unsupervised classification. Three bands or principal component axes were extracted from the 6-band data, but retain most of the variance of the original data. The PCA processed image and CIR aerial photographs were used as the primary source for the supervised classification. In the unsupervised classification, 150 classes were obtained from the 6-band TM image.

Training samples were collected from one field reconnaissance trip to the study area in August 1997. About 100 sample points were taken in the field with additional land cover sample areas marked on aerial photographs for photointerpretation and supervised

classification. We paid special attention to the ecotones between upland and bottomland types and delineation was mostly based on the field experience and changes in contour lines. We did recognize more types than we actually can identify on the TM image. Some cover types do not yield distinguishable signatures on the TM imagery. Therefore, some of the field recognized land cover types had to be combined into other types to form a composite type, such as grassland/crops and willow-birch-water elm types.

After the field trip, the 150 classes from the unsupervised procedure were subjected to a supervised classification by using training samples, the PCA processed image, and topography (contours), and by interpreting CIR aerial photographs. A total of nine types were recognized in the process.

The area of each land cover type within the inundation level of the reservoir was calculated by superimposing and masking the classified image upon a raster layer created from 300 ft. contour line (normal conservation pool level) drawn on the USGS maps.

The groundtruthing was conducted in September 1997. We chose to sample land parcels of at least 4x4 pixels, i.e. 1 hectare to assure the point would not fall out of the parcel. About 250 valid GPS points were collected during the trip. Since most of the area was privately owned and inaccessible by field crew, the collection of groundtruth points was largely restricted to the road side. Therefore, it cannot be regarded as a random sampling. Caution must be exercised in interpreting the results of the accuracy matrix. However, the procedure is necessary for in-situ verification of the classified land cover types. Another thing we have to point out is that the TM image was acquired in 1994 and does not reflect the changes thereafter. Recent logging in several areas within the reservoir area has been noticed (Frye, personal communication). Therefore, the loss of each land cover type in the bottomland areas reported here may be somewhat overestimated.

Results

1. Description of Major Land Cover Types

Nine land cover types have been recognized in the study area. Water and bare ground/soil are considered not vegetative types, although they may have sparse or partial vegetation cover. These nine types are grassland/crops, pure pine/pine plantation, oak-hickory forest, pine-oak forest, hardwood-pine forest, bottomland hardwood forest, willow-birch-water elm forest, water, and bare ground/soil.

We describe each of these vegetation types briefly:

Pure Pine/Pine Plantation

This vegetation type is dominated primarily by loblolly pine and shortleaf pine singly or in combination, at least in the overstory. In some cases, it may be planted slash pine stands or eastern red cedar in limited areas. Stand ages and density vary greatly from place to place. This vegetation type can be found in a variety of habitats except in the wettest areas. Species abundance in the understory depends on the intensity of management and the age of the stands. Southern red oak, sweetgum, water oak, dogwood, and other hardwood species can coexist. This vegetation type is a direct result of silvicultural practices, whereas cedar groves usually develop on abandoned agricultural fields. The latter can only be found in limited areas.

Oak-Hickory Forest

Species commonly found in this type are post oak, black hickory, blackjack oak, and winged elm (*Ulmus alata* Michx.). Other associate species include southern red oak, sweetgum, dogwood (*Cornus florida* L.), bluejack oak (*Q. incana* Bartr.), and eastern red cedar (*Juniperus virginiana* L.). Sassafras, farkleberry (*Vaccinium arboreum* Mars. Var. *arboreum*), blackhaw (*Viburnum rufidulum* Raf.), yaupon (*Ilex vomitoria* Ait.), American beautyberry (*Carlicarpa americana* L.) and aromatic sumac (*Rhus aromatica* Ait.) are the common shrub species. Common grasses include splitbeard bluestem (*Andropogon ternarius* Michx), sessile flowered chasmanthium (*Chosmanthium sessiliflorum* (Poir.) Yates), little bluestem (*Schizachyrium scoparium*), and Indiangrass (*Sorghastrum nutans* (L.) Nash or *S. elliotii* (Mohr.) Nash). Soils are excessively-drained, poor in nutrients, and sandy on the ridges and hill tops. The herbaceous cover is sparse owing to dry conditions. This type can also be the result of removing of pines by selective logging.

Pine-Oak Forest

This is a upland type dominated by pines (shortleaf and loblolly) with oaks (post oak, blackjack oak, southern red oak) and hickory as the codominants. Site conditions and associated shrub and herbaceous species are similar to that of the oak-hickory forest type.

Hardwood-Pine Forest

This type is a mixture of hardwood species and a few pines (shortleaf and loblolly pines) growing in the floodplain and wet areas. The habitat is similar to that of the bottomland hardwood forest type. The overstory of this type is predominantly made up of hardwood trees, with scattered pines. This type commonly occurs on better drained bottomland soils within the floodplain including old levees, flats, and higher terraces. These areas are correlated with Zone V, of the classification system developed by the National Wetlands Technical Council (Larson et al. 1981).

Common hardwood species are water oak, sweetgum, blackgum, willow oak, white oak, sugar maple (*Acer saccharum* Marsh), hawthornes (*Crataegus* spp.), buckthorn (*Rhamnus caroliniana* Walt.). Common grasses are splitbeard bluestem, sessile chasmanthium, and little bluestem. Presence of pines is minimal, and in some areas it is a result of plantation. This type was included as a bottomland hardwood and associated wetland community type 3, Forested Wetland (Palustrine) by the US Fish and Wildlife Service (1985) in an inventory of bottomland hardwood forest types in Texas. Therefore, we do consider this type a bottomland forest.

Bottomland Hardwood Forest

Bottomland hardwood forest occurs on most sites within the floodplains and flats and in areas along river channels. Species commonly found in this type include water oak, willow oak, blackgum, American elm, overcup oak, green ash, deciduous holly (*Ilex decidua* Walt.), American holly (*I. opaca* Ait.), hawthornes, swamp privet (*Foresteria acuminata* (Michx.) Poir.), American hornbeam (*Carpinus caroliniana* Walt.), and occasionally palmetto (*Sabal minor* (Jacq.) Pers.). Many vines (*Smilax* spp., *Vitis* spp., *Parthenocissus quinquefolia* L. Planch, *Rubus* spp., *Berchemia scandens* (Hill) K. Koch, etc.) can be found in this type. Common grasses are longleaf chasmanthium (*Chasmanthium latifolium* (Michx.) Yates), wildrye (*Elymus virginicus* L.), cutgrass (*Leersia lenticularis* Michx. or *L. oryzoides* (L.) Sw.), switchgrass (*Panicum virgatum* L. and *P. anceps* Michx.), etc. Periodic inundation prevents the establishment of upland species and maintains the functioning of this vegetation type.

Willow-Birch-Water Elm Forest

The willow-birch-water elm type occurs along the river and creek channels, depressed areas, and at the confluence of lakes and creeks. This type exists in narrow strips rarely exceeding 100 meters in width. Other species coexisting in this type include river birch, cottonwood (*Populus deltaoides* Marsh.), red maple, water oak, sweetgum, pecan (*Carya illinoensis* (Wangehn.) K. Koch.), blackgum, water elm (*Planera aquatica* J. G. Gmel.) , willow (*Salix nigra* Marsh. var. *nigra*), hawthornes, and American elm. Shrub, grasses and vines are similar to the bottomland hardwood type. This type may also include small swamp areas which are present in the study area but not significantly.

Grassland/Crops

These areas are dominated by grasses, graminoids (e.g. sedges and rushes), and other herbaceous species. Some of these areas were once forested, but were cut and converted to agricultural uses. This type includes pastures and natural grasslands. Species with high abundance include little bluestem, bluestems (*Andropogon gerardii*, *A. ternarius*, *A. virginicus*, *A. glomeratus*), threeawns (*Aristida* spp.), Johnson grass (*Sorghum halapense*), eastern gammagrass (*Tripsacum dactyloides*), Bahia grass

(*Paspalum dilatatum*), Bermuda grass (*Cynodon dactylon*), rushes (*Juncus* spp.) and sedges (*Carex* spp.).

Some of the grassland areas are well developed in the absence of grazing and other disturbances. This type also includes crop land which is irrigated and managed for production of grains, cotton, and other products.

Bare Ground/Soil

These are the areas with no or very little vegetation coverage. Industrial parks, pavement, mineral mines, and plowed/fallow field are some examples. However, plowed/fallow agricultural fields classified as bare soils may be temporary.

Water

This includes water bodies impounded by small dams and in oxbow lakes, and open water in river channels with no or little vegetation cover overhead. Submersed aquatic vegetation may be present in some areas.

In Part two of this report, the mixed upland forest type is equivalent to the combination of pine-oak forest and oak-hickory forest types, and the mixed bottomland hardwood forest type is the combination of three types, hardwood-pine forest, bottomland hardwood forest, and willow-birch-water elm forest (Frye and Curtis 1990).

We also call these three types as bottomland forest, or forested wetlands, as they mostly occur in the floodplain and wet areas (Larson et al. 1981. Fish and Wildlife Service 1985). In the study area, there are small parcels of a mesic type (white oak mixed with other hardwood species) in moist slopes (Marks & Harcombe 1981), but not large enough to be classified as a separate type. This type is largely lumped with the oak-hickory type.

2. Analysis of Bottomland Areas to be Lost to the Proposed Waters Bluff Reservoir

The overall accuracy (total number of correctly classified groundtruth points divided by the total groundtruth points) for the classified land cover types is 83.1% based on about 250 collected groundtruth points (Table 1). The in-situ types in Table 1 are the land cover types identified in the field and the types in the first column is the corresponding types on the classified TM image. The percentage along the diagonal of Table 1 are the percentage of groundtruth points correctly classified. Land cover types classified with high confidence include grassland/crops (96.7%), oak-hickory (100%), bottomland hardwood (91.2%), and oak-pine (81.6%). We did not collect enough samples for water, hardwood-pine, and bare ground/soil due to their limited and narrow distribution.

Table 1 Accuracy Matrix Derived from Groundtruth Samples

<i>In-situ</i> types* -->	Water	Birch-willow-water elm	Bottomland hardwood	Hardwood-pine	Oak-pine	Pine plantation	Oak-hickory	Grassland	Bare ground/soil	Total Points
Water	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1
Birch-willow-water elm	0.0	21.4	50.0	0.0	0.0	0.0	21.4	7.1	0.0	14
Bottomland hardwood	0.0	0.0	91.2	0.0	2.9	0.0	5.9	0.0	0.0	34
Hardwood-pine	0.0	0.0	0.0	55.6	22.2	22.2	0.0	0.0	0.0	9
Oak-pine	0.0	0.0	0.0	0.0	81.6	0.0	15.8	2.6	0.0	38
Pine plantation	0.0	0.0	0.0	0.0	25.0	58.3	12.5	4.2	0.0	24
Oak-hickory	0.0	0.0	0.0	0.0	0.0	0.0	100.0	0.0	0.0	28
Grassland	0.0	0.0	0.0	0.0	0.0	0.0	0.0	96.7	3.3	92
Bare ground/soil	0.0	0.0	12.5	0.0	0.0	0.0	12.5	25.0	50.0	8
Total Groundtruth Points	1	3	39	5	40	16	43	94	7	248

Note: Overall Accuracy: 83.1%

* In-situ types are the actual land cover types, and types in the first column are read from the classified TM image.

The area to be flooded is totaled at 31,018 acres (under average pool elevation of 300 ft.), of which, 68.2% (21,155 acres) is bottomland hardwood forest (Table 2). If willow-birch-water elm (704 acres) and hardwood-pine (3,072 acres) types

Table 2 Acreage Summary of Recognized Land Cover Types in Flooded Area

No	Land Cover	Hectares	Acres	% of Total
1	Water	72.4	179.0	0.6
2	Pure Pine/Pine Plantation	226.4	559.4	1.8
3	Pine-Oak Forest	97.3	240.5	0.8
4	Hardwood-Pine Forest	1,243.2	3,072.0	9.9
5	Oak-Hickory Forest	344.1	850.2	2.7
6	Bottomland Hardwood Forest	8,561.1	21,155.0	68.2
7	Willow-Birch-Water Elm Forest	284.9	704.1	2.3
8	Grassland/Crops	1,714.8	4,237.4	13.7
9	Bare Ground/Soil	8.4	20.8	0.1
Total		12,552.70	31,018.40	100.0

are included, the total loss of bottomland forests will be 81.4%. In addition, 4,237 acres (13.7%) of grassland or pasture will be inundated as well. There are two large tracts of bottomland hardwood forest within the flooded area. One is between the US Highway 69 bridge and the Texas Highway 155 bridge and the other is between the Texas Highway 155 bridge and the US Highway 259 bridge. Most of the former is a continuous tract under the stewardship of Texas Parks and Wildlife Department (Old Sabine Bottom Wildlife Management Area) and the US Fish and Wildlife Service (Little

Sandy Creek National Wildlife Refuge). The latter is not protected and shows fragmentation by pasture, plantation, and logging in some areas.

The quality assessment of and compensation calculation for these forests are addressed in Part 2 of this report.

Acknowledgements

We are grateful to Duane German for providing computer support and technical advice and Craig Scofield for GPS support. We also would like to thank Roy Frye of Wildlife Division for reviewing the manuscript and Joyce Gilliam for preparing the text layout.

References

- Cowardin, L. M., Carter, V, Colet, F. C., and Laroe, E. T. 1979. Classification of wetlands and deepwater habitats of the United States. Fish and Wildlife Service, US Department of Interior, Washington D. C.
- Dahl, T. E. and C. E. Johnson. 1991. Wetlands status and trends in the conterminous United States mid-1970's to mid-1980's: first update of the national wetlands status report. U.S. Fish and Wildlife Service, Department of Interior, Washington D. C.
- ERDAS Inc. 1997. ERDAS Field Guide, 4th edition. Atlanta, Georgia.
- Eyre, F. H. (editor). 1980. Forest cover types of the United States and Canada. Society of American Foresters, Washington D. C.
- Fish and Wildlife Service. 1985. Texas bottomland hardwood preservation program -- final concept plan. U.S. Fish and Wildlife Service, Albuquerque, N.M.
- Frye, R. G. and D. A. Curtis. 1990. Texas water and wildlife: an assessment of direct impacts to wildlife habitat from future water development projects. Austin, Texas.
- Hefner, J. M., and J. D. Brown. 1985. Wetland trends in the southeastern United States. *Wetlands* 4:1-12.
- Larson, J. S., M. S. Bedinger, C. F. Bryan, S. Brown, R. T. Huffman, E. L. Miller, D. G. Rhodes, and B. A. Touchet. 1981. Transition from wetlands to uplands in southeastern bottomland hardwood forests. pp. 225-273 in J. R. Clark and J. Benforado, eds. *Wetlands of bottomland hardwood forests. Proceedings of a workshop on bottomland hardwood forest wetlands of the southeastern United States.* Lake Lanier, Georgia.
- Marks, P. L. and P. A. Harcombe. 1981. Forest vegetation of the Big Thicket, southeast Texas. *Ecological Monographs* 51:287-305.
- McMahan, C. A., R. G. Frye, and K. L. Brown. 1984. The vegetation types of Texas including cropland. Texas Parks and Wildlife Department, Austin, Texas.

TABLES AND MAPS

Table 1 Accuracy matrix for the classified land cover types derived from groundtruth samples

Table 2 Acreage summary of recognized land cover types in the flooded areas


Map 1 Generalized study area at the proposed Waters Bluff reservoir site

Map 2 Land cover types at proposed Waters Bluff reservoir site and vicinity

ELECTRONIC DATA IN THE CD-ROM *(file name and software in parentheses)

1. Land cover types at the study site classified from TM image (wblc.img and wblc.rrd, Imagine)
2. Dam locations and layouts (damlms, ARC/INFO)
3. Normal conservation pool level or inundation line (lms300ft, ARC/INFO)
4. Road network at the study site (roads, ARC/INFO)

* Only limited copies of this final report contain CD-ROM. The data are available upon request.



Map 1 Generalized Study Area at Proposed Waters Bluff Reservoir Site



Legend	
Land Cover Type	
	Water
	Forest (Pine/Oak)
	Pine/Oak
	Forest/Grassland
	Urban/Industrial
	Barren/Unimproved
	Wetlands/Waterways
	Grasslands/Cropland
	Barren/Unimproved

	North Arrow
	Scale
	Miles



Scale
0
Miles

Data Sources:
MAPS Code Volume Aerial Photographs
Land Use Inventory, 1987, 1994
USGS 7.5 Topographic Maps



Map 2 Land Cover Types at
the Proposed Watershield
Reservoir Site, Texas

Created by
GIS Lab
Texas Parks & Wildlife Dept.
Austin, Texas
For Texas Water Development Board
August 28, 1987

Part 2

Wildlife Habitat Quality Assessment for the Proposed Waters Bluff Reservoir Site

Principal Investigators:

Roy G. Frye, Kathy Boydston, Dr. Ray C. Telfair II

Wildlife Division
Texas Parks and Wildlife Department

April 1998

Introduction

The purpose of this study was to complete analyses to determine the quality of wildlife habitat for the vegetation cover types inventoried and mapped for the proposed Waters Bluff Reservoir. A separate vegetation inventory was completed as described in Part 1 of the report. Habitat quality information will assist the Texas Water Development Board with its water planning efforts. Work focused on Bottomland Hardwood Forests as these forests account for most of the vegetational coverage within the reservoir site, have been heavily impacted in Texas, and exhibit economic and ecological value. Activities were guided by provisions of Goal 2, Appendix A (Scope of Work), Texas Water Development Board (TWDB) Interagency Contract No. 97-483-226. This analysis was conducted by staff of the Wildlife Habitat Assessment Program, Wildlife Division, Texas Parks and Wildlife Department. The wildlife habitat assessment methodology detailed below uses vegetation cover types delineated from Part 1 as a basis for calculation of mitigation requirements. The bottomland hardwood forest and other land cover types in this study may occupy some of the same areas as federally delineated wetlands. Nevertheless, the location and areal information about bottomland hardwood forests and other cover types in this study cannot be used in place of a federal jurisdictional wetland delineation since these cover type definitions do not include consideration of hydric soils and hydroperiod information. Therefore, calculations for determining compensation requirements give only a preliminary estimate of the scale of mitigation that would be required if the reservoir were constructed.

Study Area

The Waters Bluff Reservoir site lies principally within the floodplain of the Sabine River in Wood, Smith, and Upshur Counties. The communities of Hawkins and Big Sandy are just to the north of the site, while the city of Gladewater is within proximity of the northeast lower portion of the site. The site is approximately 24 miles long with a width exceeding 2 miles in several locations. The extreme western portion of the site is located within the Post Oak Savannah ecological region, while the middle and eastern portions lie within the Pineywoods region as defined by Gould et. al.(1960). Climate is subtropical, humid with warm summers and mild winters with occasional periods of sub-freezing temperatures. The average annual high temperature ranges between 75 and 76 degrees F, while the average annual low ranges between 51 and 53 degrees F. The average annual precipitation ranges from 40 to 46 inches. The average annual gross lake surface evaporation rate for this site generally ranges between 51 and 53 inches (Texas Department of Water Resources 1983).

Major vegetation cover types typical of this region have been previously mapped (McMahan et. al. 1984). These include Pine-Hardwood Forest, Willow Oak-Water Oak-Blackgum Forest, and native and/or introduced grasses. Within the broader category of Pine-Hardwood Forest, species such as loblolly pine, post oak, black hickory, eastern red cedar, hackberry, yaupon, and greenbriar commonly occur. This type is equivalent to the Mixed Upland Forest identified in Part 1 of this report. Floodplains and creek drainages are characterized by the Willow Oak-Water Oak-Blackgum association and may include other species such as overcup oak, elm, hackberry, sweetgum, willow, hawthorn, greenbriar, and water elm. This reservoir site contributes approximately 10% of the total acreage of bottomland hardwood forests occurring within the Sabine River floodplain, while the floodplain forests comprise about 22% of the remaining acreage within the major river bottoms in East Texas (Frye 1986). Cover types occurring but comprising disproportionately small acreage include pine plantations and cropland. Native and introduced grasses are generally associated with pastures, hayfields, and areas adjacent to developed urban or rural areas.

Methods

The overall quality of the occurring habitat for wildlife resources was evaluated for the Waters Bluff Reservoir Site using the wildlife habitat appraisal procedure (WHAP) (Appendix 1). The technique measures key components which contribute to the ecological condition of the classified cover types within each reservoir site and resulting overall suitability for wildlife. Habitat quality values obtained from site evaluation criteria are combined with acreage figures for each cover type to provide available Habitat Units (HU).

The method is based on the following assumptions:

1. that vegetative structure including species composition and physiognomy is itself sufficient to define the habitat suitability for wildlife;
2. that a positive relationship exists between vegetation diversity and wildlife species diversity.
3. that vegetation composition and primary productivity directly influence population densities of wildlife species.

Habitat quality scores for each cover type represent baseline conditions. The total HU's lost are numerical values that quantify initial direct impacts of reservoir construction, and to assist comparison with other projects, assume complete loss of existing vegetation cover below the proposed normal pool elevations. These numbers do not reflect annualized losses calculated over the life of the project nor account for any potential habitat gains that could be created as a result of reservoir construction.

Consequently, the compensation estimates may not be the same as estimates calculated in future site specific evaluations. Other factors which can influence these differences include changes in project assumptions, variations in project design, land use changes, and priorities for certain habitat types. The compensation estimates calculated for this report are intended to provide *preliminary* data in a format to allow comparison of reservoir site alternatives. The estimates only address direct impacts. Long term indirect impacts such as increased land use change around the proximity of the reservoirs, or any changes to vegetation composition and quality below the dams from altered instream flow are not included in this assessment.

Compensation requirements for each of the impacted cover types were calculated according to three hypothetical values representing proportional amounts (25%, 50%, and 100%) of the total potential gain in habitat quality of a compensation area which could be obtained with management. Raising the potential gain in habitat quality of a compensation area by 25% assumes relatively minimal management; an increase of 50% assumes moderate management; while achieving 100% of the potential gain assumes intensive management.

Minimal management could include marking wildlife management area boundaries, providing protection by periodic surveillance, incorporating grazing control and allowing the habitat quality to increase through natural succession. Annual estimated costs per acre for this level of management would be lower than moderate to higher management. Moderate management might include the above measures with the addition of some selected herbaceous seedings and limited vegetation manipulation through controlled burning, disking, thinning, or other means. Costs for this level would be higher than minimal management, but lower than intensive management. Intensive management would include the above measures with the addition of significant efforts to reestablish communities of grasses, forbs, woody shrubs or trees through supplemental plantings and vegetation maintenance; establishing indices of relative abundance of wildlife species and conducting research associated with wildlife needs. Annual costs for this level would involve highest funding expenditures. All three levels of management would likely include wildlife-oriented public recreational use.

All cover types evaluated for habitat quality were also classified into resource categories to denote mitigation planning goals. Such goals will be pursued by the U.S. Fish and Wildlife Service within the Federal permitting process and TPWD during the review of state water use permit applications and formulation of recommendations to the Texas Natural Resource Conservation Commission (TNRCC). A description of each resource category, designation criteria, and mitigation planning goals are provided in Table 1.

Table 1. Resource Categories and Mitigation Planning Goals.

Resource Category	Designation Criteria	Resource Planning Goal
1	High value for evaluation species or habitat, unique or irreplaceable	No loss of habitat value
2	High value for evaluation species or habitats and scarce or becoming scarce.	No net loss of in-kind habitat value.
3	High to medium value for evaluation species or habitats and commonly occurs.	No net loss of habitat value while minimizing loss of in-kind habitat value.
4	Medium to low value for evaluation species or habitats.	Minimize loss of habitat value.

Field evaluation forms used to rate the existing cover types within the two reservoir sites are provided in Appendix 2.

Field data collection required the completion of vegetation cover maps from Part 1 to allow precise location of data collection sites. Such data was not available until the first week of November, 1997. Field sampling was curtailed by extensive rainfall during the week of November 10 which hampered data collection efforts. However, a total of 20 individual sites were evaluated during the period November 10-11, 1997. The location of each site in relation to the approximate normal reservoir pool level is provided as Figure 1. Site assessments were performed by Kathy Boydston, Dr. Ray C. Telfair, II, Larry D. LeBeau, Greg Creacy, and Roy G. Frye, Wildlife Division, Texas Parks and Wildlife Department. Evaluations were conducted throughout the reservoir site as indicated by Figure 1. This included private land within the reservoir site, the Old Sabine Bottom Wildlife Management Area, and the Little Sandy National Wildlife Refuge (NWR). Existing knowledge of the general area within the reservoir site indicated that a wide variety of habitat quality existed, varying according to specific site and existing land use. The data collection efforts were structured to achieve representation throughout the reservoir site and not focus on any particular area. Results of the vegetation cover inventory in Part 1 indicated delineation of a total of 9 cover types. For the purposes of collecting field data for the Wildlife Habitat Appraisal these nine cover types were grouped as illustrated in Table 2. Note that three thematic classes were lumped into a larger mixed bottomland hardwood forest category. This included a class designated as Hardwood - Pine Forest. As discussed in Part 1, the class typifies a community characterized by mixed hardwood tree species occurring with loblolly pine located on higher, more well drained floodplain sites and is considered a bottomland hardwood type by many authorities (Eyer 1980), (FWS 1985), and (Larson, et. al 1981).


Table 2. Grouping of Classified Cover Types for Field Data Collection.

No. of Sites Evaluated	Combined Class Name (Original Mapped Name)	Acres
13	Mixed Bottomland Hardwood Forest	24,931
	(Hardwood - Pine Forest)	(3,072)
	(Bottomland Hardwood Forest)	(21,155)
3	(Willow-Birch-Water Elm Forest)	(704)
	Mixed Upland Forest	1,091
	(Pine - Oak Forest)	(241)
3	(Oak-Hickory Forest)	(850)
	Grassland/Crops	4,237
1	Pine Plantation	559
0	Other (Water, Bare Ground, soil)	200

The approximate location of sites was determined in advance of the field work to save time. This allowed for the contacting of landowners to obtain permission to access property and to assure sites adequately represented delineated cover types. Sites were also selected based on road accessibility. In all cases, investigators selected specific sites a sufficient distance from the access road to be removed from the influence of rights-of-way, easements, and fence lines. Site locations were recorded using global positioning instrumentation which was downloaded into a Department PC and subsequently plotted (Figure 1).

Proposed actions normally require mitigation measures that 1) minimize the impact by changing the project location, design, or operational plan; 2) rectify the impact by repairing or rehabilitating the affected environment; 3) reduce or eliminate the impact over time; and 4) compensate for any net wildlife losses created as a result of the impact. While the first three measures apply to reducing net losses, the latter provides a means for replacing resource losses that cannot be minimized or avoided.

The concept of compensation is based on the principle that wildlife resources are renewable and thus can be replenished through acquisition and management of suitable land. Habitat impacts and compensation credit may be quantified using habitat units (HUs). Habitat Units are calculated by multiplying habitat quality (HQ) by habitat quantity (acres of habitat). A tract of land has an existing inherent habitat value. To receive compensation credit, the land must be managed to increase its carrying capacity (as measured by HQ) so it can maintain existing wildlife populations while concurrently supporting additional populations to make up for the wildlife lost as a result of the project impacts. Management may be through physical modifications or treatments to improve the habitat or passive protection from disturbances, thus allowing natural succession. In either case, to obtain gains in habitat value for compensation from an acre of habitat, that acre must be managed to increase its existing habitat quality.



Legend

- Roads
- Reservoir Boundary
- Evaluation Site
- Little Sandy National Wildlife Refuge
- Old Sabine Bottom Wildlife Management Area (TPWD)



Scale



Figure 1 Location of Wildlife Habitat Evaluation Sites and State/Federal Lands

The formulas for determining acreage requirements to compensate for habitat losses are as follows:

Compensation acres needed = HUs lost from project ÷ Potential HQ Gain

Potential HQ Gain = Maximum HQ value for habitat type — Existing baseline HQ

For this analysis, it is assumed that the maximum HQ value obtained through management will be the maximum HQ score possible for the cover type. **The maximum HQ score for Forests is 1.0, the highest possible HQ for grasslands/crops is 0.92.** Obtaining the maximum potential HQ gain also assumes maximum (intensive) management. Compensation requirements were also calculated for the options of achieving 50% and 25% of the potential maximum HQ gain, implying moderate and minimum levels of management, respectively.

The above formulas can be applied very simply. If 10 acres of wildlife habitat with a habitat quality score of 0.6 were inundated by a reservoir, a total of 6 HUs would be lost [10 (acres) x 0.6 (HQ) = 6 (HUs)]. Full compensation to offset this loss would require an increase in habitat value of 6 HUs. If another 10 acres with an existing HQ of 0.6 were acquired, no compensation credit occurs because no increase in habitat value occurs over existing conditions. However, if by employment of habitat improvement measures, the existing habitat with an HQ score of 0.6 is increased by 0.4 to its maximum value of 1.0, then 15 acres would fully compensate [6 (HUs lost ÷ 0.4 (HQ increase) = 15 (compensation acres)]. If the existing HQ of the mitigation tract is only slightly raised through management to 0.7, then 60 acres would be required to fully compensate for the losses [6 HUs ÷ 0.1 (HQ increase) = 60 (compensation acres)]. These examples illustrate that compensation acreage can become significantly higher as the potential gain in habitat quality decreases. A summary of the Habitat Quality (HQ) assessment and compensation requirements is provided in Table 3.

Results and Discussion

Wildlife habitat appraisals were conducted for four cover types within the Waters Bluff Reservoir Site as identified in Table 2. Tabulated data is contained within Table 3. This information includes the name of the cover type evaluated, resource category of the cover type (in parenthesis following the cover type name), acres impacted within normal pool elevation, habitat quality rating obtained by field evaluation, habitat units lost, hypothetical management options, potential gain in habitat quality, and compensation requirements for each management option. Mitigation goals for habitat losses can be obtained by noting the resource category designation after the cover type name and referencing Table 1.

Table 3. Habitat Quality Assessment and Mitigation Requirements for the Proposed Waters Bluff Reservoir, Normal Pool Elevation = 300.0' msl.

Cover Type/ Resource Category ()	Acres Inventoried	Habitat Quality Value	Habitat Units Lost	Management Option	Potential Habitat Quality Gain	Compensation Requirements (Acres)
Mixed Bottomland Hardwood Forest (1 & 2) ¹	24,931	0.84	20,942	Minimum 25% Moderate 50% Maximum 100%	.040 .080 .160	523,550 261,775 130,888
Mixed Upland Forest (2 & 3)	1,091	0.75	818	Minimum 25% Moderate 50% Maximum 100%	.063 .125 .250	12,984 6,544 3,272
Grassland/Crops (3 & 4)	4,237	0.57	2,415	Minimum 25% Moderate 50% Maximum 100%	.088 .175 .350	27,443 13,800 6,900
Pine Plantation (4)	559	0.48	268	Minimum 25% Moderate 50% Maximum 100%	.130 .260 .520	2,062 1,031 515
Other	200					
Total	31,018			Minimum 25% Moderate 50% Maximum 100%		566,039 283,150 141,575

¹ Several bottomland hardwood sites were rated as unique due to exceptional large size and old growth age structure.

The total area inundated by Waters Bluff Reservoir at the proposed normal pool elevation is 31,018 acres. Mixed bottomland hardwood forests comprise 24,931 acres or 80% of the total area. Similar analyses using the WHAP appraisal technique and the HEP (Habitat Evaluation Procedure developed by the U.S. Fish and Wildlife Service) have been previously investigated (Frye and Curtis 1990) for 44 other reservoirs. In the 1990 study, HSI values obtained for particular wildlife species on previously conducted HEP evaluations were averaged for all the species occupying each cover type to provide numerical values compatible with WHAP. Data for this reservoir site was compared with information compiled for other reservoirs. With an average habitat quality rating of 0.84 the site ranks equally high in habitat quality with the Lindenau reservoir site in south central Texas, but has much more bottomland acreage than the 3,054 acres at Lindenau. In addition, the highest recorded individual habitat quality score (0.98) among all reservoirs was collected at the Waters Bluff site on the Little Sandy National Wildlife Refuge.

The influence of existing high habitat quality in calculating compensation requirements is significant. Assuming bottomland hardwood tracts of sufficient size exist and could be purchased from willing sellers, acquisition and intensive management of 130,888 acres would be required to fully compensate for in-kind losses. This represents 5.25 acres of replacement for each acre of bottomland forest impacted. For the entire reservoir project, 141,575 acres would be required to fully compensate for the loss of 30,818 acres of existing vegetation cover types or replacement of 4.59 acres for each acre lost. It is uncertain whether one or even several mitigation tracts could be found that could satisfactorily meet these mitigation requirements.

Planning Issues


Bottomland hardwood forests in Texas have been documented to contain at least 189 species of trees and shrubs, 42 woody vines, 75 grasses, and 802 herbaceous plants. This results in a food web that supports 116 species of fish, 31 species of amphibians, 54 species of reptiles, 273 species of birds, and 45 species of mammals. Many of these species would be expected to occur within the Waters Bluff Reservoir site, due to existing high vegetation species diversity and the presence of old growth forests.

Eastern Wild Turkeys have suffered major declines but are now increasing, partly through expensive releases the Department has completed at several locations within the reservoir site as part of its Eastern Wild Turkey Restoration Project. Loss of these established turkey populations will require additional compensation considerations.

Seventy-four species of threatened or endangered animals depend directly on bottomland forest systems. Bottomlands within the Waters Bluff reservoir site provide habitat for a number of rare species. Two rare species of bats, the southeastern myotis (*Myotis austroriparius*) and Rafinesque big-eared bat (*Corynorhinus rafinesquii*) occur as far west as the bottomland hardwood forests of east Texas. Declines in populations throughout other parts of their range due to disturbance and habitat loss make it imperative that habitat be protected in Texas. One maternity tree roost of the

southeastern myotis has been confirmed (Photo 1) with another suspected on the Little Sandy NWR. Another maternity tree roost has been confirmed farther downstream. In addition, several winter tree roosts have been discovered and continue to be monitored along the Sabine River corridor. It appears that these species prefer tree roosts in very large live bottomland hardwood trees where the base has been hollowed out. A disruption of this natural system could decrease available roost sites, and thus contribute to the further decline of this species.

Photo 1. Maternity Roost Tree of the Southeastern Myotis on the Little Sandy NWR.



Several tracts of particular sensitivity and importance lie within the reservoir site. These tracts were acquired by public conservation agencies for the express purpose of conserving bottomland hardwood forests and associated wetlands. The previously mentioned Little Sandy NWR is one of the few bottomland hardwood forest sites remaining in Texas (Figure 1). It contains 5 state champion trees and is of importance to many species of wildlife, especially thousands of wintering waterfowl. The refuge benefits the public by protecting and preserving forests of national value and importance. Old growth timber occurring on this site is illustrated by photo 2. Photo 3 illustrates old growth forest occurring on private land within the reservoir site.

Photo 2. Old Growth Bottomland Hardwood Forest on the Little Sandy National Wildlife Refuge.



Photo 3. Old Growth Bottomland Hardwood Forest occurring on private land within the reservoir site. Note cavities used by many species of wildlife for nesting and escape cover. Trees must reach mature or old growth stages before cavities commonly occur.



In addition to the ecological importance of the Little Sandy NWR, the Texas Parks and Wildlife Department's 5,158-acre Old Sabine Bottom Wildlife Management Area, which lies directly across the river from the Little Sandy NWR, has both ecological and mitigation values. A total of 2,694 acres was initially purchased by TPWD. An additional 2,464 acres of land was purchased by the Texas Department of Transportation. Lands acquired by TxDOT were transferred to Texas Parks and Wildlife for wildlife management purposes to provide future mitigation credits that can be used by TxDOT to offset wetland losses incurred by a number of planned highway projects in northeast Texas. Total initial investment by TPWD and TxDOT was 2.7 million dollars. This Department, TxDOT, Texas Natural Resources Conservation Commission, Corps of Engineers, Environmental Protection Agency, and the U.S. Fish and Wildlife Service worked cooperatively to establish this mitigation area with the intent of simultaneously conserving an important natural resource while addressing the requirements of a number of federal laws. TxDOT has also recently spent over 200 thousand dollars to upgrade existing roads on the area.

If it were constructed, the Waters Bluff Reservoir project would cover approximately 75% of the Old Sabine Bottom Wildlife Management area (Figure 1). Mitigation would be complicated since part of the area is already a mitigation site that is dedicated to compensating for other areas that have been altered or destroyed.

The high habitat values and level of impact expected from reservoir inundation will require substantial acquisition and management of forested wetland under any of the mitigation options.

Literature Cited

- Eyre, F.H. (ed). 1980. Forest cover types of the United States and Canada. Soc. Am. Foresters, Washington D.C. 146 pp.
- Frye, R.G. 1986. Bottomland hardwoods—current supply, status, habitat quality and future impacts from reservoirs. *In* McMahan, C.A., and R.G. Frye, eds. Bottomland Hardwoods in Texas - Proceedings of an Interagency Workshop on Status and Ecology, May 6-7, 1986. Tex. Parks and Wildl. Dep. Pub. PWD-7100-133-3/87. pp. 24-27.
- Frye, R.G. and D.A. Curtis. 1990. Texas water and wildlife- an assessment of direct impacts to wildlife habitat from future water development projects. Tex. Parks and Wildl. Dep. Pub. PWD-BK-7100-147-5/90.
- Gould, F.W., G.O. Hoffman, and C.A. Rechenthin. 1960. Vegetational areas of Texas. Tex. A&M Univ., Tex. Agric. Exp. Sta. Leaflet 492.
- Larkin, T.J., and G.W. Bomar. 1983. Climatic atlas of Texas. Tex. Dep. Water Res. LP-192, 149p.

- Larson, J.S., M.S. Bedinger, C.F. Bryan, S. Brown, R.T. Huffman, E.L. Miller, D.G. Rhodes, and B. A. Touchet. 1981. Transition from wetlands to uplands in southeastern bottomland hardwood forests. P. 225-273 *in* J.R. Clark and J. Benforado, eds. Wetlands of Bottomland Hardwood Forests. Proceeding of a Workshop on Bottomland Hardwood Forest Wetlands of the Southeastern United States. Lake Lanier, Georgia, June 1-5, 1980. Developments in Agriculture and Manage-Forest Ecology, Vol. 11. Elsevier Scientific Publishing Co. New York.
- McMahan, C.A., R.G. Frye, and K.L. Brown. 1984. The vegetation types of Texas including cropland. Tex. Parks and Wildl. Dep. Bull. 700-120, 40 p.
- U.S. Fish and Wildlife Service. 1985. Texas bottomland hardwood preservation program -final concept plan. U.S. Fish and Wildlife Service. Albuquerque, N.M. 378 pp.

Appendix 1
Wildlife Habitat Appraisal

**Biological Habitat Components
Evaluation Key**

Component 1 - Site Potential

Evaluate for all cover types.

<u>Criteria</u>^{2/}	<u>Value</u>
Substrate is composed or exhibits one or more of the following: 1) at least periodically supports predominately hydrophytic vegetation; 2) is predominately undrained hydric soil and supports or is capable of supporting hydrophytic vegetation; 3) is saturated with water or covered by shallow water during 1-2 months during the growing season of each year (swamps, bogs, marshes, and hardwood bottomlands exhibiting a high frequency of flooding).	25
Alluvial substrate although less hydric than above; only temporarily or intermittently inundated or saturated for short periods (higher terraces of hardwood bottoms, riparian drainages).	20
Uplands with thick surface layer (generally greater than or equal to 10 inches) consisting of unrestricted loam (including sandy loam) or dark well structured (granulated) clay (including sandy clay).	12
Uplands with shallow surface layer (generally less than 10 inches) consisting of shallow soil over restrictive layer (rock, gravel, claypan, etc.) or deep, leached, droughty sand or, relatively light colored, poorly structured clay or gravelly/stony sand or clay.	7
Organic matter minimal or absent at the surface. (Includes undrained or saturated hydric soils not supporting vegetation i.e., mud flats).	3
Surface contains chemical compounds which would potentially limit growth of primary producers (salt, mine overburden containing heavy metals or acid compounds, surface pollution).	1

Component 2 - Temporal Development of Existing Successional Stage

Determine currently existing successional stage (Criteria A); evaluate for all cover types except marshes. For this habitat type use Criteria B

<u>Criteria A</u> ^{3/}	<u>Value</u>
Old timber (100 or more years, trees > 25 inches*)	20
Mature timber, old brush, climax prairie (40-99 years, trees 12-25 inches)	12
Pole and young timber, mature brush (11-39 years, trees < 12 inches)	6
Grasslands in grazing disclimax** or early and mid-successional perennial grasses and forbs, hay meadows	5
Seedlings, saplings, young brush (3-10 years)	3
Annual native or introduced grasses, forbs, crops	1

* Diameter at breast height (DBH)

** Example: Texas wintergrass-silver bluestem grasslands

<u>Criteria B</u>	<u>Value</u>
(Marsh wetlands)	
Established mature communities within or adjacent to an enclosed coastal water body with a free connection to the sea and a measurable quantity of salt in its waters but with abundant or semi-abundant freshwater inflow (estuarine areas).	20
Established mature communities or intermediate to well advanced successional stages occurring in fresh, brackish, or saline environments; freshwater inflow limited to generally small tributaries and localized runoff or overflow from flood conditions.	10
Aquatic or semi-aquatic communities occurring in generally early to intermediate successional stages as a result of periodic changes in moisture gradients; highly dependent on seasonal weather conditions.	5

Component 3 - Uniqueness and Relative Abundance

1. Evaluate the habitat within the site according to the categories below. Enter the value on the Acquisition Components Evaluation Summary.

<u>Category</u>	<u>Value</u>
Highly valuable for wildlife and is very uncommon, unique or irreplaceable (USFWS Mitigation Resource Category 1)	20

^{3/}Corresponds to scarcity and abundance criteria as contained in U.S. Fish and Wildlife Service Mitigation Policy; Federal Register Vol. 46:15, Jan. 23, 1981.

Highly valuable for wildlife but is relatively scarce or becoming scarce (USFWS Mitigation Resource Category 2)	15
Exhibits high to medium value for wildlife and is relatively abundant (USFWS Mitigation Resource Category 3)	10
Exhibits medium to low value for wildlife and is relatively abundant (USFWS Mitigation Resource Category 4)	5
Exhibits very low wildlife value regardless of abundance or scarcity	0

Component 4 - Vegetation Species Diversity

Criteria A

Diversity of Woody Species

Evaluate the composition of readily observable woody species in the overstory, midstory, and understory by determining the number of species groups as represented by the following categories. Evaluate for all cover types except Swamps (Criteria C) and Marsh wetlands (Criteria D). Worksheet for Criteria A&B provided on page 25.

<u>Species Group</u> ^{4/}	<u>Examples</u>
Berry/Drupe	hackberry, mulberry, paw paw, hawthorn, winterberry, black haw, soapberry, persimmon, choke cherry, yaupon, dogwood, Am. beautyberry, greenbriar, dewberry, poison ivy, rattan vine, blackgum, grape, mulberry, holly, juniper, bumelia, huckleberry, sumac, Virginia creeper, sassafras, prickly ash, chinaberry, crab apple, agarito, lotebush, ivy tree vine
Legume/Pod	mesquite, locust, redbud, <u>Acacia</u> spp.
Acorn	white oak, red oak, live oak, water oak, willow oak, post oak, bur oak
Nut/Nutlike	hickory, pecan, walnut, wax myrtle, ironwood, ephidra
Samara (Winged Fruit)	elm, ash, box elder, maple, river birch
Cone	pine, cypress
Achene	sycamore, <u>Baccharis</u> spp., sandsage, <u>Clematis</u> spp., salt bush
All others (capsules, follicles, burrs, hairy seeds)	willow, cottonwood, sweetgum, salt cedar, yucca, cactus, buttonbush, sweetgum, bois d'arc, creosotebush

Value assigned is equivalent to the number of groups represented (Maximum=8, If none is represented then value is 0)

Criteria B**Total Number of Occurring Woody Species**

Determine the total number of readily observable woody species and assign value according to the following categories. Do not use for Swamps (Criteria C) or Marsh wetlands (Criteria D)

	<u>Value</u>
15 or more species	7
10-14 species	5
5-9 species	3
1-4 species	1
None occurring	0

Criteria C**Diversity of Vegetation in Swamps**

Evaluate swamp areas according to the following categories:^{5/}

	<u>Value</u>
Seasonally flooded mixed bottomland hardwoods; inundation resulting from freshwater inflow	15
Seasonally flooded vegetation dominated by cypress-tupelo; inundation resulting from freshwater inflow	10
Continually flooded or infrequent, abrasively flooded vegetation comprised of one or more species; inundation resulting from freshwater, brackish or saline inflow	6
Continually flooded vegetation; inundation resulting from stagnant or impounded freshwater, brackish, or saline water conditions	2

Criteria D**Diversity of Vegetation in Marshes and other similar wetland areas**

Determine the major types of wetland vegetation present according to the following categories: rooted emergent vegetation, rooted submergent vegetation, rooted vegetation with floating leaves, algal mat communities (microalgae), benthic or drifting seaweeds (macroalgae).

	<u>Value</u>
<u>High</u> - includes three or more of above categories.	20
<u>Medium</u> - includes two of the above categories.	15
<u>Low</u> - includes one of the above categories.	5

Component 5 - Vertical Vegetation Stratification⁶ⁱ

Evaluate canopy coverage of the following three categories of vegetation for all cover types except crops and marsh wetlands.

- Categories:
- 1) Vegetation greater than 12 feet high
 - 2) Vegetation 3-12 feet high
 - 3) Vegetation less than 3 feet high

<u>Criteria</u>	<u>Value</u>
All three categories present, each accounting for at least 25 percent of ground cover	5
Any two of the above categories present, each accounting for at least 25 percent of ground cover	4
Only one of the above categories present and accounting for at least 25 percent of ground cover	3
None of the categories together account for more than 25 percent of ground cover	1

Component 6 - Additional Structural Diversity Components

Evaluate for all cover types except crops. Determine the presence of brush piles, rock piles, rocky crevices, snags, fallen logs, thick grass cover, brambles or thickets according to the following categories.

<u>Criteria</u>	<u>Value</u>
<u>Abundant</u> - Three or more of the above components readily apparent and observable from most locations with the site	5
<u>Moderate</u> - Any of the above components present, and observable with very little search effort	3

Sparse - Any of the above components present, but occurring infrequently or requiring significant search effort to locate

1

Absent - None of the above components observed

0

Component 7 - Condition of Existing Vegetation - Other

Use: Criteria A&B for cover types (other than crops and marsh wetlands) containing woody and/or herbaceous vegetation.
Criteria C for cropland only.
Criteria D for marsh wetlands.

Criteria A

Value

Degree of utilization of woody vegetation by vertebrates and invertebrates

Not evident - little or no evidence of plant utilization

5

Moderate - plant utilization observable with minimal damage to leaves and/or stems

3

Severe - damage to leaves and/or stems readily observable

1

No woody vegetation present

0

Criteria B

Value

Availability of Herbaceous Vegetation. Do not evaluate for Crops (Criteria C) or Marsh Wetlands (Criteria D)

Good - Eight or more combined species of grasses and forbs readily observable.

5

Fair - Four to seven combined species of grasses and forbs readily observable

3

Poor - One to three combined species of grasses and forbs readily observable

1

None - Herbaceous vegetation lacking or absent

0

Criteria C

Value

Available Biomass (Evaluate for croplands only)

High - Biomass removed periodically, although not necessarily annually; removed biomass supplanted by other vegetation resulting from natural succession of invading species or overseeding of introduced species; (Ex. Rice or other crop on multi-year rotational system allowing for additional biomass accumulations between harvests).

10

Moderate - Most biomass removed annually or semi-annually but with some residual amount remaining during portions of the rotational period. Minimal bare ground conditions (Hay operations, crops grown for pasture or grazing, chiseled crops).

5

Low - Most biomass removed annually due to clean farming practices creating significant bare ground conditions (intensive row crop farming).

1

Criteria D

Value

Condition of Marsh Wetlands

Unaltered - Quality of water and/or associated vegetation good, no foreseeable danger of environmental intrusion including pollution, contamination, sedimentation, or stagnation

10

Stable - Quality of water and/or associated vegetation good, although evidence exists that pollution, contamination sedimentation or stagnation could occur in the future or has occurred in the past

5

Degraded - Quality of water and/or associated vegetation poor or declining or degradation imminent

1



Appendix 2
Field Evaluation Forms

WHAP
Biological Components
Field Evaluation Form

Project Waters Bluff Date: 11/10/17
Cover Type or Plant Association Shrubland

Habitat Components	* Photo #17			Components Points (From Key)				Total
	Site No.	#1	#2	#3				
1. Site Potential		22	22	20				64
2. Temporal Development								
Criteria A		3	5	1				9
Criteria B (Marsh Wetlands Only)								
3. Uniqueness and Relative Abundance		10	10					25
4. Vegetation Species Diversity								
Criteria A		4	5	5				14
Criteria B		3	5	6				14
Criteria C (Swamps Only)								
Criteria D (Marsh Wetlands Only)								
5. Vertical Stratification		3	4	4				11
6. Additional Structural Diversity Components		3	5	1				9
7. Condition of Existing Vegetation								
Criteria A (Woody Vegetation)		5	1	3				9
Criteria B (Herbaceous Vegetation)		5	5	5				15
Criteria C (Croplands Only)								
Criteria D (Marsh Wetlands Only)								

#1
LAT 32
LONG 117

#2
1132/33/51.50
W 112/15.74

#3
1132/32/51.50
W 112/15.74

Average Habitat Quality Score for all sites within this cover type = $\frac{\text{Total Points}}{\text{Total number of sites}} \times \frac{1}{100} = \frac{170}{100} = .567$

#1 Regeneration of Grass

#2 Low oil fruit day 10 recovery

Species Diversity Worksheet

Project Water BluffCover Type Grassland

Site 1	Site 2	Site 3	Site 4	Site 5	Site 6
Berry/Drupe D. Holly	Hawthorne Palmetto Sugaring C. Persimmon Yucca Simsia	Sugaring E. Red Oak Sassafras Hawthorne Dewberry			
Legume/Pod Locust	H. Locust				
Acorn Water Oak	Willow oak Sweetgum	Southern Red Oak Willow Oak			
Nut/Nutlike		Pecan			
Samara Green Ash Winged Elm	G. Ash C. Elm	C. Elm W. Elm			
Cone		Loblolly Pine			
Achene					
All Others	Sweetgum	Sweetgum			

WHAP
Biological Components
Field Evaluation Form

Project

Waters - Bluffs

Date:

11/10-11/97

Cover Type or Plant Association

Mixed Hardwood - Pine Forest

Habitat Components	Components Points (From Key)						
Site No.	HP-1	HP-2	HP-3				Total
1. Site Potential	20	20	12				52
2. Temporal Development							
Criteria A	12	20	12				44
Criteria B (Marsh Wetlands Only)							
3. Uniqueness and Relative Abundance	10	15	10				35
4. Vegetation Species Diversity							
Criteria A	6	6	6				18
Criteria B	7	7	7				21
Criteria C (Swamps Only)							
Criteria D (Marsh Wetlands Only)							
5. Vertical Stratification	5	5	5				15
6. Additional Structural Diversity Components	5	5	5				15
7. Condition of Existing Vegetation							
Criteria A (Woody Vegetation)	5	5	5				15
Criteria B (Herbaceous Vegetation)	1	3	5				9
Criteria C (Croplands Only)							
Criteria D (Marsh Wetlands Only)							

Average Habitat Quality Score for all sites within
this cover type = $\frac{\text{Total Points}}{\text{Total number of sites}} \times \frac{1}{100} =$

$\frac{71 + 86 + 67}{3} \times \frac{1}{100} =$

$\frac{1}{100} =$

.747

224

#1
N 52/35/17.79
W 95/14/32.42

#2
N 32/60/20.85
W 95/14/32.42

#3
N 32/33/5.10
W 95/03/7.67

Species Diversity Worksheet

Project Waters BluffCover Type Hardwood - Pine Forest

Site 1	Site 2	Site 3	Site 4	Site 5	Site 6
Berry/Drupe Am. Beautyberry S. Wax myrtle Black gum Palmetto C. Persimmon Rattan Vine P. Ivy Grape Smilax A. Holly Dogwood Possum Haw	Am. Beautyberry Red Mulberry Smilax Am. Holly Possum Haw E. Red Cedar Peppercorn St. Andrew's Cross Viburnum Palmetto Grape P. Ivy Rattan Vine Sassafras	Dogwood Sassafras Black gum Smilax A. Holly Grape Am. Beautyberry E. Red Cedar P. Ivy Possumhaw Yucca Peppercorn Palmetto C. Persimmon			
Legume/Pod					
Acorn Overcup oak Water oak W. Low oak	Water oak Willow oak So. Red Oak	Post Oak Water Oak Willow oak S. Red Oak			
Nut/Nutlike Mockernut Hickory Coccinifer B. Hickory	Hickory, B.	Wax Myrtle Hickory			
Samara Red Maple G. Ash	W. Elm Am. Elm	W. Elm A. Maple Am Elm			
Cone Loblolly Pine	Loblolly Pine	Loblolly Pine			
Achene					
All Others Sweetgum Buttonbush	Sweetgum	Sweetgum			

WHAP
Biological Components
Field Evaluation Form

Project Waters Bluff Date: 11-11-97
Cover Type or Plant Association Pine Plantation

Habitat Components		Components Points (From Key)						
	Site No.							Total
1. Site Potential	1	12						
2. Temporal Development								
Criteria A		5						
Criteria B (Marsh Wetlands Only)								
3. Uniqueness and Relative Abundance		3						
4. Vegetation Species Diversity								
Criteria A		6						
Criteria B		7						
Criteria C (Swamps Only)								
Criteria D (Marsh Wetlands Only)								
5. Vertical Stratification		4						
6. Additional Structural Diversity Components		3						
7. Condition of Existing Vegetation								
Criteria A (Woody Vegetation)		5						
Criteria B (Herbaceous Vegetation)		1						
Criteria C (Croplands Only)								
Criteria D (Marsh Wetlands Only)								

Average Habitat Quality Score for all sites within
this cover type = $\frac{\text{Total Points}}{\text{Total number of sites}} \times \frac{1}{100} = \frac{46}{100} = .46$

Species Diversity Worksheet

Project Waters BluffCover Type Pine Plantation

	Site 1	Site 2	Site 3	Site 4	Site 5	Site 6
Berry/Drupe <i>Devilberry</i> <i>Am Beautyberry</i> <i>Sassafras</i> <i>Elderberry</i> <i>Sugarberry</i> <i>Juniper</i> <i>Smilax</i> <i>Devilberry</i>						
Legume/Pod <i>Honey Locust</i>						
Acorn <i>Blackjack Oak</i> <i>Water Oak</i> <i>Willow Oak</i>						
Nut/Nutlike <i>Black Walnut</i> <i>Pecan</i>						
Samara <i>Am Elm</i>						
Cone <i>Loblolly Pine</i>						
Achene						
All Others						

WHAP
Biological Components
Field Evaluation Form

Project Waters Bluff Date: 11/10/97
Cover Type or Plant Association Bottomland Hardwood

Habitat Components	Components Points (From Key)								
	Site No.	Photo 2 B-1	Photo 3 B-2	Photo 4 B-3	Photo 5 B-4	Photo 6 B-5	Photo 7 B-6	Photo 8 B-7	Photo 9 Total
1. Site Potential		22	20	20	25	22	22	25	156
2. Temporal Development									
Criteria A		12	12	20	12	12	12	20	100
Criteria B (Marsh Wetlands Only)		-	-						
3. Uniqueness and Relative Abundance		15	15	15	15	15	15	20	110
4. Vegetation Species Diversity									
Criteria A		5	5	6	5	5	5	6	37
Criteria B		7	7	7	3	7	5	7	43
Criteria C (Swamps Only)		-	-						
Criteria D (Marsh Wetlands Only)		-	-						
5. Vertical Stratification		5	5	5	4	5	5	5	34
6. Additional Structural Diversity Components		5	5	5	3	5	5	5	33
7. Condition of Existing Vegetation									
Criteria A (Woody Vegetation)		3	3	5	5	5	3	5	29
Criteria B (Herbaceous Vegetation)		3	1	3	1	3	3	5	19
Criteria C (Croplands Only)		-	-						
Criteria D (Marsh Wetlands Only)		-	-						

Average Habitat Quality Score for all sites within this cover type = $\frac{\text{Total Points}}{\text{Total number of sites}} \times \frac{1}{100} = \frac{561}{670} \times \frac{1}{100} = .836$

.77 .73 .86 .73 .79 .75 .98 561
L. S. Smith
11/10/97 = .80

WHAP
Biological Components
Field Evaluation Form

Project Water Bluff
Cover Type or Plant Association Bottomland

Date: 11/11/97

Habitat Components	Components Points (From Key)						Total
Site No.	B ⁸	B ⁹	B ¹⁰	B ¹¹	B ¹²	B ¹³	
1. Site Potential	25	23	20	25	23	25	141
2. Temporal Development							
Criteria A	20	20	15	20	20	9	104
Criteria B (Marsh Wetlands Only)							
3. Uniqueness and Relative Abundance	20	20	15	20	20	15	110
4. Vegetation Species Diversity							
Criteria A	5	3	5	6	6	5	30
Criteria B	7	7	7	7	7	5	40
Criteria C (Swamps Only)							
Criteria D (Marsh Wetlands Only)							
5. Vertical Stratification	5	5	5	5	5	3	28
6. Additional Structural Diversity Components	5	5	5	5	5	3	28
7. Condition of Existing Vegetation							
Criteria A (Woody Vegetation)	5	5	3	5	5	5	28
Criteria B (Herbaceous Vegetation)	5	3	3	1	5	1	18
Criteria C (Croplands Only)							
Criteria D (Marsh Wetlands Only)							527

Average Habitat Quality Score for all sites within
this cover type = $\frac{\text{Total Points}}{\text{Total number of sites}} \times \frac{1}{100}$

$\frac{527}{6} \times \frac{1}{100} = .836$

$\frac{1}{100}$

$.836$

Total for #10 N32/32/52.02
W05/12/19.7

#9 N32/30/25.32
W05/10/27.73

Species Diversity Worksheet

Project Water BluffCover Type Mixed Bottomland Hardwood Forest

	Site 1	Site 2	Site 3	Site 4	Site 5	Site 6
Berry/Drupe	Pocumbaw Smilax Rattan Vine Sugar Hackberry Poison Ivy Red Mulberry Japanese Honeyuckle Ilex Smilax	Mexican Plum Pepper Vine Palmetto Pocumbaw Sugar Hackberry Poison Ivy Rattan Vine Red Mulberry Ilex Smilax Red Cedar	A. Holly Sugarberry Rattan Vine Grape Dec. Holly Smilax Smilax 1 Smilax 2 Peppercorn Mulberry Palmetto	Common Persimmon Swamp Privet	Smilax Poison Ivy Poison Ivy R. Mulberry Sugarberry Peppercorn Rattan Vine	P. Ivy Rattan Vine Sugarberry Smilax Poison Ivy Red Mulberry Dewberry
Legume/Pod		Red bud				
Acorn oak Willow oak		Overcup Oak Willow Oak	Water Oak Overcup.	Overcup Oak	Willow oak Water Oak	Willow Oak Overcup Oak
Nut/Nutlike Hickory			Carpinus W. Hickory	Water Elm Butternut		Carpinus
Samara Cedar Elm Green Oak Am Elm		Green Elm Winged Elm Cedar Elm	Cedar Elm Green Ash Am Elm		Gr. Ash Cedar Elm Red Maple Am Elm	Gr. Ash C. Elm
Cone						
Achene Teardrop Vine		Teardrop Vine	Teardrop Vine	Teardrop Vine	Teardrop Vine	Teardrop Vine
All Others			G. Lane Sweetgum	B. Willow	Sweetgum	

Species Diversity Worksheet

Project Water BluffCover Type Mixed Bottomland Forest

	7 Site 1	8 Site 2	9 Site 3	10 Site 4	Site 5	Site 6
Berry/Drupe	Red Mulberry Hawthorne Blackgum Palmetto P. Ivy Rattanvine Smilax Sugarberry Grape Dewberry	Blackgum A. Holly Palmetto Rattanvine P. Ivy Smilax Sugarberry Red Mulberry Grape Dogwood Poplarvine	Possumhaw Smilax Sugarberry Palmetto Grape Smilax Dewberry St. Andrew's Peppercorn Rattanvine	Sugar Berry Possumhaw Smilax #1 Peppercorn P. Ivy R. Mulberry Smilax #2 Grape C. manneal St. Andrew's	Possumhaw Rattanvine Smilax 1 Smilax 2 P. Ivy Palmetto Rattanvine Blackgum Sugarberry Hawthorn	Possumhaw Palmetto Smilax 1 Smilax 2 Rattan V. P. Ivy Peppercorn Hawthorne
Legume/Pod	Red Bud					Eve's Neckhane
Willow oak Acom Water oak	Water oak	Willow oak Overcup oak Water oak B. Post oak	Overcup Willow oak	Willow oak Overcup oak Water oak	Willow oak Overcup oak	Willow oak Overcup oak
Nut/Nutlike Carpinus	Black Hickory Carpinus Bitter Pecan			Bitter Pecan Carpinus	Bitter Pecan Hickory	Carpinus B. Halpecan
Green Ash Samara W. Elm Red Maple	W. Elm Am Elm River Birch Red Maple	Am Elm C. Elm	Water Elm Am Elm C. Elm	C. Elm A. Elm	C. Elm G. Ash	Loblolly Pine
Cone					Loblolly Pine	Loblolly Pine
Achene						
All Others Sweetgum G. Lane	Sweetgum Black Willow			Sweetgum	Sweetgum G. Lane	Sweetgum

Species Diversity Worksheet

Project Water BluffCover Type Bottomland

Site 1 ¹³	Site 2	Site 3	Site 4	Site 5	Site 6
Berry/Drupe Palmetto Passum Haw C. Passum Smilax Popovine P. Fry					
Legume/Pod					
Acorn Willow Oak Overcup Oak					
Nut/Nutlike B. Ha Pecan					
Samara C. Elm					
Cone					
Achene					
All Others Sweet gum					

Attachment: A Copy of TWDB Comments on TPWD Draft Report

TWDB COMMENTS ON TPWD DRAFT REPORT: ANALYSIS OF BOTTOMLAND
HARDWOOD AREAS AND ASSESSMENT OF WILDLIFE HABITAT QUALITY AT
THE POTENTIAL FUTURE WATERS BLUFF RESERVOIR SITE
(TWDB CONTRACT NO. 97-483-226)

All tasks in the contract scope of work have been addressed. A few specific comments are detailed below and should be considered in preparing the Final Report for submittal to the Texas Water Development Board. Overall, the draft report supplies data and information that meets or exceeds the contract requirements.

Part One, page one, Introduction: Several federal agencies recognize a wetland type called “bottomland hardwood forest.” When this type of wetland is altered or destroyed by construction of a water development project, mitigation is required and may involve compensatory land acquisition to make up for the wetland loss. The exact area that will require mitigation is determined through a federal delineation of the wetlands, which employs three major criteria—the vegetation present, the hydroperiod of the area, and the presence of hydric soils. However, the draft report also recognizes a land cover type called “bottomland hardwood forest,” which is based only on a vegetation analysis using satellite imagery, aerial photography, and ground-truthing selected sites. Since no hydroperiod or hydric soils information was used in TPWD’s determination, a clarifying statement should be included in the report’s Introduction which explains to readers that bottomland hardwood forest areas identified in the report are not exactly the same as those that might be identified using the federal delineation procedures.

Part One, page 9, line 3: The contents of Table 1 could use a little more explanation. It was not clear to some reviewers which axis in the table represents the classified image (horizontal axis?) and which axis represents the ground-truth determination (vertical axis?). In addition, please include a brief description of how the overall 83.1% accuracy was calculated.

Part Two, page 1, line 4: Add the word “Board” after the words “Texas Water Development.”

Part Two, page 2, line 3: Note that Appendix 1 and Appendix 2 were not included in the review copy of the draft report; therefore, TWDB requests copies of the appendices to review before the report is finalized.

Part Two, page 4, line 11: There seems to be a font problem with the word “Area.”

Part Two, page 8, line 11: The text refers to a tree roost for the “southeastern” myotis, but the caption on Photo 1 refers to the “eastern” myotis. Shouldn’t these common names be the same?