

Table 1.9 Heating Degree-Days by Census Division

Census Divisions	July				
	Normal ^a	2011	2012	Percent Change	
				Normal to 2012	2011 to 2012
New England Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, Vermont	11	4	5	NM	NM
Middle Atlantic New Jersey, New York, Pennsylvania	6	0	1	NM	NM
East North Central Illinois, Indiana, Michigan, Ohio, Wisconsin	9	1	0	NM	NM
West North Central Iowa, Kansas, Minnesota, Missouri, Nebraska, North Dakota, South Dakota	15	2	1	NM	NM
South Atlantic Delaware, Florida, Georgia, Maryland and the District of Columbia, North Carolina, South Carolina, Virginia, West Virginia	0	0	0	NM	NM
East South Central Alabama, Kentucky, Mississippi, Tennessee	0	0	0	NM	NM
West South Central Arkansas, Louisiana, Oklahoma, Texas	0	0	0	NM	NM
Mountain Arizona, Colorado, Idaho, Montana, Nevada, New Mexico, Utah, Wyoming	19	1	0	NM	NM
Pacific^b California, Oregon, Washington	24	21	11	NM	NM
U.S. Average^b	9	4	2	NM	NM

^a "Normal" is based on calculations of data from 1971 through 2000.

^b Excludes Alaska and Hawaii.

NM=Not meaningful (because "Normal" is less than 100 or ratio is incalculable).

Notes: Degree-days are relative measurements of outdoor air temperature used as an index for heating and cooling energy requirements. Heating degree-days are the number of degrees that the daily average temperature falls below 65° F. Cooling degree-days are the number of degrees that the daily average temperature rises above 65° F. The daily average temperature is the mean of the maximum and minimum temperatures in a 24-hour period. For example, a weather station recording an average daily temperature of 40° F would report 25 heating degree-days for that day (and 0 cooling degree-days). If a weather station recorded an average daily temperature of 78° F, cooling degree-days for that station would be 13 (and 0 heating degree days).

Web Pages: • See <http://www.eia.gov/totalenergy/data/monthly/#summary>

for current data. • See <http://www.eia.gov/totalenergy/data/annual/#summary> for historical data.

Sources: There are several degree-day databases maintained by the National Oceanic and Atmospheric Administration. The information published here is developed by the National Weather Service Climate Prediction Center, Camp Springs, MD. The data are available weekly with monthly summaries and are based on mean daily temperatures recorded at about 200 major weather stations around the country. The temperature information recorded at those weather stations is used to calculate statewide degree-day averages based on population. The State figures are then aggregated into Census Divisions and into the national average. The population weights currently used represent resident State population data estimated for the 2000 Census by the U.S. Department of Commerce, Bureau of the Census. The data provided here are available sooner than the Historical Climatology Series 5-1 (heating degree-days) developed by the National Climatic Data Center, Asheville, NC, which compiles data from some 8,000 weather stations.