Office of Applied Studies U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES

Substance Abuse and Mental Health Services Administration http://dawninfo.samhsa.gov/

DRUG ABUSE WARNING NETWORK, 2005: SELECTED TABLES OF NATIONAL ESTIMATES OF DRUG-RELATED EMERGENCY DEPARTMENT VISITS

In August 2008, the Substance Abuse and Mental Health Services Administration (SAMHSA) released *Drug Abuse Warning Network, 2006: National Estimates of Drug-Related Emergency Department Visits.* This annual report included estimates for 2006 and comparisons of the emergency department estimates for 2004, 2005, and 2006. This report may be accessed on the Web at http://dawninfo.samhsa.gov/files/ED2006/DAWN2k6ED.pdf.

The 2006 estimates reflect several improvements made to the Drug Abuse Warning Network (DAWN) sampling and estimation methodology. To ensure comparability between estimates for 2004, 2005, and 2006, the improvements were applied retrospectively to 2004 and 2005 data, resulting in small changes as compared with early published estimates.

Tables for 2005 that have not already been published in the 2006 ED report are now available online. The 2005 tables and charts follow the same title and numbering system as the 2006 ED report. Links to the revised 2005 tables are provided below.

General information about DAWN is at http://dawninfo.samhsa.gov/. Information on other sources of data on substance abuse and mental health from SAMHSA's Office of Applied Studies (OAS) is located at http://oas.samhsa.gov/.

LIST OF TABLES

Table 1	Drug misuse and abuse in ED visits in the U.S., by type of drug involvement: 2005
Table 2	Illicit drugs in ED visits: 2005
Table 3	Illicit drugs, by patient characteristics: 2005
Table 4	Alcohol in drug-related ED visits: 2005
Table 5	Alcohol in combination, by patient and visit characteristics: 2005
Table 6	Drugs most frequently reported with alcohol: 2005
Table 7	Alcohol in drug-related ED visits in patients under age 21: 2005
Table 8	Alcohol only (age < 21), by patient and visit characteristics: 2005

Table 9	Nonmedical use of pharmaceuticals: 2005
Table 10	Nonmedical use of pharmaceuticals, by patient and visit characteristics: 2005
Table 16	Suicide attempts: 2005
Table 17	Suicide attempts, by patient and visit characteristics: 2005
Table 19	Seeking detox: 2005
Table 20	Seeking detox, by patient and visit characteristics: 2005
Table C2	Drug-related ED visits, by type of case: 2005
Table C5	Population by age and gender: 2005

PUBLIC DOMAIN NOTICE

All material appearing in these tables is in the public domain and may be reproduced or copied without permission from SAMHSA. However, this publication may not be reproduced or distributed for a fee without the specific, written authorization of the Office of Communications, SAMHSA. Citation of the source is appreciated. Suggested citation:

Substance Abuse and Mental Health Services Administration, Office of Applied Studies. *Drug Abuse Warning Network, 2005: Selected Tables of National Estimates of Drug-Related Emergency Department Visits.*Rockville, MD, 2009.

This publication was prepared by the Substance Abuse and Mental Health Services Administration, Office of Applied Studies, and by RTI International under Contract No. 280-03-2602.

ORIGINATING OFFICE

Office of Applied Studies
Substance Abuse and Mental Health Services Administration
1 Choke Cherry Road, Room 7-1044, Rockville, MD 20857

DATE OF ORIGINAL POSTING

June 2009

Table 1
Drug misuse and abuse in ED visits in the U.S., by type of drug involvement: 2005

	Estimated	Davaget	Relative	95% Confidence interval			
Drug involvement ¹	visits ²	Percent of visits	standard error (RSE)	Lower bound	-	Upper bound	
All types of drug misuse/abuse	1,616,311	100%	9.1	1,329,073	-	1,903,548	
Illicit drugs only	517,558	32%	17.5	339,699	-	695,417	
Alcohol only (age < 21)	110,599	7%	11.3	86,147	-	135,052	
Pharmaceuticals only	444,309	27%	7.2	381,282	-	507,337	
Combinations							
Illicit drugs with alcohol ³	221,823	14%	19.2	138,406	-	305,240	
Illicit drugs with pharmaceuticals	127,245	8%	13.6	93,304	-	161,186	
Alcohol with pharmaceuticals	140,275	9%	6.7	121,764	-	158,787	
Illicit drugs with alcohol and pharmaceuticals	54,500	3%	13.9	39,601	-	69,400	

¹ The classification of drugs used in DAWN is derived from the Multum *Lexicon*, © 2007, Multum Information Services, Inc. The classification was modified to meet DAWN's unique requirements (2007). The Multum Licensing Agreement governing use of the *Lexicon* can be found on the Internet at http://www.multum.com.

² These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

³ DAWN excludes alcohol-only visits for adults. Alcohol, when present with other drugs, is included for all ages.

Table 2 Illicit drugs in ED visits: 2005

	Estimated	Relative	95% Confi	iden	ce interval
Drug category and selected drugs ¹	visits ^{2,3,4}	standard error (RSE)	Lower bound	-	Upper bound
Drug-rel	ated ED visits				
Total drug misuse/abuse ED visits	1,616,311	9.1	1,329,073	-	1,903,548
ED visits, illicit drugs	921,127	15.2	647,030	-	1,195,224
Cocaine	483,865	16.5	327,426	-	640,304
Heroin	187,493	19.5	115,953	-	259,034
Marijuana	279,664	12.4	211,755	-	347,572
Stimulants	137,650	24.7	71,051	-	204,248
Amphetamines	34,928	15.8	24,102	-	45,753
Methamphetamine	109,655	28.9	47,540	-	171,770
MDMA (Ecstasy)	11,287	16.2	7,695	-	14,879
GHB	1,036	19.5	640	-	1,431
Flunitrazepam (Rohypnol)				-	
Ketamine	303	32.1	112	-	494
LSD	2,001	19.6	1,231	-	2,771
PCP	14,825	33.0	5,224	-	24,427
Miscellaneous hallucinogens	3,194	19.2	1,991	-	4,398
Inhalants	5,156	21.6	2,968	-	7,344
Combinations not tabulated above (NTA)	3,201	39.3	732	-	5,669

Table 2 (continued)

Illicit drugs in ED visits: 2005

	ED visits	Relative	95% Confidence interva		
Drug category and selected drugs ¹	per 100,000 population ^{2,3,4}	standard error (RSE)	Lower bound	-	Upper bound
ED visits per 1	00,000 population				
Total drug misuse/abuse ED visits	545.3	9.1	448.0	-	642.6
ED visits, illicit drugs	310.8	15.2	218.2	-	403.3
Cocaine	163.2	16.5	110.4	-	216.0
Heroin	63.3	19.5	39.1	-	87.4
Marijuana	94.4	12.4	71.4	-	117.3
Stimulants	46.4	24.7	24.0	-	68.9
Amphetamines	11.8	15.8	8.1	-	15.4
Methamphetamine	37.0	28.9	16.0	-	57.9
MDMA (Ecstasy)	3.8	16.2	2.6	-	5.0
GHB	0.3	19.5	0.2	-	0.5
Flunitrazepam (Rohypnol)				-	
Ketamine	0.1	32.1		-	
LSD	0.7	19.6	0.4	-	0.9
PCP	5.0	33.0	1.8	-	8.2
Miscellaneous hallucinogens	1.1	19.2	0.7	-	1.5
Inhalants	1.7	21.6	1.0	-	2.5
Combinations not tabulated above (NTA)	1.1	39.3		-	

¹ The classification of drugs used in DAWN is derived from the Multum *Lexicon*, © 2007, Multum Information Services, Inc. The classification was modified to meet DAWN's unique requirements (2007). The Multum Licensing Agreement governing use of the *Lexicon* can be found on the Internet at http://www.multum.com.

² These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

³ ED visits often involve multiple drugs. Such visits will appear multiple times in this table (e.g., a visit involving both cocaine and heroin will appear twice in this table). Summing ED visits as reported will produce incorrect and inflated counts of ED visits.

⁴ Three dots (...) indicate that an estimate with an RSE greater than 50% or an estimate less than 30 has been suppressed. **SOURCE**: Office of Applied Studies, SAMHSA, Drug Abuse Warning Network, 2005 (03/2008 update).

Table 3 Illicit drugs, by patient characteristics: 2005

Patient				Se	elected drug	s ¹			
characteristics	All illicits	Cocaine	Heroin	Marijuana	Stimulants	MDMA (Ecstasy)	GHB	LSD	PCP
Drug-related ED visits ^{2,3}									
ED visits, illicit drugs	921,127	483,865	187,493	279,664	137,650	11,287	1,036	2,001	14,825
Gender									
Male	606,714	308,559	131,665	187,440	89,015	7,009	796	1,716	10,721
Female	314,037	175,191	55,787	92,081	48,515	4,274	240	285	4,007
Unknown	375	116							
Age									
0-5 years	1,102	106							
6-11 years	557								
12-17 years	53,400	7,848	1,033	38,607	12,028	1,929			691
18-20 years	70,746	21,112	9,360	38,254	14,055	2,528		343	
21-24 years	105,420	39,468	19,546	44,597	23,751	2,206		488	2,288
25-29 years	113,258	53,136	22,570	41,480	23,186	2,512	277	250	3,384
30-34 years	114,255	62,640	24,053	31,574	18,880	947	119		
35-44 years	261,943	174,023	54,218	50,131	31,071	877	283		3,651
45-54 years	161,962	105,165	43,301	28,107	11,712				946
55-64 years	31,914	16,348	11,504	5,498	1,999				
65 years and older	5,320	2,834		714					
Unknown	1,250		71						
Race/ethnicity									
White	402,160	177,820	80,522	147,087	79,085	5,053	633	1,194	5,851
Black	283,519	203,124	47,537	66,877	8,413	2,298			6,218
Hispanic	99,126	38,881	26,719	26,220		1,017			1,074
Race/ethnicity not tabulated above (NTA)	9,117	4,016	1,119	2,688		468			
Unknown	127,206	60,025	31,596	36,792	21,796	2,451	187	468	1,589

Table 3 (continued)
Illicit drugs, by patient characteristics: 2005

Patient		Selected drugs ¹								
characteristics	All illicits	Cocaine	Heroin	Marijuana	Stimulants	MDMA (Ecstasy)	GHB	LSD	PCP	
			ED visits	per 100,000) population	2,3			_	
ED visits, illicit drugs	310.8	163.2	63.3	94.4	46.4	3.8	0.3	0.7	5.0	
Gender										
Male	415.6	211.3	90.2	128.4	61.0	4.8	0.5	1.2	7.3	
Female	208.8	116.5	37.1	61.2	32.3	2.8	0.2	0.2	2.7	
Age										
0-5 years	4.6	0.4								
6-11 years	2.3									
12-17 years	209.5	30.8	4.1	151.5	47.2	7.6			2.7	
18-20 years	568.2	169.6	75.2	307.3	112.9	20.3		2.8		
21-24 years	625.4	234.1	116.0	264.6	140.9	13.1		2.9	13.6	
25-29 years	564.4	264.8	112.5	206.7	115.6	12.5	1.4	1.2	16.9	
30-34 years	569.1	312.0	119.8	157.3	94.0	4.7	0.6			
35-44 years	597.2	396.7	123.6	114.3	70.8	2.0	0.6		8.3	
45-54 years	381.2	247.6	101.9	66.2	27.6				2.2	
55-64 years	105.1	53.9	37.9	18.1	6.6					
65 years and older	14.5	7.7		1.9						

¹ The classification of drugs used in DAWN is derived from the Multum *Lexicon*, © 2007, Multum Information Services, Inc. The classification was modified to meet DAWN's unique requirements (2007). The Multum Licensing Agreement governing use of the *Lexicon* can be found on the Internet at http://www.multum.com.

² These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

³ Three dots (...) indicate that an estimate with an RSE greater than 50% or an estimate less than 30 has been suppressed. **SOURCE**: Office of Applied Studies, SAMHSA, Drug Abuse Warning Network, 2005 (03/2008 update).

Table 4
Alcohol in drug-related ED visits: 2005

	Estimated Relative		95% Confidence interval			
Drug category and selected drugs ¹	visits ^{2,3}	standard error (RSE)	Lower bound	-	Upper bound	
Total drug misuse/abuse ED visits	1,616,311	9.1	1,329,073	•	1,903,549	
ED visits, alcohol	527,198	10.0	424,345	-	630,052	
Alcohol in combination	416,599	11.1	325,805	-	507,392	
Alcohol alone	110,599	11.3	86,147	-	135,052	

The classification of drugs used in DAWN is derived from the Multum *Lexicon*, © 2007, Multum Information Services, Inc. The classification was modified to meet DAWN's unique requirements (2007). The Multum Licensing Agreement governing use of the *Lexicon* can be found on the Internet at http://www.multum.com.

² These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

³ Estimates are all expressed in visits.

Table 5
Alcohol in combination, by patient and visit characteristics: 2005

Patient characteristics	Estimated visits ^{1,2}	Visit characteristics	Estimated visits ^{1,2}
ED visits, alcohol in combination	416,599		
Gender		Number of drugs involved	
Male	261,482	Single drug	
Female	155,042	Multiple drugs	416,599
Unknown	***	Alcohol involved	416,599
Age		Disposition	
0-5 years	•••	Treated and released	217,029
6-11 years	***	Discharged home	176,516
12-17 years	19,720	Released to police/jail	12,250
18-20 years	27,784	Referred to detox/treatment	28,263
21-24 years	45,688	Admitted to this hospital	136,951
25-29 years	49,962	ICU/critical care	26,310
30-34 years	45,275	Surgery	267
35-44 years	122,860	Chemical dependency/detox	15,506
45-54 years	82,520	Psychiatric unit	48,065
55-64 years	15,460	Other inpatient unit	46,803
65 years and older	6,412	Other disposition	62,619
Unknown		Transferred	38,987
Race/ethnicity		Left against medical advice	8,598
White	223,734	Died	
Black	94,620	Other	
Hispanic	40,413	Not documented	4,042
Race/ethnicity not tabulated above (NTA)	3,428		
Unknown	54,403		

¹ These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

² Three dots (...) indicate that an estimate with an RSE greater than 50% or an estimate less than 30 has been suppressed. **SOURCE**: Office of Applied Studies, SAMHSA, Drug Abuse Warning Network, 2005 (03/2008 update).

Table 6
Drugs most frequently reported with alcohol: 2005

Drugs reported with alcohol ¹	Estimated visits ²
No other drug	110,559
Cocaine only	95,557
Marijuana only	40,483
Cocaine and marijuana only	19,552
Heroin only	17,123
Stimulants only	15,415
Alprazolam only	8,005
Cocaine and heroin only	9,306

¹ The classification of drugs used in DAWN is derived from the Multum *Lexicon*, © 2007, Multum Information Services, Inc. The classification was modified to meet DAWN's unique requirements (2007). The Multum Licensing Agreement governing use of the *Lexicon* can be found on the Internet at http://www.multum.com.

² These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

Table 7
Alcohol in drug-related ED visits in patients under age 21: 2005

	Estimated	Relative	95% Confidence interval			
Drug category and selected drugs ¹	visits ²	standard error (RSE)	Lower bound	-	Upper bound	
Pati	ients aged 12-17				_	
ED visits, alcohol	62,459	9.7	50,578	-	74,339	
Alcohol in combination	19,720	9.5	16,066	-	23,373	
Alcohol alone	42,739	11.6	32,983	-	52,495	
Pati	ients aged 18-20				_	
ED visits, alcohol	95,166	11.7	73,382	-	116,949	
Alcohol in combination	27,784	12.2	21,146	-	34,422	
Alcohol alone	67,382	13.0	50,154	-	84,610	

¹ The classification of drugs used in DAWN is derived from the Multum *Lexicon*, © 2007, Multum Information Services, Inc. The classification was modified to meet DAWN's unique requirements (2007). The Multum Licensing Agreement governing use of the *Lexicon* can be found on the Internet at http://www.multum.com.

² These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

Table 8 Alcohol only (age < 21), by patient and visit characteristics: 2005

Patient characteristics	Estimated visits ^{1,2}	Visit characteristics	Estimated visits ^{1,2}
ED visits, alcohol only (age < 21) ³	109,174		
Gender		Number of drugs involved	
Male	61,973	Single drug	109,174
Female	47,194	Multiple drugs	
Unknown		Alcohol involved	109,174
Age		Disposition	
0-5 years		Treated and released	97,942
6-11 years		Discharged home	90,025
12-17 years	42,259	Released to police/jail	6,455
18-20 years	66,437	Referred to detox/treatment	1,463
21-24 years		Admitted to this hospital	5,569
25-29 years		ICU/critical care	1,209
30-34 years		Surgery	174
35-44 years		Chemical dependency/detox	
45-54 years		Psychiatric unit	788
55-64 years		Other inpatient unit	3,162
65 years and older		Other disposition	5,663
Unknown		Transferred	3,014
Race/ethnicity		Left against medical advice	1,063
White	62,911	Died	
Black	7,860	Other	
Hispanic	18,689	Not documented	746
Race/ethnicity not tabulated above (NTA)	1,632		
Unknown	18,083	1	

¹ These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

² Three dots (...) indicate that an estimate with an RSE greater than 50% or an estimate less than 30 has been suppressed.

³ This table is limited to ED visits classified as "alcohol only (age < 21)" and excludes visits classified as either "suicide attempt" or "seeking detox." Therefore, the estimate of total visits is slightly lower than reported in Table 4.

Table 9 Nonmedical use of pharmaceuticals: 2005

	Estimated	Relative	95% Confidence interval			
Selected drug categories and selected drugs ¹	visits ^{2,3,4}	standard error (RSE)	Lower bound	-	Upper bound	
ED visits, nonmedical use	669,214	7.0	577,800	-	760,628	
PSYCHOTHERAPEUTIC AGENTS	308,655	8.3	258,552	-	358,758	
Antidepressants	67,051	8.2	56,287	-	77,816	
MAO inhibitors				-		
SSRI antidepressants	30,374	11.6	23,467	-	37,280	
Tricyclic antidepressants	14,515	21.2	8,482	-	20,548	
Miscellaneous antidepressants	7,452	18.9	4,697	-	10,206	
Antipsychotics	44,393	9.7	35,922	-	52,864	
Anxiolytics, sedatives, and hypnotics	227,486	10.6	180,381	-	274,590	
Barbiturates	14,693	27.6	6,735	-	22,652	
Benzodiazepines	189,704	11.6	146,433	-	232,974	
Alprazolam	57,419	11.0	45,033	-	69,806	
Clonazepam	30,648	10.9	24,111	-	37,185	
Diazepam	18,433	14.1	13,350	-	23,516	
Lorazepam	23,210	13.4	17,132	-	29,288	
Benzodiazepines NOS	61,486	29.1	26,461	-	96,510	
Misc. anxiolytics, sedatives, and hypnotics	35,561	12.0	27,214	-	43,908	
Diphenhydramine	10,294	13.7	7,534	-	13,055	
Hydroxyzine	2,179	25.8	1,078	-	3,279	
Zolpidem	14,730	16.2	10,050	-	19,410	
Anxiolytics, sedatives, and hypnotics NOS	4,421	19.3	2,751	-	6,091	
CNS stimulants	10,965	12.4	8,307	-	13,623	
Amphetamine-dextroamphetamine	2,669	23.9	1,417	-	3,920	
Caffeine	4,567	16.6	3,080	-	6,054	
Dextroamphetamine				-		
Methylphenidate	2,519	21.1	1,478	-	3,560	

Table 9 (continued)
Nonmedical use of pharmaceuticals: 2005

	Estimated	Relative	95% Confidence interval			
Selected drug categories and selected drugs ¹	visits ^{2,3,4}	standard error (RSE)	Lower bound	-	Upper bound	
CENTRAL NERVOUS SYSTEM AGENTS	336,900	8.2	282,659	-	391,142	
Analgesics	294,251	8.5	245,312	-	343,191	
Antimigraine agents	1,018	26.1	497	-	1,539	
Cox-2 inhibitors	765	34.2	253	-	1,276	
Opiates/opioids	217,594	10.4	173,143	-	262,045	
Opiates/opioids, unspecified	52,670	19.3	32,702	-	72,638	
Narcotic analgesics	168,376	10.5	133,645	-	203,108	
Buprenorphine/combinations		52.3		-		
Codeine/combinations	6,180	11.9	4,737	-	7,622	
Fentanyl/combinations	11,211	24.2	5,904	-	16,519	
Hydrocodone/combinations	47,192	15.0	33,335	-	61,048	
Hydromorphone/combinations	4,714	22.9	2,595	-	6,83	
Meperidine/combinations	383	39.0	90	-	67	
Methadone	42,684	14.5	30,520	-	54,84	
Morphine/combinations	15,762	26.5	7,563	-	23,96	
Oxycodone/combinations	52,943	13.3	39,157	-	66,72	
Propoxyphene/combinations	7,648	15.4	5,335	-	9,96	
Nonsteroidal anti-inflammatory agents	28,837	9.4	23,541	-	34,13	
Ibuprofen	22,268	11.1	17,428	-	27,10	
Naproxen	5,190	12.2	3,951	-	6,42	
Salicylates/combinations	12,123	10.7	9,569	-	14,67	
Miscellaneous analgesics/combinations	51,881	8.0	43,699	-	60,06	
Acetaminophen/combinations	43,558	8.3	36,434	-	50,68	
Tramadol/combinations	5,918	23.6	3,178	-	8,65	
Tramadol	5,427	26.9	2,569	-	8,28	
Acetaminophen-tramadol		61.0		-		
Analgesic combinations not tabulated above (NTA)	653	29.7	273	-	1,032	
Anorexiants	1,757	20.9	1,038	-	2,476	
Anticonvulsants	27,641	8.8	22,897	-	32,38	
Antiemetic/antivertigo agents	1,771	38.6	432	-	3,10	
Anti-Parkinson agents	1,692	27.4	785	-	2,599	
General anesthetics				-		

Table 9 (continued)
Nonmedical use of pharmaceuticals: 2005

	Estimated	Relative	95% Confidence interval			
Selected drug categories and selected drugs ¹	Selected drug categories and selected drugs ¹ visits ^{2,3,4}		Lower bound	-	Upper bound	
Muscle relaxants	33,695	14.3	24,235	-	43,154	
Carisoprodol	20,082	18.5	12,791	-	27,374	
Cyclobenzaprine	7,629	16.0	5,230	-	10,028	
Miscellaneous CNS agents	900	32.0	336	-	1,464	
RESPIRATORY AGENTS	28,017	12.1	21,383	-	34,651	
Antihistamines	4,429	14.1	3,206	-	5,652	
Bronchodilators	3,043	27.1	1,427	-	4,659	
Decongestants	1,309	39.0	308	-	2,310	
Expectorants	1,960	25.7	971	-	2,949	
Upper respiratory combinations	15,837	14.9	11,209	-	20,465	
Respiratory agents NTA	3,692	14.4	2,651	-	4,733	
CARDIOVASCULAR AGENTS	37,095	9.5	30,224	-	43,966	
Antiadrenergic agents, centrally acting	5,125	19.9	3,124	-	7,127	
Beta-adrenergic blocking agents	9,824	13.3	7,260	-	12,388	
Calcium channel blocking agents	5,434	16.8	3,642	-	7,226	
Diuretics	5,332	15.2	3,739	-	6,924	
Cardiovascular agents NTA	18,881	12.0	14,443	-	23,320	

¹ The classification of drugs used in DAWN is derived from the Multum *Lexicon*, © 2007, Multum Information Services, Inc. The classification was modified to meet DAWN's unique requirements (2007). The Multum Licensing Agreement governing use of the *Lexicon* can be found on the Internet at http://www.multum.com.

² These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

³ ED visits often involve multiple drugs. Such visits will appear multiple times in this table (e.g., a visit involving oxycodone and hydrocodone will appear twice in this table). Summing ED visits as reported will produce incorrect and inflated counts of ED visits.

⁴ Three dots (...) indicate that an estimate with an RSE greater than 50% or an estimate less than 30 has been suppressed. **SOURCE**: Office of Applied Studies, SAMHSA, Drug Abuse Warning Network, 2005 (03/2008 update).

Table 10 Nonmedical use of pharmaceuticals, by patient and visit characteristics: 2005

Patient characteristics	Estimated visits ^{1,2}	Visit characteristics	Estimated visits ^{1,2}
ED visits, nonmedical use	669,214		•
Gender		Number of drugs involved	
Male	308,072	Single drug	305,688
Female	360,975	Multiple drugs	363,526
Unknown	167		
Age		Disposition	
0-5 years	8,840	Treated and released	381,881
6-11 years	4,753	Discharged home	343,810
12-17 years	62,174	Released to police/jail	14,097
18-20 years	46,674	Referred to detox/treatment	23,973
21-24 years	65,239	Admitted to this hospital	196,314
25-29 years	66,936	ICU/critical care	58,733
30-34 years	58,483	Surgery	371
35-44 years	144,758	Chemical dependency/detox	
45-54 years	110,565	Psychiatric unit	33,909
55-64 years	50,646	Other inpatient unit	101,280
65 years and older	49,443	Other disposition	91,018
Unknown		Transferred	54,384
Race/ethnicity		Left against medical advice	11,479
White	442,305	Died	1,827
Black	73,380	Other	
Hispanic	38,173	Not documented	6,960
Race/ethnicity not tabulated above (NTA)	7,044		
Unknown	108,312		

¹ These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

² Three dots (...) indicate that an estimate with an RSE greater than 50% or an estimate less than 30 has been suppressed. **SOURCE**: Office of Applied Studies, SAMHSA, Drug Abuse Warning Network, 2005 (03/2008 update).

Table 16 Suicide attempts: 2005

	Estimated	Relative	95% Confidence interva			
Drug category and selected drugs ¹	visits ^{2,3,4}	standard error (RSE)	Lower bound	-	Upper bound	
Total drug-related ED visits, suicide attempts	151,568	8.0	127,715	-	175,421	
Major su	bstances of abuse					
Alcohol	47,891	8.4	39,981	-	55,800	
Alcohol in combination	46,806	8.3	39,204	-	54,408	
Alcohol alone	1,085	35.4	332	-	1,838	
Non-alcohol illicits	33,784	17.8	21,979	-	45,588	
Cocaine	19,628	20.4	11,765	-	27,492	
Heroin	3,167	37.0	870	-	5,463	
Marijuana	11,955	17.6	7,842	-	16,068	
Stimulants	5,410	31.9	2,031	-	8,789	
Amphetamines	1,646	28.2	735	-	2,557	
Methamphetamine	3,853	36.6	1,092	-	6,614	
MDMA (Ecstasy)	529	36.7	148	-	909	
GHB				-		
Flunitrazepam (Rohypnol)				-		
Ketamine				-		
LSD				-		
PCP		68.1		-		
Miscellaneous hallucinogens				-		
Inhalants	794	40.0	171	-	1,416	
Combinations not tabulated above (NTA)				-		
Othe	er substances					
PSYCHOTHERAPEUTIC AGENTS	82,144	7.4	70,250	-	94,038	
Antidepressants	27,086	7.7	22,976	-	31,196	
MAO inhibitors				-		
SSRI antidepressants	13,377	9.9	10,779	-	15,97	
Tricyclic antidepressants	3,008	18.0	1,950	-	4,06	
Miscellaneous antidepressants	2,681	22.1	1,520	-	3,84	
Antipsychotics	17,129	11.6	13,235	-	21,02	
Anxiolytics, sedatives, and hypnotics	52,022	8.8	43,090	-	60,95	
Barbiturates	1,219	23.7	654	-	1,78	
Benzodiazepines	35,676	10.3	28,439	-	42,91	
Alprazolam	14,530	16.2	9,915	-	19,14	
Clonazepam	9,064	12.8	6,789	_	11,33	

Table 16 (continued)
Suicide attempts: 2005

	Estimated	Relative	95% Confidence interval			
Drug category and selected drugs ¹	category and selected drugs ¹ visits ^{2,3,4} visits ^{2,3,4} error (RSE)		Lower bound	-	Upper bound	
Diazepam	3,968	16.4	2,694	-	5,242	
Lorazepam	5,182	18.6	3,296	-	7,067	
Benzodiazepines NOS	3,343	33.0	1,179	-	5,508	
Misc. anxiolytics, sedatives, and hypnotics	17,522	9.8	14,161	-	20,883	
Diphenhydramine	6,583	18.5	4,200	-	8,967	
Hydroxyzine	1,795	24.2	944	-	2,647	
Zolpidem	4,972	14.4	3,572	-	6,373	
Anxiolytics, sedatives, and hypnotics NOS	2,147	27.8	975	-	3,319	
CNS stimulants	1,782	17.8	1,160	-	2,404	
Amphetamine-dextroamphetamine				-		
Caffeine	450	42.4	76	-	823	
Dextroamphetamine				-		
Methylphenidate	818	33.5	280	-	1,356	
CENTRAL NERVOUS SYSTEM AGENTS	66,321	7.4	56,726	-	75,916	
Analgesics	54,858	7.1	47,250	-	62,466	
Antimigraine agents				-		
Cox-2 inhibitors	514	48.2	28	-	999	
Opiates/opioids	20,359	9.0	16,770	-	23,948	
Opiates/opioids, unspecified	2,819	26.7	1,343	-	4,296	
Narcotic analgesics	17,801	10.6	14,117	-	21,485	
Buprenorphine/combinations				-		
Codeine/combinations	2,656	28.0	1,199	-	4,113	
Fentanyl/combinations				-		
Hydrocodone/combinations	7,035	15.1	4,955	-	9,115	
Hydromorphone/combinations				-		
Meperidine/combinations				_	•••	
Methadone	1,596	33.2	557	_	2,636	
Morphine/combinations	1,210	27.3	564	_	1,856	
Oxycodone/combinations	4,229	15.6	2,933	_	5,524	
Propoxyphene/combinations	2,129	24.7	1,099	_	3,159	
Nonsteroidal anti-inflammatory agents	14,117	9.2	11,572	_	16,662	
Ibuprofen	10,917	10.0	8,771	-	13,064	
Naproxen	3,224	18.7	2,042	_	4,406	
Salicylates/combinations	4,645	16.2	3,173	_	6,118	

Table 16 (continued) Suicide attempts: 2005

	Estimated	Relative	95% Confidence interval			
Drug category and selected drugs ¹	Estimated visits ^{2,3,4}	standard error (RSE)	Lower bound	-	Upper bound	
Miscellaneous analgesics/combinations	22,692	9.3	18,576	-	26,808	
Acetaminophen/combinations	21,017	9.3	17,167	-	24,866	
Tramadol/combinations	1,515	29.8	631	-	2,399	
Tramadol	1,079	37.9	278	-	1,879	
Acetaminophen-tramadol				-		
Analgesic combinations NTA				-		
Anorexiants				-		
Anticonvulsants	9,389	16.8	6,299	-	12,480	
Antiemetic/antivertigo agents				-		
Anti-Parkinson agents	543	46.2	51	-	1,035	
General anesthetics				-		
Muscle relaxants	5,785	14.8	4,112	-	7,458	
Carisoprodol	2,038	21.1	1,197	-	2,879	
Cyclobenzaprine	2,784	22.7	1,545	-	4,022	
Miscellaneous CNS agents				-		
RESPIRATORY AGENTS	7,662	15.4	5,345	-	9,979	
Antihistamines	1,650	20.7	980	-	2,320	
Bronchodilators				-		
Decongestants		58.6		-		
Expectorants	474	42.6	79	-	869	
Upper respiratory combinations	4,207	22.4	2,359	-	6,056	
Respiratory agents NTA	1,244	35.1	388	-	2,100	
CARDIOVASCULAR AGENTS	5,814	17.8	3,788	-	7,840	
Antiadrenergic agents, centrally acting	912	28.2	408	-	1,416	
Beta-adrenergic blocking agents	1,916	27.4	885	-	2,947	
Calcium channel blocking agents	193	38.9	46	-	341	
Diuretics	539	41.4	102	-	976	
Cardiovascular agents NTA	3,024	18.6	1,922	-	4,126	

¹ The classification of drugs used in DAWN is derived from the Multum *Lexicon*, © 2007, Multum Information Services, Inc. The classification was modified to meet DAWN's unique requirements (2007). The Multum Licensing Agreement governing use of the *Lexicon* can be found on the Internet at http://www.multum.com.

² These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

³ Three dots (...) indicate that an estimate with an RSE greater than 50% or an estimate less than 30 has been suppressed.

⁴ This column denotes statistically significant (*p* < 0.05) increases or decreases between estimates for the periods shown. **SOURCE**: Office of Applied Studies, SAMHSA, Drug Abuse Warning Network, 2005 (03/2008 update).

Table 17
Suicide attempts, by patient and visit characteristics: 2005

Patient characteristics	Estimated visits ^{1,2}	Visit characteristics	Estimated visits ^{1,2}
Total drug-related ED visits, suicide attempts	151,568		
Gender		Number of drugs involved	
Male	58,775	Single drug	58,541
Female	92,682	Multiple drugs	93,027
Unknown			
Age		Disposition	
0-5 years		Treated and released	28,662
6-11 years		Discharged home	21,950
12-17 years	17,869	Released to police/jail	3,377
18-20 years	13,763	Referred to detox/treatment	3,335
21-24 years	14,989	Admitted to this hospital	74,595
25-29 years	18,761	ICU/critical care	28,297
30-34 years	14,074	Surgery	
35-44 years	39,140	Chemical dependency/detox	105
45-54 years	22,057	Psychiatric unit	24,395
55-64 years	6,745	Other inpatient unit	21,691
65 years and older	4,079	Other disposition	48,311
Unknown		Transferred	41,846
Race/ethnicity		Left against medical advice	
White	89,172	Died	
Black	26,229	Other	
Hispanic	13,353	Not documented	1,842
Race/ethnicity not tabulated above (NTA)	2,132		
Unknown	20,682		

These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

 $^{^2 \ \ \, \}text{Three dots (...) indicate that an estimate with an RSE greater than 50\% or an estimate less than 30 has been suppressed.}$

Table 19 Seeking detox: 2005

	Fatimated	Relative	95% Confidence interval			
Drug category and selected drugs ¹	Estimated visits ^{2,3,4}	standard error (RSE)	Lower bound	-	Upper bound	
Total drug-related ED visits, seeking detox	126,226	12.0	96,495	-	155,957	
Major	substance of abuse					
Alcohol	47,494	14.0	34,502	-	60,485	
Alcohol in combination	47,154	14.0	34,194	-	60,114	
Alcohol alone				-		
Non-alcohol illicits	101,244	13.1	75,203	-	127,286	
Cocaine	56,061	11.3	43,647	-	68,475	
Heroin	40,895	14.3	29,451	-	52,340	
Marijuana	22,486	25.0	11,462	-	33,511	
Stimulants	15,402	47.2	1,149	-	29,656	
Amphetamines		63.1		-		
Methamphetamine		50.7		-		
MDMA (Ecstasy)	511	34.1	170	-	853	
GHB				-		
Flunitrazepam (Rohypnol)				-		
Ketamine				-		
LSD				-		
PCP	729	45.1	85	-	1,373	
Miscellaneous hallucinogens				-		
Inhalants				-		
Combinations not tabulated above (NTA)	191	40.0	41	-	340	

¹ The classification of drugs used in DAWN is derived from the Multum *Lexicon*, © 2007, Multum Information Services, Inc. The classification was modified to meet DAWN's unique requirements (2007). The Multum Licensing Agreement governing use of the *Lexicon* can be found on the Internet at http://www.multum.com.

² These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

³ Estimates are all expressed in visits. Visits cannot be summed across drugs because drug-related visits often involve multiple drugs.

⁴ Three dots (...) indicate that an estimate with an RSE greater than 50% or an estimate less than 30 has been suppressed.

Table 20 Seeking detox, by patient and visit characteristics: 2005

Patient characteristics	Estimated visits ^{1,2}	Visit characteristics	Estimated visits ^{1,2}
Total drug-related ED visits, seeking detox	126,226		
Gender		Number of drugs involved	
Male	82,539	Single drug	46,006
Female	43,643	Multiple drugs	80,220
Unknown			
Age		Disposition	
0-5 years		Treated and released	64,598
6-11 years		Discharged home	28,247
12-17 years	1,776	Released to police/jail	907
18-20 years	9,500	Referred to detox/treatment	35,444
21-24 years	16,073	Admitted to this hospital	42,582
25-29 years	17,345	ICU/critical care	185
30-34 years	16,421	Surgery	
35-44 years	38,291	Chemical dependency/detox	27,664
45-54 years	22,584	Psychiatric unit	7,877
55-64 years	2,989	Other inpatient unit	6,852
65 years and older	1,176	Other disposition	19,046
Unknown		Transferred	12,752
Race/ethnicity		Left against medical advice	2,364
White	71,147	Died	
Black	27,105	Other	2,328
Hispanic	9,399	Not documented	1,602
Race/ethnicity not tabulated above (NTA)	790		
Unknown	17,786		

These are estimates of ED visits based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

 $^{^2 \ \ \, \}text{Three dots (...) indicate that an estimate with an RSE greater than 50\% or an estimate less than 30 has been suppressed.}$

Table C2
Drug-related ED visits, by type of case: 2005

	Unweighted Weighted Relative			95% Confi	onfidence interval			
Type of case	sample data	estimates ¹	standard error (RSE)	Lower bound	-	Upper bound		
Drug-related ED visits								
Suicide attempt	11,523	151,568	8.0	127,715	-	175,421		
Seeking detox	22,166	126,226	12.0	96,495	-	155,957		
Alcohol only (age < 21)	9,636	109,174	11.2	85,281	-	133,068		
Adverse reaction	88,278	1,250,377	11.5	969,563	-	1,531,191		
Overmedication	23,279	290,627	7.6	247,216	-	334,039		
Malicious poisoning	697	6,527	16.7	4,392	-	8,662		
Accidental ingestion	3,866	56,517	9.2	46,285	-	66,750		
Other	108,683	1,018,009	13.4	750,423	-	1,285,594		
Total drug-related ED visits	268,128	3,009,025	7.6	2,560,019	-	3,458,031		
		Drugs ²						
Suicide attempt	24,391	318,526	7.5	271,423	-	365,629		
Seeking detox	45,057	253,213	13.8	184,611	-	321,815		
Alcohol only (age < 21)	9,636	109,174	11.2	85,281	-	133,068		
Adverse reaction	111,218	1,597,947	12.3	1,214,031	-	1,981,863		
Overmedication	40,048	504,041	8.3	422,223	-	585,860		
Malicious poisoning	1,239	12,998	17.6	8,502	-	17,494		
Accidental ingestion	4,997	77,042	10.1	61,762	-	92,322		
Other	183,834	1,751,514	12.4	1,326,667	-	2,176,360		
Drugs in all drug-related ED visits	420,420	4,624,455	7.7	3,922,142	-	5,326,768		

¹ These are estimates based on a representative sample of non-Federal, short-stay hospitals with 24-hour EDs in the United States.

² These are estimates of drugs. A single ED visit may involve multiple drugs.

Table C5
Population by age and gender: 2005¹

Gender and age	Total United States	Males	Females
Total	296,410,404	145,999,746	150,410,658
0-5 years	24,211,671	12,381,683	11,829,988
6-11 years	23,767,612	12,160,248	11,607,364
12-17 years	25,490,701	13,059,495	12,431,206
18-20 years	12,450,315	6,400,184	6,050,131
21-24 years	16,856,810	8,702,127	8,154,683
25-29 years	20,065,702	10,268,169	9,797,533
30-34 years	20,077,210	10,153,091	9,924,119
35-44 years	43,862,464	21,940,039	21,922,425
45-54 years	42,482,265	20,895,355	21,586,910
55-65 years	30,355,541	14,626,718	15,728,823
65 years and older	36,790,113	15,412,637	21,377,476

Population estimates for 2005, as of July 2006, from U.S. Census Bureau County Population Dataset NST-EST2005-ALLDATA (see http://www.census.gov/popest/national/files/NST_EST2005_ALLDATA.csv).