Image Based Biomarkers of Breast Cancer Risk: Analysis of Risk Disparity Among Minority Populations

Fengshan Liu¹, Predrag R. Bakic², David D. Pokrajac¹, Charlie Wilson¹, Xiguan Shi¹, Despina Kontos², Andrew D.A. Maidment²

¹ Applied Mathematics Research Center, Delaware State University, Dover, DE 19901 ² Department of Radiology, University of Pennsylvania, Philadelphia, PA 19104

{FLiu,DPokrajac,CWilson,XShi}@desu.edu,{Predraq.Bakic,Andrew.Maidment}@uphs.upenn.edu



ABSTRACT

Delaware State

University

Advances in understanding breast cancer risk factors, coupled with advances in screening methodologies and prevention strategies, are critical to the development of nersonalized breast cancer surveillance and prevention. This HRCLI/MI Partnership Training Award has built upon a collaboration between DSU and Penn, with the objectives to: (1) extend the skills of DSU faculty to become competitive breast cancer researchers (2) establish an independent research program at DSU by performing a joint research project focused on breast cancer risk disparity in minority populations; and (3) produce high quality publications and funded grants at DSU to sustain breast cancer research.

DSU Faculty Training Related to Breast Cancer Research

The training program developed for DSU, which combines didactic instruction covering breast cancer fundamentals with ongoing research training, is designed to inculcated the DSU faculty with the culture and language of breast cancer research.

Each DSU faculty member attended up to 3 courses at Penn to augment their knowledge of breast cancer. The courses also provided exemplary didactic methods, so that DSU could create new curricula for DSU students relevant to breast cancer. Courses included GCB535: Intro to Bioinformatics

CAMB512 Cancer Biology and Genetics BE545/CIS537 Biomedical Image Analysis

GCB / CAMB 752: Seminar in Genomics

BE483 Molecular Imaging
EP801 Fundamentals of Epidemiologic Study Designs.

In addition, a seminar course on the Basics of Breast Cancer was taught by Penn faculty at DSU, covering epidemiology, anatomy, pathology, imaging, and clinical oncology

The research training component of the grant is focused on the assessment of the racial dependence of image-based risk biomarkers (IBRB) of breast cancer. Our long-term goal is to develop an accurate method for breast cancer risk estimation in minority populations by including IBRB and genetic risk factors. Despite evidence of racial disparities in breast cancer statistics, reported results to date have not established a racial dependence on IBRB.

The first aim of our project is to assess IBRB racial dependence in a large clinical trial using images from the ACRIN DMIST study.[1] During preliminary work, a large number of women were identified for whom multiple images were taken to capture the whole breast. As a result, the IBRB cannot be estimated for these women using existing methods. Thus, we have additionally undertaken efforts to develop a method to combine IBRB data from multiple images. To validate these methods, we have used synthetic mammograms generated from software breast phantoms.[2]



The flow-chart of research steps aimed

Preparing Clinical Data from ACRIN DMIST Study

After IRB approval, we created a database of 11.106 (5.553 minority and corresponding Caucasian controls) cancer-free women gathered as part of the ACRIN DMIST study. The anonymized database consisted of both study images and selected study forms. The images have been stored in a CTP/MIRC database, facilitating queries and convenient interface with other processing steps. Scripts have been developed for querying MIRC XML files to perform quality control and to prepare the data for IBRB

Previously, we demonstrated that in a database of 657 women with mammography exams, 85 (i.e., 13%) cases has only partial breast visualization in a single image due to breast size. To date, we have observed a similar percentage in an initial assessment of 990 ACRIN cases. This high prevalence has required that we develop a strategy to measure IBRB in breasts spanning multiple images.










Simulation of Large Breasts



Sections (left) and a synthetic mammogram (right) of a sample software breast phanton

Jointly we have added two specific features to simulate large breasts, including simulating the partial volume of tissues in a voxel and improving the efficiency of the software through parallelization of the phantom generation procedure. The method developed is octree-based. This algorithm has been shown to almost achieve the theoretical lower bound in computational complexity of such simulations.

We are currently developing a method to combine images with partial breast visualization, to be validated using synthetic mammograms of an anthropomorphic software breast phantom developed at Penn. The simulation of large breasts required that we improve the software phantom design in terms of simulation time, memory requirements, and scalability.[2]



Estimate Image-Based Risk Biomarkers

We have been developing an interface between the MIRC study database and IBRB image analysis pipeline, which are physically located in two different labs at Penn. The proposed interface will allow remote access to both the MIRC and pipeline servers. The analysis protocol is designed so that any incoming image is automatically analyzed. From the MIRC server it will be possible to start an automatic sequence pushing images to the pipeline. After each image has been processed, the resulting breast density data and texture images are automatically pushed to MIRC and stored as associated record series. This organization would allow running additional image analyses remotely from DSU, after the migration of the study database from Penn to DSU.



The flow-chart of the breast image analysis pipel

The IBRB image analysis pipeline employs a fully-automated breast percent density (PD%) estimation technique based on an adaptive multi-class fuzzy-c-means algorithm. The segmented dense tissue regions are also used to create a regional tissue mask from which the imaging parenchymal descriptors are extracted. Parenchymal texture analysis is performed by extracting multiple descriptors, offering a range of optimization and image processing parameters, as well as the first-order statistics of the original breast image and the stochastic fractal dimension image, and second-order statistics extracted from the co-occurrence and run-length matrices. Statistical information for any desired feature (e.g., feature merging approach combining principal component analysis (PCA) with independent component







Perform Statistical Analysis of Risk Biomarkers

The extracted breast density and parenchymal texture descriptors will be used to test racial dependence of image-based biomarkers of breast cancer risk. We plan to publish our results and compare them with reports in the literature (e.g. [4]). In addition, the results of the pipeline analysis of ACRIN data will be used in our long term research goal of developing a an accurate method for breast cancer risk estimation in minority populations by including IBRB and genetic risk factors.

Conclusion and Impact

Our ultimate hope is that this DoD BCRP HBCU/MI award will provide an excellent environment at DSU for their faculty to conduct state of the art breast cancer research. Our success is evident by the expanding number of faculty and staff involved in breast cancer research at DSU. and the high-quality of the presentations and publications resulting from this proje Moreover, DSU is becoming an integral ons resulting from this project. part of a network of established breast cancer investigators at Penn and beyond. For the rest of the award time we will focus preparing strong grant applications to



List of journal papers, conference and invited presentations, and patent applications:

- Lef journal pagests, conference and invited presentations, and patent applications:
 Puriage, D. Maderica, A.D., Balloc P.P. ("pinited Generation of High Peculition Times Anthropomorphic Software Phartoms," submitted for patients in Medical Physics, 2011.
 N. S., Lei, F. Maderica, A.D., Balloc, P.P. ("Arrimentes to the Deviamation Model of An Arthropomorphic Computer Generation of High Peculition Times and Peculitics ("Arriments to the Polysics, 2011.
 No. 3, 1989, 1

- Pokraiac, D.D., Bakic, P.R., Maidment, A.D.A., "Techniques for Simulation of Multi-dimensional Spatial Regions. ation, University of Delaware/Delaware State University, (DSU No. 12-01), 2011

References

- Plazo ED, Galtorios C, Hendrick E, et al. (2003) Diagnosis performance of digital versus film mammography for breast-forcers reserving. N Eur J J Mer 533 (T), 1773-83.

 N Evaluage C), Malester, B, March P, (2011) A mode of the Chipfer Meridine or share in heast phartons. Revising C) March P, Rose P, (2011) A mode of the Chipfer Meridine year because C meridine J Meridine S (Ander E), Wang Y et al. (2011) A Fully A-fundamed Software Psychie for Pizerschyntal Pattern Assights in March P, (2011) A Fully A-fundamed Software Psychie for Pizerschyntal Pattern Assights in March P, (2011) A Fully A-fundamed Software Psychie for Pizerschyntal Pattern Assights in March P, (2011) A Fully A-fundamed Software Psychie for Pizerschyntal Pattern Assights in March P, (2011) A Fully A-fundamed Software P, (2011) A Fully A-fundamed Softw

Acknowledgements

This project is funded by the US Department of Defense BCRP HBCUMN Partmership Training Award (#BC083639). The authors are grateful of Ms. Fatime Boulant, Mr. Felyu Chen, Mr. Joseph Chui, Dr. Brad Keller, Ms. Talyana Nu