


Priorities for Effective Management of Coral Diseases


Andrew W. Bruckner



Black-band disease, *Stephanocoenia intersepta*



White-band disease, *Acropora palmata*



White plague, *Montastraea annularis*



U.S. Department of Commerce
National Oceanic and Atmospheric Administration
National Marine Fisheries Service

NOAA Technical Memorandum NMFS-OPR-22
August 2002

Priorities for Effective Management of Coral Diseases

Andrew W. Bruckner

Office of Protected Resources, Rm 13604
NOAA National Marine Fisheries Service
1315 East-West Highway
Silver Spring, MD 20910-3282

**NOAA Technical Memorandum
NMFS-OPR-22
August 2002**



U.S. Department of Commerce
Donald L. Evans, Secretary

National Oceanic and Atmospheric Administration
Vice Admiral Conrad C. Lautenbacher, Jr., USN (Ret.)
Under Secretary for Oceans and Atmosphere

National Marine Fisheries Service
William T. Hogarth, Assistant Administrator for Fisheries

Table of Contents

EXECUTIVE SUMMARY	1
INTRODUCTION	2
INFORMATION NEEDS FOR EFFECTIVE MANAGEMENT OF CORAL DISEASES	7
Early warning systems	7
The spatial distribution and temporal variations of coral disease outbreaks	7
Relationships with environmental factors, anthropogenic stresses and natural disturbances ..	10
Standardized nomenclature to describe, identify and differentiate diseases	13
Factors that facilitate the occurrence, spread and transmission of pathogens	14
Effects on population dynamics, community structure and ecosystem function	16
Measures to mitigate disease impacts	19
PROPOSED STRATEGIES TO ADDRESS MANAGEMENT NEEDS FOR CORAL DISEASES	20
Early warning systems	20
The spatial distribution and temporal variations of coral disease outbreaks	21
Relationships with environmental factors, anthropogenic stresses and natural disturbances	23
Standardized nomenclature to describe, identify and differentiate diseases	24
Factors that facilitate the occurrence, spread and transmission of pathogens	26
Effects on population dynamics, community structure and ecosystem function	26
Measures to mitigate disease impacts	27
CONCLUSION	27
ACKNOWLEDGMENTS	29
APPENDIX I: Monitoring and Assessment Approaches	30
Atlantic and Gulf Rapid Reef Assessment (AGRRA)	30
Caribbean Coastal and Marine Productivity Network (CARICOMP)	33
EPA Florida Keys	37
APPENDIX II. Priorities for research on coral diseases linked to management objectives in Florida.	40
CORAL DISEASE LITERATURE	41

Priorities for Effective Management of Coral Diseases

EXECUTIVE SUMMARY

Diseases and syndromes of scleractinian corals and associated species have increased in number in recent years, and they are now recognized as important phenomena capable of altering the structure and composition of coral reefs. Since the early 1990s there has been a concerted effort to characterize coral diseases, including the application of novel molecular tools to confirm identities of pathogens and understand mechanisms of host response and resistance. Most of the causative agents of emerging diseases, factors contributing to their occurrence and spread, and consequences on coral populations remain incompletely understood, however. A long-term, multidisciplinary research and monitoring program for coral diseases is necessary to assist resource managers in identifying and responding to emerging coral diseases. These efforts should involve management-driven strategies that include 1) an early warning system to predict and identify disease outbreaks; 2) documentation of spatial distribution and temporal variations of coral diseases at local to global scales; 3) elucidation of relationships of environmental stressors, localized anthropogenic impacts, and widespread phenomena such as global warming and El Niño on coral health, disease, degradation and recovery; 4) development of standardized terminology for diseases through characterization of the visual appearance, pathology and etiology; 5) development of molecular probes and other tools to identify and verify microbial pathogens in the field; 6) identification of factors that facilitate the introduction, spread and transmission of pathogens; 7) long-term monitoring and targeted research on the effects of disease on coral species and populations, associated species, and ecosystem structure and function; and 8) implementation of measures to mitigate disease impacts, including strategies that reduce anthropogenic stressors responsible for the proliferation or spread of diseases and the development of novel techniques to treat affected corals and improve habitat quality. Progress in coral disease research has been hindered by a lack of coordinated research, practical difficulties in correctly identifying diseases and characterizing their etiology and epizootiology, limited availability of funding, and limited awareness and recognition by policy makers and the public of the importance of coral diseases and their potential global implications. Given existing limitations in funding, disease research should be focused towards those efforts that have the greatest management benefits with applications on local to regional scales.

INTRODUCTION

Coral reef ecosystems are being degraded at an accelerated rate. The underlying causes of reef decline are diverse, and include pollution, sedimentation, fishing impacts, habitat destruction, invasive species, bleaching, disease, global climate change, and other factors. In recent years, disease outbreaks have caused widespread mortalities to scleractinian corals, gorgonians, sea urchins, reef fishes, sponges, algae, and other coral reef organisms (Peters, 1997; Harvell et al., 1999; Williams and Bunkley-Williams, 2000). Characterization of the cause, prevalence, and consequences of most of these events is limited or non-existent. For instance, hundreds of studies have been published on coral diseases (Appendix III) since the first reported coral disease (Antonius, 1973), and yet the causative agent has been confirmed for only four diseases (Richardson, 1998; Patterson et al., 2002). In addition, very limited quantitative data are available on the spatial and temporal distribution of coral diseases, synergistic and cumulative effects of environmental and anthropogenic stressors, and long-term consequences on the composition, structure and function of reef communities (Green and Bruckner, 2000).

Although quantitative field observations on coral diseases are restricted to a few locations and relatively small numbers of corals, there is a growing consensus that coral diseases have proliferated in the 1990s (Santavy and Peters, 1997; Goreau et al., 1998; Hayes and Goreau, 1998; Richardson, 1998; Harvell et al., 1999; Williams and Bunkley-Williams, 2000). In an examination of all information published on coral diseases between 1973-1999, diseases were reported in over 102 coral species from 54 countries, with a disproportionate number of records (66%) from the Wider Caribbean (Green and Bruckner, 2000). Currently, the majority of these reports are for black-band disease (BBD), white-band disease (WBD) and white plague. These diseases were first documented from the wider Caribbean in the 1970s, but they were reported at a low prevalence or from few locations (Antonius, 1977; Garrett and Ducklow, 1975; Dustan, 1977). Outbreaks of BBD and WBD were reported from numerous locations in the Caribbean in the 1980s and 1990s, and both are now circumtropical in distribution (Antonius, 1987; Green and Bruckner, 2000). White plague was first identified as a significant infectious disease in the Florida Keys in 1975, and it was still common in 1984 on 19 different reefs in the Key Largo region (Dustan, 1977; 1993). A more virulent form of white plague (plague type II) emerged in 1995 on reefs in the northern Florida Keys (Richardson et al., 1998a,b). Additional outbreaks of white plague (plague type II and type III) were also observed in Florida and other locations around the Caribbean between 1999-2001 (Weil, 2001; Nugues, 2002; Miller et al., in press; Richardson and Aronson, in press; Bruckner, unpubl. Obs). To date, 27 other diseases of scleractinian corals and gorgonians were reported from the western Atlantic (Table 1) and 16 from the IndoPacific (Table 2). Western Atlantic reefs have

Table 1. Diseases, syndromes, abnormal tissue conditions, and parasitic infestations of scleractinian corals and gorgonians on coral reefs in the tropical western Atlantic. Data are compiled from publications and web sites on coral disease identified through 2002. Other names are used in the literature to describe similar disease signs but these have been combined under the following diseases: (1) BBD is also called black line disease; (2) WBD and white plague are also called white line disease; (3) WBD is also used to describe massive corals in the Caribbean with signs similar to white plague; (4) YBD is also called yellow-band disease and yellow pox disease; (5) rapid wasting disease was renamed rapid wasting syndrome, and recently reported to be synonymous with parrotfish white spot biting; and (6) ridge mortality has been called damselfish-ridge denuding syndrome.

Condition	First observed	Known Range	Affected species	Source
Black-band disease (BBD) ¹	Belize; Florida; Puerto Rico	W. Atlantic 20 countries	faviids, agaricids gorgonians 22 species total	Antonius, 1972
White-band disease (WBD) ²	St. Croix	Aruba, Bahamas, Belize, Bonaire, British Virgin Islands (BVI), Cuba, Curacao, Florida, Guadalupe, Jamaica, Puerto Rico, Tobago, Nicaragua, St. Lucia, US Virgin Islands (USVI)	<i>A. palmata</i> <i>A. cervicornis</i> <i>A. prolifera</i>	Gladfelter, 1977
WBD type II	Bahamas	Bahamas	<i>A. cervicornis</i>	Richie and Smith, 1995
Shut-down reaction	Florida, Belize	Belize, Florida	massive corals, acroporids	Antonius, 1977
White plague ³	Florida	Bahamas, Bermuda, Bonaire, BVI, Curacao, Florida, Guadalupe, Jamaica, Nicaragua, Puerto Rico, Tobago, St. Lucia, USVI	faviids, agaricids, mussids, poritids; 12 species	Dustan, 1977
Plague type II	Florida	Bonaire, Curacao, Dominica, Florida, Mexico, Puerto Rico, St. Lucia, USVI, Venezuela	33 species	Zorpette, 1995
Plague type III	Florida	Caribbean	large massive corals (<i>M. faveolata</i> , <i>C. natans</i>)	Richardson and Aronson, 2001
Yellow-blotch disease (YBD) ⁴	Florida	Bahamas, Bonaire, Curacao, Dominica, Florida, Honduras, Jamaica, Mexico, Panama, Puerto Rico, St. Lucia, Venezuela	<i>Montastraea annularis</i> complex; <i>C. natans</i>	Reeves, 1994
White pox	Florida	Bahamas, Florida	<i>A. palmata</i>	Porter, 1996
Patchy necrosis	Puerto Rico	Caribbean	<i>A. palmata</i>	Bruckner and Bruckner, 1997
Rapid wasting disease ⁵	Bonaire	Caribbean	<i>Montastraea</i> spp., <i>C. natans</i>	Cervino et al., 1997
White spot syndrome	Caribbean	?	massive corals	Global Coral Reef Alliance web pages (Cervino, pers. comm)
Ridge mortality ⁶	Flower Gardens, Gulf of Mexico	Caribbean	<i>C. natans</i> , <i>D. strigosa</i>	Abbott, 1979
Dark-spots disease (DSD)	Columbia	Caribbean	<i>M. Annularis</i> , <i>S. siderea</i> , <i>S. radians</i> , <i>S. intersepta</i>	Gil-Agudelo and Garzón-Ferreira, 2001

Table.1 continued. Diseases, syndromes, abnormal tissue conditions, and parasitic infestations of scleractinian corals and gorgonians in the tropical western Atlantic continued. ** Mortalities of sea fans were observed in the 1980s, but they may be caused by disease or other factors.

DSD- II	Bermuda, Bonaire, Colombia, Puerto Rico and Venezuela	Caribbean	<i>S. intersepta</i> ; <i>M. annularis</i> ; <i>M. favedata</i> ; <i>M. cavernosa</i> ; <i>C. natans</i> ; <i>C. amaranthus</i> ; <i>S. siderea</i>	Weil, 2001
Red-band disease (RBD) type I	Belize, Jamaica, Puerto Rico	Belize, Curaçao, Dominica, Jamaica, Puerto Rico	<i>Gorgonia</i> , <i>Agaricia</i> , <i>Colpophyllia</i> , <i>Mycetophyllia</i> and <i>Stephanocoenia</i>	Rützler et al., 1983 Santavy and Peters, 1997
RBD type II	Bahamas	Bahamas	<i>D. stigosa</i> , <i>M. annularis</i> , <i>M. cavernosa</i> , <i>P. astreoides</i> and <i>S. radians</i>	Richardson, 1992
Hyperplasia	Puerto Rico, US Virgin Islands, Jamaica, Netherlands Antilles, Trinidad, Belize	Caribbean	massive corals	Loya et al., 1984
Neoplasia	Florida	Florida, Netherlands Antilles, Puerto Rico, Trinidad	<i>A. palmata</i>	Peters et al., 1986
Coccidiosis	Jamaica, Puerto Rico, USVI		<i>A. agaricites</i> , <i>D. cylindicus</i> , <i>D. strigosa</i> , <i>M. meandrites</i> , <i>M. cavernosa</i> , <i>P. astreoides</i> , <i>P. porites</i> 8 species	Upton and Peters, 1986
Nematopsis spores	USVI		<i>Porites</i> spp.	Peters, 1984
Stress-related necrosis	Puerto Rico			Peters, 1984
Ring disease	Bermuda	Florida, Honduras	<i>D. labyrinthiformis</i>	Weil, 2001
Finger coral denuding syndrome	?		<i>Porites porites</i>	Williams and Bunkley-Williams, 2000
Star coral polyp necrosis	?		<i>M. cavernosa</i>	Williams and Bunkley-Williams, 2000
Algal tumors	Bonaire, Trinidad	Caribbean	<i>Gorgonia</i> spp. <i>Pseudoplexaura</i> ; <i>Plexaura</i>	Morse et al., 1977
Aspergillosis	Columbia, Costa Rica, Panama, Trinidad**	Caribbean	<i>Gorgonia</i> spp.	Nagelkerken et al., 1997
Fire coral fungal disease	Florida	Florida	<i>Millepora</i> spp.	TeStrake et al., 1988

Table 2. Diseases, syndromes, and anomalies reported to affect stony corals in the tropical Pacific Ocean, Indian Ocean and Red Sea.

Condition	Location	Species affected	Source
Black-band disease	Australia, Egypt, Fiji, India, Papua New Guinea, Philippines, Saudi Arabia, Tonga, United Arab Emirates	<i>Favia, Favites, Acropora, Montipora, Pachyseris, Platygyra, Goniastrea, Leptoria, Turbinaria, Montastraea, Hydnophora</i> 20 species	Antonius, 1987; Chesher, 1984 Glazebrook and Steiner, 1994 Littler and Littler, 1995 Miller, 1996 Korrubel and Riegl, 1998 Fenner, 1998; Cervino, 1998
White-band disease	Australia, Egypt, Fahl Island, Guam, India, Mauritius, Oman, Papua New Guinea, Philippines, Saudi Arabia, United Arab Emirates	<i>Acropora, Goniastrea, Platygyra, Hydnophora, Leptoria, Favia, Pocillopora, Stylophora, Lobophyllia, Porites, Leptoseris, Echinopora, Mycedium, Podabacia, Monitopora, Symphyllia</i> 33 species	Antonius, 1987 Coles, 1994 Korrubel and Riegl, 1998
Shutdown reaction	Saudi Arabia, Tonga	<i>Acropora</i>	Antonius, 1981 Chesher, 1984
White blotch disease	Australia	<i>Acropora</i>	Dinsedale, 1994
Brown-band disease	Australia	<i>Acropora</i>	Dinsedale, 1994
Black aggressive band	Mauritius	<i>Acropora</i>	Antonius, 1995a
Black overgrowing cyanobacteria	Indian Ocean	<i>Acropora, Pocillopora, Porites, Favia</i>	Antonius, 1995a
<i>Halofolliculina corallasia</i> , ciliate			Antonius, 1999
Skeleton eroding band	Egypt	massive corals	Antonius, 1999
Pink line disease	Papua New Guinea, Sri Lanka	<i>Porites</i>	Goreau/Cervino, coral list server
Yellow-band disease	United Arab Emirates; Arabian Gulf	<i>Acropora, Porites, Cyphastrea, Turbinaria</i> 10 species	Korrubel and Riegl, 1998
Fungal infection	Adaman Islands, Indian Ocean	<i>Porites, massive corals</i>	Raghukumar and Raghukumar, 1991; Ravindran et al., 2001
<i>Porites</i> ulcerative white spot disease	Philippines	<i>Porites</i>	Raymundo and Harvell, 2000
Tumors (calicoblastic epitheliomas)	Australia (GBR), Indian Ocean, Hawaii, Guam, Enewatak, French Polynesia, Gulf of Oman	<i>Platygyra</i> ; 8 families; most common in <i>Acropora</i>	Loya et al. 1984 Glazebrook and Steiner, 1994 Coles and Seapy, 1998
Pink-blue disease	Israel	<i>Acropora, Porites</i>	Red Sea Marine Park, 2001
Rapid tissue necrosis	Aquarium Corals	<i>Acropora, Pocillopora, Seriatopora</i> ; 11 others	Borneman, 2001

become a “hot spot” for diseases, due to increased virulence and rapid spread of pathogens, a rapidly expanding host range of coral diseases, a recent emergence of new diseases, and the relatively small, partially enclosed and interconnected nature of the Caribbean basin (Porter and Tougas, 2001; Goreau et al., 1998; Hayes and Goreau, 1998).

It is now recognized that coral diseases have played an important role in the loss of coral cover over the last two decades. For instance, there is good evidence that mortality from WBD has modified the structure and composition of reefs in the wider Caribbean by removing two of the most common and locally abundant species (Aronson and Precht, 2001). However, environmental perturbations and anthropogenic stressors that may have triggered disease outbreaks or contributed to their spread and severity are not fully understood. In addition, our ability to predict the long-term impact of emerging syndromes and coral diseases is hindered by an incomplete understanding of 1) ecological processes and interactions at the species, population and community level; 2) connectivity among reefs and the importance of regional and local water movement patterns as vectors for dispersal of (coral) propagules and water-borne pathogens; and 3) factors that influence the resilience of coral reef ecosystems and their ability to persist under increasing environmental and anthropogenic impacts.

While much research is needed to achieve a better understanding ecological and oceanographic processes affecting coral reef ecosystems and responses of these ecosystems to chronic impacts, there is also an urgency to understand the causes of diseases and their consequences on coral populations. Regional and global coral reef monitoring programs [ReefCheck, Caribbean Coastal Marine Productivity (CARICOMP), Atlantic and Gulf Rapid Reef Assessment (AGRRA)] are providing valuable information on reef condition and threats. Some of these programs also include methodologies for disease surveys (Appendix I). However, there remains a need for targeted studies on coral disease etiology and epizootiology¹, to identify and characterize disease-causing microorganisms, and to understand processes and stressors that may trigger outbreaks of disease and exacerbate their impacts. In addition, resource managers need effective tools to control disease epizootics and reduce coral mortality.

The purpose of this paper is to summarize the type of information needed by resource managers to improve their ability to effectively manage coral diseases, and to discuss impediments hindering progress in coral disease research. Specific examples from the literature highlight some of the existing limitations of available data. The focus of this paper is on the use of rapid ecological assessments, long-term monitoring, and targeted research to address gaps in the understanding of coral diseases, with a

¹Epizootiology is the science that deals with outbreaks of animal diseases, including the prevalence and variability at local, regional and temporal scales, factors involved in the occurrence and spread, and the effect on host populations. It is the equivalent of human epidemiology.

discussion of possible strategies to link coral disease research with management objectives. Descriptions of regional coral disease monitoring protocols and techniques used by scientists illustrate the existing methodologies used to characterize coral disease epizootiology (Appendix A). Priorities for management of diseases identified by the Florida Keys National Marine Sanctuary (FKNMS) in their draft science plan are also summarized (Appendix B).

Key priorities for effective management of coral reef diseases

1. Early warning systems

The challenge: Coral bleaching and disease outbreaks may be linked to unusual environmental events. Yet, it is difficult to reliably predict their occurrence or potential impacts, or to separate the relationships of these events from natural variations in disease occurrence.

The health of coral reef ecosystems is closely linked to physical properties of the environment and ecological linkages among species. Certain global or regional-scale disturbances such as coral bleaching occur during periods of elevated water temperature, high UV radiation, excessive sedimentation and similar stressful events. Such events have been numerous in the Florida Keys (FKNMS) and some of these have been followed one to two years later by the appearance of a disease outbreak (B. Causey, pers. comm, Jan. 2002). However, the relative importance of large-scale phenomena such as global warming and El Niño as a cause of coral mortality, and how much these phenomena affect the prevalence and severity of coral diseases are unknown. By understanding the synchronization of various stressful events and the occurrence of coral diseases, managers may be able to predict potential threats in a timely manner and develop strategies to lessen impacts. This information may also assist in interpreting changes in reef condition that occur in response to these events, and separate these from natural variations. Unfortunately, the wide distribution and remote location of many coral reefs presents challenges for *in situ* monitoring of unusual environmental events and their relationships with bleaching and disease outbreaks.

2. The spatial distribution and temporal variations of coral disease outbreaks

The problem: Limited quantitative information exists on the prevalence and incidence of diseases. Existing data may not be representative over large geographic or temporal scales, or even at local scales, due to the small sample sizes of corals, few reefs examined, and infrequent sampling that is characteristic of most studies.

A summary of all *in situ* observations (up to 1999) on the prevalence, range of species affected, global geographic distribution, and mortality for the “major” diseases has been compiled by Green and Bruckner (2000). This review highlights the large geographic scale over which

diseases have been reported, but many of the records represent qualitative observations only. Currently, detailed quantitative data on the prevalence of diseases are available only for a few locations (e.g., Florida) and diseases (primarily BBD, WBD and white plague). There have been a few recent regional studies to evaluate the health of coral reefs and the occurrence of diseases. This includes the Atlantic and Gulf Rapid Reef Assessment (AGRRA) surveys conducted between 1998-2000 in 22 countries (Kramer et al., *in press*), and a CARICOMP survey of 19 reef sites from 6 countries in the Caribbean (Weil, *in press*).

A. White-band disease (WBD)

White-band disease has been reported from most countries in the Caribbean, and it is believed to be the major factor responsible for the regional decline of *Acropora cervicornis* and *A. palmata* (Aronson and Precht, 2001). However, quantitative data on disease prevalence and incidence are available for few locations, and these show large local, regional, and seasonal variations (Table 3). In addition, long-term monitoring of WBD on relevant temporal scales was conducted only in the USVI (Gladfelter et al., 1977; Gladfelter, 1982; Gladfelter, 1991). WBD was observed on acroporids in many other locations, and extensive mortality occurred at some later date, but conclusive proof to verify that WBD was the leading (or only) source of mortality is not available. Localized losses of these species have been attributed to other factors (storms, predation, bleaching etc.), yet some authors have concluded that a new disease, white pox, is actually the major source of mortality to *A. palmata* in Florida (Patterson et al., 2002). The importance of these factors versus WBD, or their relationships with WBD have not been fully characterized, and research should be targeted to answer these questions.

Table 3. An example of the variation in prevalence of white-band disease reported during the 1980s.

Prevalence of WBD	Location	Author
2-5%; up to 40%	Florida, USVI, Belize	Antonius, 1981
5-26%	British Virgin Islands	Davis et al, 1986
1-2%; up to 64%	U.S. Virgin Islands	Gladfelter et al., 1977
20-33%	La Parguera, Puerto Rico	Goenaga and Boulon, 1992
> 80%	Jamaica; Netherlands Antilles	Rogers, 1985

B. White plague

Studies from the Florida Keys suggest that white plague may have been a significant factor contributing to the declines in live coral cover over the last 20 years (Dustan, 1977; Dustan and Halas, 1987; Richardson et al., 1998 a,b). Plague type II has been one of the most severe disease events in terms

of the number of coral species affected, the rates of tissue loss, extent of mortality, and the rapid spread along the Florida reef tract (Richardson and Aronson, in press). However, it is difficult to evaluate long-term impact on coral populations in Florida because the disease has not been monitored repeatedly (weekly to monthly) over single years or continuously for multiple years (Table 4). The value of frequent monitoring to accurately track the spread of a coral disease and evaluate its effect on coral populations is illustrated by a study from the USVI where researchers monitored coral populations monthly for about 4.5 years. They identified frequencies of plague type II that ranged from 3-58%, with new infections observed each month and an absence of a seasonal correlation in disease prevalence (Miller et al., in press).

The examples presented in Table 4 illustrate the high percentage of stressed, diseased, and dying corals identified in surveys, but these studies often do not distinguish between cause and effect of mortality. For instance, Dustan and Halas (1987) identified numerous dead corals in their transects which may have died from white plague, BBD, sediment-related tissue loss or other factors. Also, diseased corals were identified during surveys, a decline in coral abundance was noted at a later date, but it is not possible to verify that the identified disease was the sole cause of this decline.

Table 4. Reports of white plague from the Florida Keys.

Species	Description of event	Location	Author
<i>M. ferox</i> , <i>M. lamarckiana</i> , <i>C. natans</i> , <i>M. annularis</i> , <i>P. astreoides</i> , <i>S. michelini</i>	1) Outbreak observed in 1975; 2) Highest prevalence in <i>M. ferox</i> (24-73%); 3) Spread of up to 3.1 mm/day in <i>M. ferox</i> ; 4) Mortality of 20-30% of <i>M. ferox</i> in one year.	Carysfort, Elbow, French, Key Largo Dry Rocks, Molasses	Dustan, 1977
Species with plague not identified	1) Corals at 10-21 m declined in abundance between 1974-1982; 2) Skeletons of <i>M. ferox</i> , <i>M. lamarckiana</i> , <i>S. hyades</i> and <i>C. natans</i> recorded along transects 3) BBD, white plague and sediment-related tissue loss prevalent	Carysfort	Dustan and Halas, 1987
Species with plague not identified	1) 362 out of 9800 (3.69%) colonies with white plague; 2) Prevalence same on all reefs (no "hot spots").	19 reefs in Key Largo	Dustan, 1993
<i>A. agaricites</i> , <i>A. lamarcki</i> , <i>C. natans</i> , <i>D. cylindricus</i> , <i>D. stokessi</i> , <i>D. labyrinthiformis</i> , <i>D. strigosa</i> , <i>E. fastigiata</i> , <i>M. decactis</i> , <i>M. mirabilis</i> , <i>M. areolata</i> , <i>M. meandrites</i> , <i>M. annularis</i> , <i>M. cavernosa</i> , <i>S. siderea</i> , <i>S. bournoni</i> , <i>S. michelinii</i>	1) Outbreaks of plague type II from 1995-1997; 2) Plague type II spread throughout the Florida Reef Tract; 3) Epizootics confined to the middle keys in 1995, southern keys and the Dry Tortugas in 1996, and reefs north of Miami in 1997; 4) Up to 38% mortality of one species; 5) Seasonal occurrence (June-October)	23 reefs over 200 km reef tract	Richardson et al., 1998a
<i>C. natans</i> , <i>M. faveolata</i>	Outbreaks of plague type III reported from Florida in 1999		Richardson and Aronson, in press

There are several possible pitfalls associated with infrequent monitoring and sampling of a small number of reefs in one location (Bruckner and Bruckner, 1997):

- Diseases often occur at a low frequency, but they may be clumped in occurrence. Surveys that are not representative, or are too few number, may miss localized aggregations of diseased corals and underestimate disease prevalence or they may focus on localized areas with multiple diseased corals and overestimate disease prevalence.
- Diseases often affect individual hosts for a relatively short time, with new diseases appearing on surrounding corals at frequent intervals. Surveys conducted infrequently are unable to accurately assess the duration of individual diseases and their role in coral mortality.
- Some diseases exhibit seasonal variations in abundance or occurrence. Surveys conducted during certain times of the year may incorrectly estimate disease prevalence or incidence, and the impact of diseases.
- Coral mortality may be caused by numerous factors, in addition to disease. Infrequent surveys may overestimate the role of disease in coral mortality, by not identifying other causes of mortality (e.g., injury, predation, sedimentation) and their relative importance.

3. Relationships with environmental factors, anthropogenic stresses and natural disturbances

The problem: The recent emergence of diseases is often presumed to be related to habitat degradation, increased human impacts, or poor water quality, but linkages with specific factors have not been conclusively determined. Reef managers need to recognize and partition various threats that may affect the occurrence, prevalence, or severity of diseases so their sources can be identified, and possibly reduced through better management practices.

Although it has been difficult to identify the cause of most new diseases, the sudden emergence may be related to deteriorating and changing environmental conditions (Santavy and Peters, 1997). For example, eutrophication may be an important source of stress that compromises a coral's immune system and decreases the resistance to disease, and it may also promote proliferation of pathogens, allowing new diseases to emerge (Hayes and Goreau, 1998). Green and Bruckner (2000) mapped the spatial distribution of coral diseases, and noted that 97% of all locations in the Caribbean affected by coral diseases correspond to areas where human activities are expected to have medium to high impacts, based on the Reefs at Risk analysis².

Porter and Tougas (2001) presented a coral disease stress model that links environmental quality with incidence of disease. In this model, a variety of stresses (elevated temperature, increased storm frequency, elevated nutrients and toxins and decreased water clarity) are predicted to suppress the

²*Reefs at Risk (1998)* presents an indicator of potential threats from coastal development, overexploitation and destructive fishing, impact of inland pollution and erosion, and marine pollution based on a series of distance relationships correlating mapped locations of human activity. It also considers population demography, rainfall, and topography of associated land masses.

immune and disease defense systems in corals. The consequences of reduced health include increases in the number of pathogenic organisms, susceptible species, locations affected by disease, and mortality rates. The authors suggest that the model is applicable to the Florida Keys, where a dramatic increase in the number of diseases, the number of species affected, and the rates of coral mortality were recorded in 40 sites between 1996-1998.

A. Black-band disease

Previous studies quantifying the spatial and temporal distribution and abundance of BBD illustrate the widespread distribution of the disease, but infections exhibit significant variations in prevalence and dispersion (Table 5). BBD occurs on most reefs at a low abundance, although infections may be aggregated, and outbreaks have been reported from numerous locations (Peters, 1993; Bruckner et al., 1997). Goreau et al. (1998) report that BBD often first appears in polluted areas, and infections spread radially outward. They suggest that the abundance of BBD mimics the distribution of human influenced areas, with the largest impacts near sewage outflows and areas of high turbidity from eroded soil. Peters (1993) also noted that BBD prevalence is related to adverse environmental conditions, including warmer than normal temperatures, nutrient loading, increased sedimentation and turbidity, predation, and toxics. Antonius (1998a) identified BBD on reefs in the Red Sea located in close proximity to industrialized areas, with a notable absence of the disease in other locations. In Jamaica, the incidence of BBD progressively increased over 19 months, with the largest increase during or just after a period of unusual rainfall and run-off (Bruckner et al., 1997a). In this study, the incidence of BBD was not correlated with run-off in the most commonly affected species (*M. annularis*; *M. cavernosa* and *Diploria* spp.). However, one species that is thought to be highly resistant to BBD (*S. siderea*) exhibited few infections prior to the rainfall event, with a dramatic increase in bleaching, white plague and BBD in the second year of the study, corresponding to periods of high rainfall and run-off (Bruckner, et al., 1997). Richardson (1995) compiled information on BBD incidence versus nutrients, temperature, light, turbidity, salinity, coral cover, and coral diversity for 200 sites in the Florida Keys. This study may further clarify the role of environmental quality; however, at least initially, she noted that there is no clear correlation.

An extensive, multi-year study conducted in Puerto Rico by Bruckner (1999) suggests that a negative relationship may exist between BBD prevalence and turbidity or elevated nutrients. The lowest prevalence of BBD overall was found near Mayaguez and Ponce, which are some of the most polluted and turbid sites in Puerto Rico due to high sedimentation and nitrification associated with river discharge, agricultural runoff, and direct input of untreated sewage. In addition, the incidence of BBD on a fringing reef off the west coast (Rincon) was highest in spring (May-June) when water clarity was high,

with infections disappearing during the rainy period (July-August) when run-off increased and visibility declined, even though temperatures were approaching their annual maxima. Furthermore, high turbidity was one of the major factors limiting the spatial distribution of BBD in La Parguera. Infections were restricted to shallow water on turbid inshore reefs, even though species susceptible to BBD occurred in shallow and deep water, while BBD occurred to depths of 30 m on offshore reefs with high water clarity. The disease was also common in remote locations around Mona Island, which is 70 km from the mainland of Puerto Rico and lacks permanent inhabitants, industry or agriculture (there is a small Puerto Rico Department of Natural and Environmental Resources facility and camping is permitted) and reefs are unaffected by river discharge (Bruckner, 1999).

Table 5. A comparison of published quantitative studies on black-band disease from the Caribbean.

Location	Survey information	BBD prevalence	Habitat and environmental characters	Author
USVI	22 sites on 7 reefs; 9204 corals; 4 surveys in 1 year.	0.2% of all corals; 5.5% of <i>D. strigosa</i> ; random dispersion; BBD declined in November.	high density of corals (1.3 colonies/m ²) but only 24% of all corals in study area were susceptible to BBD; undeveloped watershed.	Edmunds, 1991
Florida	19 reefs; 9800 corals; single survey.	6%; same prevalence on all reefs.	surveys in 1980s; higher coral cover and abundance than in the 1990s.	Dustan, 1993
Florida	10 sites on 3 reefs; 1397 corals; 3 surveys in 1 year.	0.72%; clumped distribution; seasonal occurrence, but some corals infected year-round.	low density of corals (0.15 colonies/m ²); large annual decline in temperature; possible water quality issues.	Kuta and Richardson, 1996
Jamaica	20 sites on 4 reefs; 5564 corals; bi-monthly surveys for 20 months.	5.2% total over 20 months; max 1.2% at one time; clumped distribution; new infections during each survey; no decline in winter, except in <i>M. cavernosa</i> .	>90% of the corals in sites were susceptible to BBD; moderate density of corals (0.9 colonies/m ²); seasonal periods of high rainfall and runoff.	Bruckner et al., 1997a

The role of physical factors (temperature, light, wave action), anthropogenic stressors (nutrients, sediments, pollutants) and other environmental stressors must be carefully evaluated before concrete conclusions of relationships with disease can be made. Most coastal environments are affected by a suite of anthropogenic inputs, making it difficult to identify any one specific cause associated with deteriorating health or an increased prevalence of disease. Differences in prevalence, severity or impact of diseases may be related to specific anthropogenic stressors, other biotic or abiotic parameters or multiple factors. For instance, habitat characteristics and composition including the cover and abundance of susceptible corals, the amount of macroalgae, or presence or absence of certain key indicator species such as *Diadema* may influence coral diseases.

4. Standardized nomenclature to describe, identify and differentiate diseases

The problem: Inconsistent terminology for coral diseases, absence of standard requirements for reporting of new diseases, and lack of field identification tools hinders our ability to obtain accurate information on the types of diseases, their prevalence, and their impacts.

There is considerable disagreement among coral disease specialists about many of the fundamental aspects of the newly emerging diseases. This includes a profusion of names describing so-called “new diseases” which may represent conditions caused by pathogens, predators, environmental perturbations, or other factors (Goreau et al., 1998; Bruckner and Bruckner, 1998, Richardson, 1998). In the majority of cases, researchers have assigned names to coral syndromes based on the appearance of the microbial community (e.g., BBD), the color of the affected tissue (YBD), or the pattern of loss of coral tissue (WBD). In a recent review of coral disease literature, Green and Bruckner (2000) noted that 29 different “disease states” had been reported. In some cases, a variety of similar names were used to describe different diseases; in others, different names were used by separate researchers to describe apparently similar disease signs. For instance, the term white-band disease (which occurs only in *Acropora* spp. in the Caribbean) has been used to describe similar signs in massive and plating corals that are reported elsewhere as white plague (Table 1). Also, similar gross signs of coral abnormalities that are classified as the same disease have been later shown to have different histological characteristics (Santavy and Peters, 1997). There has also been an uncertainty as to whether what is being described as a coral disease is actually a biotic or abiotic disease, or whether it is caused by other factors. This includes rapid wasting disease (later renamed rapid wasting syndrome), which has been shown to be the result of focused biting by *Sparisoma viride* (Bruckner and Bruckner, in press), a phenomena that was documented over 100 years ago. Another example is ridge mortality disease (Abbott, 1979), a condition that affects brain corals and other species. This condition is always associated with *Stegastes planifrons* predation and the development of algal lawns (Bruckner, 2001), and has been termed damselfish ridge-denuding syndrome by other authors (Williams and Bunkley-Williams, 2000).

In the last five years there have been numerous “review articles” on coral diseases (Cervino and Smith, 1997 a,b; Santavy and Peters, 1997; Peters, 1997; Bruckner and Bruckner, 1997c; Richardson, 1998; Goreau et al., 1998; Hayes and Goreau, 1998; Harvell, et al., 2000; Williams and Bunkley-Williams, 2000; Borneman, 2001; Bruckner, 2001; Richardson and Aronson, in press). Many of these provide descriptions and lists of reported diseases, but the summaries often lack a thorough review of existing literature, and may be heavily biased towards work by one or a few researchers. In addition, some of these articles have been published without a thorough and unbiased peer-review. In some cases,

these articles have discounted many of the presumed diseases, while they identify additional “new” syndromes without any supporting quantitative field or laboratory studies. In other cases, even after considerable supporting evidence is available that discounts a presumed disease, references to this condition continue to appear in scientific literature and popular articles. Along with the continued controversy regarding many of the presumed new diseases, many of the articles draw conclusions on the recent spread, prevalence, associated factors, or potential impact based on qualitative information, in absence of sound quantitative science. Without a coordinated and cooperative effort among all researchers working on coral diseases, this misinformation is likely to continue to proliferate, severely affecting our ability to respond to a growing coral reef crisis.

5. Factors that facilitate the occurrence, spread and transmission of pathogens

The problem: The source of emerging diseases and their mode of spread and transmission is unknown.

It is currently unclear whether the recent emergence of diseases is associated with the introduction of pathogenic organisms from terrestrial environments, or whether disease-causing microbes have always been present but have only recently become pathogenic due to deteriorating environmental conditions and/or reduced host resistance. A critical area of research is the identification of disease sources and vectors, and natural reservoirs for pathogens. Possible sources include the introduction of novel pathogens via aerosols (e.g., African dust), run-off, sewage, vessel discharge, the introduction of exotic species as well as the movement of water masses as a vector for dispersal (NCORE, 2001).

Black-band disease now occurs in most reef environments in the western Atlantic. In many locations BBD increases seasonally in summer and fall, with infections becoming rare or absent during winter when water temperatures decline (Kuta and Richardson, 1996). Factors responsible for the reappearance the following year are not completely understood, but the occurrence of the dominant microorganism (*Phormidium corallyticum*) outside of the black-band association has been reported. During surveys in Belize, Florida and the Bahamas, Taylor (1983) reported the occurrence of *P. corallyticum* from shaded areas under reef limestone not in association with living corals. Richardson (1997) identified non-pathogenic biofilms that contained *P. corallyticum* and other cyanobacteria in sediment patches on the surface of healthy, BBD-susceptible corals. However, none of these developed into active BBD after 12-17 months. Santavy and Peters (1997) also identified algal mats that contained *P. corallyticum* filaments at the base of a reef in the Bahamas.

Although *P. corallyticum* was not identified in plankton tows, injured colonies have become infected with BBD when placed downstream from an infected colony, suggesting that the disease is transmitted in the water column (Rützler and Santavy, 1983; Antonius, 1985b). In Jamaica, a BBD epizootic was observed at the eastern end of a reef system and it progressed to the west over 20 months, in the direction of the major water current (Bruckner et al., 1997 a). In this study, portions of the BBD mat became dislodged during periods of high wave action, and floated towards other corals down current. Although the dislodged BBD mat was not observed to land on and infect a new host, a high number of new infections were identified down-current 1 to 3 weeks after periods of high wave action (Bruckner et al., 1997 a). While water motion provides a vector for transmission of BBD, under extreme wave conditions, water turbulence may actually limit BBD progress and prevent new infections from becoming established. On exposed shallow (1-2 m) reefs on the north coast of Puerto Rico, a high incidence of BBD has been observed during spring and summer, coinciding with periods of low to moderate wave surge. In fall, although temperatures are approaching their maximum (29°C), BBD virtually disappears from exposed reefs; the decline corresponds to the onset of fall and winter storm waves. In certain protected locations, BBD reaches its maximum annual abundance during this period (Bruckner, 1999).

There is the possibility that pathogenic microorganisms may be transported into coastal environments along with soil during periods of run-off (Epstein et al., 1998). *Aspergillus sydowii*, a fungus identified as the cause of aspergillosis in sea fans, is reported to be of terrestrial origin. One hypothesis is that *A. sydowii* was transported to coral reef environments via run-off; however, fungal isolates from the terrestrial environment are not pathogenic to corals. An alternate hypothesis is that *A. sydowii* was transported to Caribbean reefs in clouds of dust from western Africa (Weir et al in press).

Another new disease, white pox, has been recently characterized as being caused by a common bacteria, *Serratia marcescens* (Patterson et al., 2002). These authors conclude that the bacteria is associated with the human gut and may be introduced into the marine environment via human waste. However, others have noted that the bacteria is very common in the intestines of other animals, including fish. One fish, *Sparisoma viride*, forms territories within areas dominated by *A. palmata*, and the fish will frequently defecate as they swim over standing elkhorn coral colonies. The fecal matter will accumulate on the surface of elkhorn branches during calm periods, and will cause mortality. It is possible that the parrotfish are the main vector for the bacterium that causes white pox, negating the linkage with human sewage.

6. Effects on population dynamics, community structure and ecosystem function

The problem: Long-term consequences of diseases on coral reefs are largely unknown. Managers need information on the fates of corals, the resilience of coral populations, potential effects on associated species, and long-term impacts on coral reef composition, structure and function.

Hayes and Goreau (1998) concluded that mortality from coral reef diseases has the potential to be more detrimental to the survival of coral reefs than sedimentation, pollution, physical degradation, or all other threats combined. It is clear that coral diseases can have a major impact on the structure and composition of coral reef ecosystems, as evidenced by the regional spread of WBD and devastating levels of mortality to acroporid populations. However, the fates of individual colonies, the resilience of coral populations, and the long-term effects on associated species and the ecosystem are a complex interaction of multiple factors, of which disease may be one important factor. For instance:

- The virulence of pathogens is likely to vary among diseases and affected species, with additional seasonal and spatial variations and possible synergistic impacts from other disturbances.
- The role of coral disease as a community structuring force depends on the magnitude, duration and frequency of reappearance of the disease, and the recovery time of the system.
- The recovery time depends on the life history strategies of the affected species, including rates of recruitment and growth.
- The impact from disease is likely to vary depending on the overall condition of the reef, including the cover and composition of reef building corals, the presence and abundance of critical indicator species (e.g., *Diadema*), and the degree of physical, environmental and anthropogenic stressors.

A. Fates of infected corals

Coral diseases can cause rapid rates of tissue mortality, but rate of tissue loss within a colony is highly variable depending on the disease, affected species, location and season (Table 6). Diseases may produce mortality of affected corals over varying periods of time, although diseases more often cause only partial mortality (Bruckner and Bruckner, 1997 c). While corals have the potential for recovery once the insult is eliminated, the reduced size of affected colonies may affect growth, reproduction, resistance to injury and other diseases, and competitive abilities of the surviving polyps (Edmunds, 1991). Colonies affected by disease are vulnerable to reinfection, infection by other diseases, or subsequent attacks by predators and overgrowth by competitive invertebrates and algae (Kuta and Richardson, 1996; Bruckner and Bruckner, 1997 a; Antonius and Riegl, 1997; 1998). For instance, Bruckner and Bruckner (1997 a) reported a recurrence of BBD later in the season or in subsequent years (1995-1996) in 39 corals (11.4%) identified with BBD in 1992 or 1993.

B. Effects on coral populations

Coral diseases, once they invade a reef system, may become a chronic feature of that system which undergo periodic escalations in abundance and virulence (Bruckner, 1999; Richardson and Aronson, in press). From an evolutionary standpoint it is not beneficial for a microbial pathogen to eliminate all of its host species. A coral species may occasionally suffer periods of high mortality from disease, however survivors are likely to persist by becoming resistant, the pathogen may undergo a decrease in virulence, or that disease may occur on a wider range of host species or over a wider distribution. For instance, Dustan (1977) observed an epizootic of white plague in 1975 that was predicted to result in the disappearance of *Mycetophyllia ferox* from Carysfort Reef in the Florida Keys. This species still occurred on this reef in 1984, and most colonies were not observed with white plague, suggesting that this species developed a resistance to the pathogen and new colonies recruited onto this reef to replace those lost to disease.

Table 6. Rates of tissue loss and extent of mortality from black-band disease.

Affected species	linear spread	Tissue loss (observation period)	Total loss/season	Location & Source
<i>M. annularis</i> (n=2) <i>D. strigosa</i> (n=1)	0.3-6.2 (mm/day) 2.4-4.0 (mm/day)	64-746 cm ² (41 days)	1509-4820 cm ²	Belize; Rutzler et al., 1983
<i>D. strigosa</i> (n=12)	no data		7 colonies lost >75% live tissue; 3 lost 50-74%; 2 lost <25%	USVI; Edmunds, 1991
<i>S. siderea</i> (n=93) <i>M. cavernosa</i> (n=48)	no data	1810 cm ² (2 years) 1630 cm ² (2 years) Small colonies (mean=239 cm ²) died in 1-8 months.	Large colonies of Ss lost 50% live tissue in 2-3 years; large colonies of Mc lost 68% of live tissue.	Jamaica; Bruckner and Bruckner, 1997a
<i>M. annularis</i> <i>D. strigosa</i>	up to 1 cm/day			Antonius, 1981
<i>M. annularis</i> <i>M. faveolata</i> <i>M. franksi</i> <i>M. cavernosa</i> , <i>D. strigosa</i> <i>D. clivosa</i> <i>C. natans</i> <i>S. siderea</i> (n=210)	0.5-8.1mm/day Variations among species, reefs and depths. Slowest spread among <i>S.</i> <i>siderea</i> in deep water and fastest among <i>M. annularis</i> in shallow water.	1.3-100 cm ² /day 107-1329 cm ² total Mean duration of BBD= 46-220 days.	Mortality rates depended on species, reef and depth and also size of coral, length of BBD and duration of BBD. Mean loss (pooled by species) ranged from 18%-96%.	Puerto Rico; Bruckner, 1999
<i>M. faveolata</i> (n=9) <i>D. strigosa</i> (n=51)	4.1 mm/day 2.0 mm/day			Honduras; Griffin, 1998

(Dustan, 1987). In contrast, *M. faveolata* colonies showed few signs of recent tissue loss in 1975, while many were infected with white plague in 1985 (Dustan, 1987). More recently, Richardson et al. (1998) reported an outbreak of white plague in the Florida Keys which initially targeted *Dichocoenia stokessi*, a species that was not affected by plague in the 1970s or 1980s. Between 1995-1997, white plague spread among 17 coral species in the Florida Keys and the rate of tissue loss was much greater than that observed in earlier years. These observations led Richardson (1998) to conclude that this disease outbreak involved a more virulent pathogen than that reported by Dustan (1977). The outbreak was one of the most severe disease events in terms of diversity of species affected, mortality rates, and rapid rate of spread throughout the reef tract, but was believed to have minimal long-term impacts, as disease incidence was greatly reduced in these sites in subsequent years, and the pathogen identified as the cause no longer appeared to induce disease signs (Richardson and Aronson, in press). Another outbreak (termed plague type III by Richardson) was observed in 1999 on the same reefs; rates of tissue loss exceeded that reported for earlier outbreaks, and large faviid corals (*M. faveolata* and *C. natans*) appeared to be most susceptible (Richardson and Aronson, in press).

C. Effects on ecosystem structure and function

When diseases have particularly severe impact on populations of a single species, the effects can extend over the entire ecosystem. Diseases may exert a major control on biodiversity by eliminating certain highly susceptible species and losses may reduce the capacity for the ecosystem to sustain associated species. Until about 1980, more than 95% of the total coral coverage in the reef crest and shallow fore reef consisted of *A. palmata* and *A. cervicornis*, with colonies forming extensive high relief thickets (Gladfelter et al., 1977). Populations of acroporids were reduced to unprecedented levels throughout the Caribbean during the 1980s and 1990s (Aronson and Precht, 2001). The combination of coral mortality, declining herbivory (associated with the *Diadema* die-off and overfishing of parrotfish and surgeonfishes), and possibly elevated nutrients have precipitated a phase shift from coral to macroalgal dominance (Hughes, 1994; Aronson and Precht, 2000). In many locations, *A. palmata* colonies suffered near total mortality, but erect skeletons remained in growth position, continuing to provide high-relief habitat. However, exposed skeletal surfaces were rapidly colonized by invertebrates and algae and skeletons are being progressively bioeroded, dislodged and removed during storms. The collapse of acroporid skeletons is associated with the elimination of high relief, three-dimensional topography. Long-term consequences may include loss of critical habitat and refuge for a wide number of associated species, reduced rates of reef accretion, and reduced shoreline protection from tropical storm waves.

7. Measures to mitigate disease impacts

The problem: Resource managers need tools to combat disease epizootics, prevent future outbreaks, and reduce the time needed for recovery of affected reefs and species

In the past two decades numerous reef mitigation and restoration projects have been undertaken in response to coral mortality, in attempt to offset these losses. While the majority of these projects have been undertaken in response to direct human impacts (e.g. ship groundings), limited restoration efforts have been conducted to rehabilitate areas degraded by coral mining, blast fishing and other human impacts. A variety of treatments have also been attempted for corals with disease, including application of antibiotics, removal of the microbial community, shading, and biological controls (Peters, 1997; Griffin, 1998; Bruckner, 1999; Franklin, 1999; Borneman, 2001). In addition, attempts have also been undertaken to treat corals with YBD (A.W. Bruckner, unpubl. data) and white plague (S. Miller, et al., in press) involving application of putty or clay over affected tissue. These approaches have had varying rates of success, but most attempts have been conducted on a very small scale and often only limited efforts have been made to determine the long-term benefits. As an alternative to direct treatment of diseased corals, novel biological restoration approaches are being applied in Florida and other locations. These approaches are designed to enhance recruitment of corals, reduce coral mortality, and improve habitat quality, and include the reintroduction of *Diadema antillarum* as a tool to reduce macroalgae and induced settlement of coral larvae (M. Miller, unpubl data).

In response to an outbreak of BBD in the Florida Keys in the 1980s, Harold Hudson attempted to treat colonies through aspiration of the BBD using a boat-operated vacuum pump, followed by application of modeling clay over the affected area to prevent reinfection. Unfortunately, colonies were not tagged or followed to determine their fates. In 1998, Hudson and Franklin repeated their initial experiments in response to another outbreak of BBD (Hudson, 2001). An experimental approach to treat BBD corals was undertaken by Bruckner (1999) in Puerto Rico. The experimental design included 1) removal of the BBD using a large hypodermic syringe to aspirate the BBD. In these experiments, the affected area was unmanipulated, covered with modeling clay, or covered with underwater putty; 2) application of underwater putty directly over the BBD; 3) shading using various grades of sun screen; and 4) addition of *Diadema antillarum* to cages containing BBD-infected corals. All treatments were effective to varying degrees, with the highest success (>95%) achieved with the use of underwater putty, and very low success when removal was combined with modeling clay (Bruckner and Bruckner, 1998d). Putty proved to be a highly efficacious treatment which prevented further destruction of century-old colonies, and may be most suitable when applied on a small scale for those colonies most important to

the persistence of the reef (e.g., the largest colonies of slow-growing massive species). However, it may be impractical to treat a large number of corals due to the time required to manipulate infections and the expense associated. Because BBD is spread by water motion, care must also be taken to remove the bands without liberating filaments into the water column, which could result in additional infections downstream.. There are a few other limitations associated with treatments discussed for BBD: 1) aspiration of BBD followed by application of clay may be unsuccessful because any remaining *Phormidium* filaments are capable of emerging from within clay; 2) shading may cause coral bleaching; 3) *Diadema* may injure coral tissue when grazing pressure exceeds available algal resources.

Hayes and Goreau (1998) recommended rapid isolation and removal of diseased colonies to provide protection for other reef constituents against the spread of disease. However, this approach is likely to have minimal benefits, as the removal of an infected host may actually promote dispersal and spread of the pathogen, and it may result in considerable habitat damage. Furthermore, low background levels of disease may be a normal occurrence on most reefs that require minimal management interventions, while a disease epizootic is of much greater concern and yet it would be very impractical to remove all corals affected during a disease epizootic. Another proposed strategy, that may have severe long-term consequences, involves the application of antibiotics. Broad application of antibiotics are not recommended, as they may affect beneficial microbes and can lead to the proliferation of antibiotic-resistant pathogens. However, some researchers have proposed adding antibiotics to putty or clay, to restrict treatment to affected areas and contain the antibiotic, which may be much more plausible.

Proposed strategies to address management needs for coral diseases

1. Early warning systems

Remote sensing of coral reefs, using high resolution satellite sensors, airborne LIDAR and hyperspectral coverage, and acoustic mapping methods are important tools to characterize ecosystem types and features, classify human use patterns, and determine sea surface temperature, water mass movements, light exposure, and other physical properties. Satellite remote sensing measurements (ocean color, surface winds, and insolation data) are being used in NOAA's Coral Reef Watch Program to conduct thermal bleaching surveillance of coral reefs world-wide. In this program, oceanographic data are integrated with a GIS spatial database to produce high resolution "HotSpot" (bleaching anomaly)

charts available on the world wide web³. Also, a series of moored buoys are being deployed as part of a Coral Reef Early Warning System (CREWS) in the NWHI, U.S. Pacific, Florida, the Bahamas and other locations. These permanent oceanographic stations provide detailed *in situ* measurements of a variety of water quality parameters including water and air temperature, light (PAR and UV), nutrients, wind speed and direction, and water currents. Through the Coral Health and Monitoring Program (CHAMP) coral-list server, NOAA regularly announces sites that are affected by high sea water temperatures as a warning for potential bleaching. Researchers are then able to verify the onset of bleaching, and subsequently monitor impacts associated with the event. For instance, these data demonstrated accurate capabilities for detection of El Niño-induced coral reef bleaching in 1998. Some coral diseases become more prevalent during warm-water periods, and “HotSpot” data may also prove useful in understanding relationships between coral bleaching and disease. It is important to link oceanographic data with rapid assessments for disease. Real-time information on environmental stressors could provide valuable insight into identifying indicators of reef health, and factors that trigger disease outbreaks.

2. The spatial distribution and temporal variations of coral disease outbreaks

Specific monitoring programs need to be developed to provide information on disease occurrence and progression at a variety of temporal and spatial scales using standardized techniques suited to answer specific questions on coral diseases. Because diseased corals often occur at a low frequency and in a clumped distribution, surveys must encompass large areas to quantify the distribution and abundance of diseased colonies (Edmunds, 1991). Surveys must also be conducted relatively frequently to identify temporal variability, quantify synergistic impacts, and distinguish mortality from disease from mortality caused by other factors.

An effective program should include a three-tiered approach. First, rapid assessments conducted on a regional scale can elucidate large-scale processes such as the transmission of regional-scale water-borne pathogens, and serve as an early warning system to identify and track disease outbreaks before irreparable harm occurs. Targeted monitoring can provide information on a particular disease, and intensive seasonal or weekly monitoring of specific reefs can help managers understand the effect of a particular disease at a local scale, or the role of a localized stressor on disease processes and impacts. Monitoring of individual colonies affected by disease is also necessary to document patterns of spread, the rate of spread and amount of mortality sustained from individual infections, impacts of diseases at

³NOAA/NESDIS Coral Reef Watch incorporates the Coral Reef Early Warning System (CREWS) and OAR/AOML international coral reef information system. This information is available through the Coral Health and Monitoring Program (CHAMP) Web Page: <http://www.coral.noaa.gov>.

colony or population levels, and relationships between disease occurrence and other natural or anthropogenic impacts.

A variety of approaches have been used to identify the prevalence (number of affected corals divided by the number of corals examined) and incidence (number of new colonies with disease occurring in a population over a given time divided by the number of unaffected individuals at the beginning of the time period) of coral diseases, including quantitative, semi-quantitative and qualitative surveys (Table 7; Appendix I). Many monitoring programs rely on repeated sampling of permanently marked transects, quadrats, or radial study sites. Detailed monitoring of particular reefs, through the establishment of permanent sites, can provide information on disease incidence. In addition, long-term monitoring of individual tagged colonies affected by microbial pathogens can provide detailed information on the duration of infections, frequency of reinfection, and effect at the species level.

Monitoring programs that involve the reexamination of permanent sites may be impractical for sampling more than a few sites or reefs, and may underestimate or overestimate the prevalence or potential impacts. Because diseases often occur at a low frequency or they exhibit a clumped distribution, a large number of transects may be required within each reef to accurately estimate disease prevalence. To obtain a more realistic measure of disease prevalence overall at a larger (country-wide or regional) scale, and to determine factors responsible for the increase and spread of diseases, a repeated measures approach consisting of multiple random belt transects positioned throughout each reef environment is necessary.

Specifically, a monitoring program implemented for coral diseases at the community level is needed to obtain information on:

- The type of disease, coral species affected, and variability among species.
- The prevalence and incidence of diseases, the duration of a disease outbreak or bleaching event, and the consequences to affected species.
- Physical parameters (depth, water clarity, temperature etc.), biological factors (predators, algal abundance, etc.), and environmental stressors and anthropogenic impacts associated with the outbreak.
- Local, regional, and global distribution of diseases and temporal and spatial variations.

Table 7. Monitoring and assessment techniques used for coral diseases.

Monitoring Approach	Condition examined. Description of technique	Author
Timed swims	<i>All syndromes.</i> Diver swims for 30 minutes along a single depth gradient and examines all corals in a band approximately 2 m wide recording the type of disease, species and number of affected corals. Disease occurrence is categorized as rare (1-3 cases), moderate (4-12 cases), frequent (13-25 cases), abundant (26-50 cases), epidemic (51-100 cases) or catastrophic (greater than 100 cases).	Antonius, 1995a
Radial sites	<i>Prevalence and incidence of BBD:</i> Total number of infected and uninfected colonies of each species present are recorded within a circular area (10 m radius; 314 m ²).	Edmunds, 1991; Kuta and Richardson, 1996; Bruckner et al., 1997a
Radial belt transect	<i>All diseases on stony corals and gorgonians.</i> Total number of affected and un affected colonies of each species present are recorded within the outer 8- to 10-m of an arc radius (circular area, 10 m radius; 314 m ²).	Santavy et al., 2001
Linear transects	<i>A. Changes in composition, size and cover.</i> Twenty 25m transects on Carysfort Reef from 0.3 to 21m depth. Permanent transects were extended parallel to each other and subdivide an area approximately 25 m X 300 m. Mean colony size (length under the transect line) and distance between colonies were recorded; colonies were inspected for signs of mortality (disease, physically damaged areas, sediment covered tissue or other blemish). <i>B. Condition of stony corals.</i> Minimum of ten 10m transects per reef/depth; surveys at 3m and 10m depth; record species, size, percent recent and old mortality of all colonies that touch the transect line; examine whole colony for diseases or other source of mortality.	A. Dustan, 1985 B. AGRRA, 1998
Belt transects	<i>A. Sea fan disease.</i> One or two 4m X 12.5-50m transects per country. <i>B: All diseases; corals and gorgonians.</i> Ten 10 X 2m belt transects (level 1) and three 20 m X 2 m belt transects per depth; minimum 9 transects at each site (level 2). <i>C. Reef condition.</i> Three 20m X 5m transects per reef; divers note presence and proposed type of disease and affected species, but do not collect quantitative information.	A. Nagelkerken et al, 1997; B. CARICOMP (Appendix I) C. Reef Check
Quadrats	<i>A. BBD in Gorgonians.</i> Prevalence: three 100 X 250m quadrats; incidence: one 10 X 10m quadrat; on e reef <i>B. Sources of mortality.</i> Photostations 2 X 2.25 m; 2 per reef; 2 reefs to identify sources and amount of mortality or time; complemented with permanent chain transects extended over 25m to measure changes in percent coral cover, diversity, and colony size. <i>C. Coral reef monitoring.</i> 160 stations, each 2m X 22m; sampled 40 sites in the FKNMS. 15-minute survey of each quadrat to complete a species inventory, identify species that are affected with bleaching or disease, and record the type of disease. No data collected on disease prevalence. Also run video transects. Sites reexamined once each year from 1996-2000.	A. Feingold, 1988; B. Porter and Meier, 1992 C. Jaap et al., 2000

3. Relationships with environmental factors, anthropogenic stressors and natural disturbances

If there is a link between anthropogenic pollutants, nutrients, and sedimentation and the prevalence of coral diseases, disease occurrence is unlikely to be randomly distributed at regional scales. An understanding of the global epizootiology of diseases constitutes a first step in examining relationships between human impacts and the occurrence of coral diseases. A comprehensive database needs to be developed linking all accurate field reports by trained observers as well as observations and photographs from amateur observers to understand the impacts of coral reef diseases on a global scale

(Goreau et al., 1998). The Global Coral Disease Database⁴, developed by NMFS in conjunction with UNEP's World Conservation Monitoring Centre (WCMC) provides a first attempt at analyzing the global distribution of diseases (Green and Bruckner, 2000). This web-accessible database provides information on the reported types of disease, species affected, and global distribution of each disease. Disease occurrence can be mapped by particular disease, location, or year. Currently, the database contains all published records identified through 2000, as well as volunteer disease reports; a form is available for reporting disease data electronically. The database is being expanded to include new regional monitoring data from AGRRA, CARICOMP and other sources, with a separate category for volunteer diver reports of disease (non-experts). Furthermore, additional layers will be added to the GIS database that map environmental and anthropogenic parameters. This information will facilitate interpretation of disease epizootics and causative factors involved in the development of these outbreaks.

In addition, targeted research that combines both laboratory and field methods is needed to understand the influence of various stressors on coral diseases, bleaching and mortality. These efforts should focus on establishing cause-effect thresholds of anthropogenic pollutants and environmental stressors, and capabilities for providing early warnings of disease outbreaks. For example this could include studies on:

- Mechanisms and factors that increase/decrease host susceptibility to disease.
- Cellular, mechanical, chemical and behavioral mechanisms that impart host resistance and host defense to diseases, and the ability of a host to counteract disease agents.
- Genetic adaptations and resistance of specific clones of coral colonies.
- Identification of biomarkers of exposure or effect of stress to corals resulting from anthropogenic impacts or environmental perturbations.

4. Standardized nomenclature to describe, identify and differentiate diseases

A method to categorize disease terminology and standardize the list of recognized diseases is essential in attempting to assess the global implications of disease. Classifying the various visual categories of recent mortality as well as deviations from normal coloration or morphology is necessary to correlate damage with causes (Santavy and Peters, 1997). Field observations must be combined with techniques from biochemistry, histopathology, and microbiology to determine whether diseases are microbial or abiotic in nature, to identify and differentiate disease-causing microorganisms, megafauna

⁴<http://www.unep-wcmc.org/marine/coraldis/home.htm>

(corallivores) and environmental stressors associated with each condition, and to develop molecular tools to validate field identification of diseases that are difficult to visually identify (e.g., aspergillosis).

As a first step, the list of all known and presumed coral diseases (Table 1 and 2) should be evaluated for its validity. Accepted terminology should reflect the first name assigned to a condition, and terminology to describe presumed “new diseases” should be discarded if the condition was previously described. For instance, the condition “rapid wasting syndrome” described in 1998 was called “parrotfish white spot biting” in literature from the 1980s. Separate terminology should also be used for syndromes that visually differ to avoid confusion. This includes “yellow-band disease” which was first described in acroporids and other species in the Arabian Gulf, and later used to describe an unrelated condition in faviid corals (primarily *M. annularis* complex) from the western Atlantic (Cervino et al., 2001). Santavy and Peters (2000) recommended the use of yellow-blotch disease for the condition affecting *M. faveolata* and *M. annularis*, thereby differentiating it from the yellow-band disease affecting Pacific corals.

Disease nomenclature may evolve from descriptive terminology to etiological and pathognomonic characterizations (e.g., aspergillosis) (Hayes and Goreau, 1998). Hayes and Goreau (1998) suggest that conditions where no microbial pathogens are identified may be better classified as stress responses, while pathogenic diseases should be renamed to indicate the specific organisms implicated as the causative agent. However, several contributing microbes may be required to produce a disease sign, such as that observed in BBD, making a single name (e.g., cyanobacteriosis) impractical or incorrect. In other cases, the proposed pathogen may not be the initial cause, but a secondary invader. Conclusive demonstration of the causative agent (and verification according to Koch’s postulate) has proven difficult, and positive identification of the cause of some diseases may not be possible in the immediate future. Because diseases manifest on corals in a limited number ways, corals that exhibit specific disease signs in the field may be affected by a variety of pathogens that differ spatially or temporally (e.g., the pathogen for plague may differ depending on the location, affected species, or other factors).

There are several sources of photographs and descriptions of coral diseases that can assist in developing a list of recognized coral diseases and descriptions to allow field identification of these diseases. These include 1) a coral diseases web site by McCarty and Peters⁵; 2) underwater coral disease and predator identification cards by Bruckner and Bruckner designed to help differentiate signs of disease from predation (NOAA, 1999); and 3) a recent chapter “Coral health and mortality: recognizing

⁵http://ourworld.compuserve.com/homepages/mccarty_and_peters/coraldis.htm

signs of disease and predation” in Humann and Deloach, *Reef Coral Identification, Florida, Caribbean, Bahamas* 2nd edition (Bruckner, 2001), which contains extensive descriptions and photographs of diseases, signs of predation, and coral overgrowth. A coral disease identification CD-ROM is being developed by NMFS and UNEPs-WCMC for distribution to dive shops throughout the Caribbean to complement volunteer diver disease identification programs. One additional source of information under development is a comprehensive coral disease website within the CHAMP database ⁶ designed to provide help in disease identification and research approaches for laymen, scientists, managers, and disease specialists. This will contain an identification key for diseases, high resolution photographs showing diseased corals, tissue sections, and electron micrographs, a Frequently Asked Questions (FAQ) list, a glossary, contact information for coral disease researchers, a list of publications on disease, and PDF files of relevant publications.

5. Factors that facilitate the occurrence, spread and transmission of pathogens

A multidisciplinary effort is necessary to conduct controlled experiments that evaluate possible sources of pathogens, identify mechanisms of dispersal and spread, and develop strategies to reduce the potential for introduction and spread of pathogenic organisms. This includes:

- Evaluation of the role of water circulation as a vector for dispersal of pathogenic microorganisms.
- Examination of microorganisms transported in African dust, through species introductions, from discharge of ballast water, and other possible external sources of pathogens.
- Examination of microbial populations found in sediment adjacent to corals and microbiota associated with coral mucus.
- Evaluation of other possible sources of pathogens, including algal and cyanobacterial mats that occur in benthic habitats outside of disease associations.
- Identification of conditions and factors that promote the proliferation of pathogenic microorganisms and/or affect their virulence.

6. Effects on population dynamics, community structure and ecosystem function

Long-term monitoring of individual colonies and species affected by disease should be conducted to determine their fates, factors that may increase or decrease the amount of mortality and their potential for recovery or recolonization, and the effects of coral mortality on associated species and coral reef community structure and function. Relevant studies include:

⁶<http://www.coral.noaa.gov/>

- Monitoring of individual colonies to determine the rate of tissue loss, extent of mortality, effect on physiological processes such as growth and reproduction (for corals that experience partial mortality), differences among and within diseases, coral species, seasons, depths and locations, and synergistic effects of other stressors.
- Evaluation of the fates of coral skeleton exposed by diseases, including the potential for corals to regrow over the affected areas, extent of colonization by algae and invertebrates (especially organisms that bioerode coral skeletons), and the extent of coral recruitment.
- Examination of relationships among reef-building coral species and associated species and in the composition, structure and function of reef communities, and the role of diseases in structuring communities and influencing relationships among predators, competitors and other reef-associated organisms.

7. Measures to mitigate disease impacts

The ability for resource managers to effectively address coral disease epizootics and mitigate their impacts relies on an understanding of the causes, species affected, spatial and temporal prevalence and incidence, and synergistic effects of climate change, human impacts and other stressors (e.g., hurricanes). Research should focus on the development of techniques to prevent the occurrence of diseases, cure diseases, and minimize the effects of disease. This includes:

- Strategies to reduce relevant stressors that are correlated with the occurrence, virulence or spread of diseases.
- Novel biological, ecological and physical techniques to improve habitat quality and water quality, and potential reintroductions of ecologically important species that were formerly abundant but have declined due to human pressures or other disturbances, including predators, competitors, or other organisms that may help counteract the disease-causing agent.
- Treatments for corals with disease, including efforts to reduce host contact with pathogens and increase host defense and resilience.
- Restoration approaches to enhance coral recruitment and introduce disease-resistant clones of species impacted by disease that are unlikely to recover naturally in the short term.

All efforts to mitigate diseases should involve hypothesis-driven experiments to determine the effectiveness of these approaches.

CONCLUSION

Most managers and policy makers are aware of the continued, rapid degradation of coral reefs, and they recognize that many of the stressors responsible for these changes are increasing in severity. Coral diseases are receiving increased attention by scientists and the media as an important source of coral mortality, but several factors hinder the development and implementation of strategies necessary for

rapid response to diseases, and management of disease-related impacts. This includes difficulties in obtaining funding for monitoring and targeted research needed to address many unanswered questions regarding diseases. It is also apparent that there is a lack of coordination among research scientists which must be overcome. Research on coral diseases requires an approach that combines ecological monitoring with biochemistry, molecular biology, histology, toxicology, physical oceanography, ecology, taxonomy and other laboratory and field methods. An interdisciplinary approach is necessary to identify, differentiate and characterize coral diseases and their consequences, and understand relationships among diseases and other biotic and abiotic factors. To achieve an understanding of diseases at local to global scales requires considerable investment of time and effort to survey large areas of reef, and to conduct these surveys at frequencies that are sufficient to document the duration of the condition, seasonal patterns, or chronic effects. These studies must be combined with research to elucidate relationships with environmental and anthropogenic stressors, which may help explain differences observed at regional to local scales, and to identify approaches to mitigate disease impacts.

The policy and decision-making processes necessary for effective conservation of reef ecosystems, including the development of appropriate actions to manage coral diseases, are influenced by equally complex socioeconomic and political systems. There is a limited awareness and recognition of the potential global implications of disease by policy makers and the public who may have the tendency to overlook diseases, partly because they are not charismatic. Important decisions that must be addressed, such as the regulation of coastal development or implementation of strategies to control land-based runoff must occur over large spatial scales and be in effect for long time periods. The science in support of these decisions is often conducted on much smaller spatial and temporal scales. Also, it may involve highly specialized research with results that are difficult to apply directly to policy decisions, or the science may be completely lacking. Given limitations of existing funding, coral disease research should be targeted towards efforts with the greatest management benefits that can be applied over large spatial scales, and research must also be relevant to understanding diseases and their impacts, and mitigating their effects.

ACKNOWLEDGMENTS

The National Ocean Service (NOS) convened a workshop in Charleston, South Carolina on 22-25 January 2002 as part of the Coral Disease and Health Consortium's (CDHC) efforts to develop a National Research Plan to improve our understanding of coral diseases. This paper was developed to provide an overview of what is known about diseases and to identify information needs of coral reef managers that could be provided through rapid ecological assessments, long-term monitoring, and targeted research. I am grateful to Dr. Cheryl Woodley, the head of CDHC, for the opportunity to present this information to the workshop participants. Comments on an early draft were provided by Dr. Tom Hourigan and Robin Bruckner. The final preparation of this report was greatly improved from reviews by Dr. Esther Peters, Dr. Deborah Santavy and Eric Borneman. Dr. Ernesto Weil and Dr. Debbie Santavy provided information on the monitoring protocols described in Appendix I, and Dr. Brian Keller and Billy Causey provided information on the management and science objectives for the FKNMS. Finally, I want to recognize the support of the National Marine Fisheries Service Office of Habitat Conservation and the Coral Reef Program in encouraging the preparation of this document, in supporting its publication, and in its distribution to those working on coral reef issues. The opinions and views expressed in this document are those of the author and do not necessarily reflect those of the National Marine Fisheries Service, National Oceanic and Atmospheric Administration or the U.S. government.

APPENDIX I: Monitoring and Assessment Approaches

I. Atlantic and Gulf Rapid Reef Assessment Protocol

The following description represents the benthic survey method for corals. The assessment technique involves 10m transects placed along depth gradients at haphazard, non-overlapping intervals. The condition of corals that touch the transect are assessed. The full protocol can be obtained at: <http://coral.aoml.noaa.gov/agra/method.htm>

The AGRRA method is focused on assessing the condition of the principal scleractinian and hydrozoan corals that contribute most to the three-dimensional structure and complexity of reefs. The vitality of these corals responsible for the construction and maintenance of reef framework is important for the long-term persistence of a coral reef. The AGRRA method assesses overall coral cover, and for each coral, assesses the amount of partial coral mortality (both recent and old), size and height, incidence of bleaching/diseases, causes of mortality, and number of damselfish.

AGRRA makes a further distinction of mortality into partial mortality that includes both “recently” dead and “old” dead. “Recently dead” is defined as any non-living parts of the coral in which the corallite structures are white and either still intact or covered over by a thin layer of filamentous algae or mud. In contrast, “old dead” is defined as any non-living parts of the coral in which the corallite structures are either gone or covered over by organisms that are not easily removed (e.g., certain algae and invertebrates), even if the outline of the dead corallites is still visible below (e.g., certain zooxanthella-containing boring sponges that overgrow their substrates).

Causes of partial mortality include, but are not limited to grazing (e.g., parrotfish bites), extended bleaching, sedimentation, algal overgrowth, disease (e.g., black band), and corallivores. To regenerate injured tissue, corals must reallocate energy that would otherwise be used for energy storage, growth or reproduction. It is fairly common for corals to experience partial mortality and to regenerate these areas if the lesions are not too large, relative to their remaining live tissues. Most partial mortality is small in size, but above certain sizes, which vary with the species and the kind of injury that has been received, dead skeletal areas tend to become overgrown or eroded by algae or other bioeroding organisms. In cases of large lesions, corals often will put energy into marginal growth instead of regenerating new tissue over the dead area.

By quantifying the amount of recent mortality, you sum the extent of any damage and obtain information on whether the corals are likely to recover. Several studies have shown that the extent and amount of partial mortality is related to differences in size, morphology, species identity, and differences in the spatial and temporal distribution of predators. Normally, small corals often have either no injuries or total mortality, while larger corals are likely to survive partial mortality. The AGRRA data can be used to calculate size frequency distributions as well as mortality patterns related to size and, when sample sizes are large enough, species identity. This allows the added benefit of being able to examine coral size-frequency statistics to assess a population structure and interpret levels of mortality. Although there are studies showing the importance of size-frequency distributions in population dynamics, additional research is required to determine “normal” size-frequency distributions for major reef-building species in various habitats.

Areas of recent mortality and adjacent coral tissues should be examined closely to determine if calices are disrupted or damaged (fish bites), if corallivores are present (e.g. gastropods or polychaetes), if polipary and coenostial structures are abnormally large or form aberrant masses (hyperplasm or

neoplasm), or if other disease conditions exist. Diseases are recorded based on color, including black (black band), white (white band or other white-type diseases in acroporids; white plague in massive corals), yellow (yellow blotch) or red (red band). White areas must be examined closely to distinguish areas without living tissue (e.g. exposed skeleton caused by disease or predation) from bleached tissue. Dark-spot disease also has a mottled appearance, however it is manifested as irregular patches that are darker than the normal pigmentation and these areas may be slightly depressed. Predation can often be differentiated from disease (and bleaching) by examining tissues at the interface of white, exposed skeleton; diseased tissue will appear irregular and may be sloughing off the skeleton. Elevated “chimneys” on *Acropora* spp. and “ridge mortality” on brain corals are induced by repeated three spot damselfish (*Stegastes planifrons*) bites.

1. At each SITE, haphazardly lay the 10-m transect line just above the reef surface in a direction that is parallel to the long axis of the reef. Make sure the line is taut.

Note: Be sure to avoid or cross any other transects that are being set by your companions, and stay away from the edges of the reef. Also try to avoid areas with abrupt changes in slope, deep grooves, large patches of sand or unconsolidated coral rubble. You want to place the transect in areas where corals are likely to grow. Unusual reef features should only be included to the extent appropriate to their relative abundance at the site.

2. Approximate live coral cover by swimming along the transect line with your 1 m measuring device and roughly estimating how many meters of the line is underlain by living coral (to the nearest 10 cm). Be sure to include all live stony corals below the line, regardless of size or species density (but remember do not include dead skeleton). Record this total. If the reefs are too small for you to avoid sandy patches, record how much of the transect line crosses sand, to allow a later calculation of the number of coral heads encountered/m of reef surface.

3. Swim towards the other end of the transect and stop at the first coral head, cluster, or thicket (or a portion) of the appropriate species that is located directly beneath the transect line, is at least 10 cm wide, and which is in original growth position. For a colony that has fallen or been knocked over, only assess it if it has become reattached to the substratum or is too large to move. If the colony is loose and rolling around, then skip it. For each coral surveyed, record each of the following:

A. Name (genus and species)

B. Record the water depth at the top of the corals at the beginning and end of each transect. In cases where bottom topography is very irregular, or the size of the individual corals is very variable, record the water depth at the top of each coral beneath the transect line at any major change in depth (>1m).

C. Identify the colony's boundaries based on connective or common skeleton, connected live tissue, polyp size, and polyp color. Using a measuring device, measure to the nearest cm, its maximum projected diameter (live + dead areas) in plan view and maximum height (live + dead areas) from the base of the colony' substratum, not the reef's substratum or ocean floor. The diameter should be measured perpendicular to the axis of growth. The height should be measured parallel to the axis of growth. Plan view is assessed from an angle that is parallel to the axis of growth.

Note 1: Some surveyors may want to measure all sizes of corals (even <10 cm max. diameter) on reefs on which large colonies are scarce. They are encouraged to do so, but only data based on corals greater than

10 cm will be directly comparable to other AGRRA results). In previous years we suggested measuring corals >25 cm, but based on consensual agreement we have changed it to >10cm.

D. Estimate the percent (%) of the coral that is "recently dead" and the % of the coral that is "**Long dead**" as viewed from above in "plan" or "map" view. Plan view is assessed from an angle that is parallel to the axis of growth (be prepared to tilt your head to find the axis of growth and establish the proper plan view).

"**Recently dead**" is defined as any non-living parts of the coral in which the corallite structures are white and either still intact or covered over by a layer of algae or fine mud. For recent mortality, there are several "stages" of recent

Very recent = white intact skeleton is still visible (dead w/in 1 month or less)

Recent = corallite may be covered by thin turf algae or sediment (up to 6 months)

Older recent = corallite structure may be slightly eroded or covered but can still identify to genera (< 1-2 yrs), unless covered by clionid sponge

E. Combine all of these recent mortality categories into RECENT. In some cases circular or oblong lesions or excavations caused by fish biting may result in destruction of the corallite. If fish bites are identifiable and constitute part of the mortality, consider it as recent mortality.

Note 1: "Long dead" is defined as any non-living parts of the coral in which the corallite structures are either gone or covered over by organisms that are not easily removed (certain algae and invertebrates). If it is entirely "long dead", indicate this on your data sheet as 100% "long dead", as long as you can identify it to generic level based on morphology (e.g., *Acropora palmata*) or skeleton (e.g., *Diploria* sp.).

Note 2: In some cases, a coral may be partially or completely overgrown by the brown encrusting sponge, *Cliona* sp. At a quick glance, it may look like live coral tissue, but if you look closely you may observe the in/ex-current holes of the sponge and sponge tissue and will not observe live coral polyps. If you can see the coral skeletal structure underneath and are able to identify it to genus (e.g., *Diploria* and *Montastraea*), this should be considered old mortality and you should note *Cliona* overgrowth in comments.

Note 3: If a coral like columnar *M. annularis* or *Dendrogyra* has been knocked over, has either reattached to the substrate or is too large to move, has started to regrow towards the water's surface, it now has a different diameter and height because of its new growth direction. You should measure diameter, height, and mortality in the correct plan view along the new axis of growth. Only measure the recently "live" part of the *M. annularis* when estimating mortality and not the old "base" because this will skew old mortality. Quickly scan over the surviving portions of the ENTIRE coral colony and note if there are any DISEASES and/or BLEACHED tissues present. Characterize any DISEASES by the following color categories:

- BB = Black band
- WB = White band
- WS = White spots, patches or pox
- WP = White plague
- YB = Yellow blotch (sometimes called yellow band)
- RB = Red band
- UK = Unknown

II. CORAL-OCTOCORAL DISEASE SURVEY - PROTOCOL LEVEL-2 CARICOMP. Modified by E.Weil June of 2000.

OBJECTIVES:

To assess the occurrence (incidence at population and community levels) and local/geographic distribution of coral and octocoral diseases across the wider Caribbean region.

More specifically : 1- To quantify relative frequencies of infected colonies in each coral species affected within and across localities. 2- To quantify relative frequencies of the different diseases within and across species and within and across localities. 3- To determine if there is a correlation between frequencies of diseased corals with depth, and distance to anthropogenic affected areas.

LEVEL- 1 - PROTOCOL:


Protocol will work with many different sampling methods (quadrats, chain transects, belt transects, etc.). This is to be carried out at the two reef sites of the CARICOMP reef monitoring using the chain. The survey is organized in a hierarchical way that will facilitate the decision-making process. A set of bilingual-underwater-laminated cards (Bruckner & Bruckner & Weil) was selected to be distributed amongst all site directors. These ID cards are to be used as the ONLY guide to identify the different diseases and other causes of coral mortality. This will serve as a tool to standardize the identification of the different diseases and causes of mortality. There are ten, 10 m long chain sections per locality. Survey should be conducted over a belt transect of two meters wide (one meter on each side of the chain) by ten meter long (20 m²), for a total of 200 m² per locality at one depth interval. All coral and octocoral colonies within this area should be checked and counted. The sample unit is one colony (ramet). Each coral colony should be identified to species (or genus at least) and/or to either one of the five colony form descriptors specified in the CARICOMP manual (branching, crustose, massive, foliose or milleporid). Octocorals should be identified at least at the genera level, or using the four categories (rod, fan, feather, whip) in the manual.

The first discrimination-category for each colony is whether it is healthy or unhealthy. Healthy colonies are those without signs of disease, recent injury and/or bleaching. Unhealthy colonies are those with any signs of disease, recent, open injuries and/or bleaching. A non-healthy colony will then be put into one of three categories, bleached, injured or diseased. If a colony has a disease and any of the other two signs (bleaching, injuries) it will be included in the disease category. Notes about the other signs should be recorded. Bleaching should be verified by looking for the presence of live tissue (but decolorated) and at the pattern of bleaching (usually not a band-like pattern like most “white-type” diseases; e.g., **pale, blotchy or white**). Injured colonies include those with damage from predation (often with skeletal damage), anchors or other physical damage, and overgrowth. If you can identify the type of damage (small lesions like individual bite marks or large excavations such as anchor damage), please do so. If the colony is being overgrown by other organisms (algae, zoanthid, sponge, hydroid, tunicates, etc.) it is important to note which organism (s) are overgrowing it. Colonies with old dead areas covered with algae but with healthy looking tissues fall into the healthy category.

A diseased colony will then be carefully observed to identify the disease that is affecting it. Observations should be compared with the information in the ID-cards to put the colony within any of the following 9 categories (see below and the flow diagram):

- 1- Black-Band Disease (BBD)
- 2- White-Band Disease (WBD)
- 3- White Plague-II (WP-II)
- 4- Yellow-Blotch Disease (YBD)
other favid species.
- 5- Dark Spots Disease I (DS-I)
- 6- Dark Spots Disease II (DS-II)
- 7- Red Band Disease (RBD)
been reported for
- 8- Aspergillosis (ASP)
- 9- White POX (WPX)
- 9- Other -

Multispecific
Types I and II (only in acroporids for the moment)
Multispecific
Only described in species of *Montastraea* but also reported in other favid species.
Small, dark areas with no apparent tissue mortality. Common in *Siderastrea spp.*
Large dark areas, larger than the DSD-I; common in *M. annularis* and *S. intersepta*.
Careful here since BBD can be seen as red bands too. RBD has been reported for *Gorgonia spp* and Agaricids in the Caribbean.
Gorgonia ventalina, *G. flabellum*, *P. americana*; other octocorals (*Plexaura flexuosa*).
Round or irregular white areas on surface of Acroporids. this category includes all other Aunconfirmed-pathogen-produced diseases (tumors, hyperplasia, and all the blotches).



LEVEL 2 PROTOCOL - CORAL DISEASES .

The purpose of this activity is to assess the incidence of diseases in each coral and octocoral species at different depth intervals (habitats) and in at least two different reef sites per locality. The diseases will be identified following the protocol level-1 described above. The field experimental design is as follows :

1- At least two separated (more than 5 km) reef sites should be selected at each geographic locality (CARICOMP chapter), and preferably, one near and one far from anthropogenic-impacted areas. A preliminary qualitative, visual assessment of current levels of disease incidence and depth distribution within the two different reef localities should be done to determine if coral diseases are common or rare.

2- A quantitative assessment will follow using a modified protocol from the standard CARICOMP coral survey. Three depth intervals would be selected in those reefs with a depth profile to at least 45 feet deep. The depth intervals are: 0-15 feet; 16-35 feet and 35-50 feet, so major habitats would be surveyed (shallow, branching *Acropora* zone and boulder-crustose platform zone; intermediate depth *Montastraea* zone - massive, boulders and platy spp.; and the deeper more diverse zone - foliose, massive and platy spp). Re-bars or other permanent means of marking the transect position should be hammered into the substrate for future monitoring.

In each depth interval, at least three 20 meter long x 2 m (40 m²) wide band-transects should be surveyed (a minimum of 9 transects/reef, better if 12 transects per reef are surveyed). Two sets of 6 band-transect each should be separated at least 100 m. In each depth interval a set of two band-transects could be done using a 50 m long, plastic tape. Once the tape is positioned, two 20 x 2 m bands separated by a 10 m interval will be surveyed (from 0 to 20 m and 30 to 50 m). A one m long pvc section, marked in cm (100 cm) will be moved along each side of the transect line and every colony of coral and of *Gorgonia* spp. will be surveyed using the protocol level 1 described above.

Once a colony is found to have a disease, two diameters (maximum and minimum) will be measured with the pvc stick to estimate surface area. **ONLY diseased colonies (corals and *Gorgonia* spp) will be measured.**

Separation between the consecutive band transects (at same depth interval) should be at least 10 m. If the reef areas do not have a depth profile or slope, or are such that only two depth intervals can be selected, the minimum of 12 band-transects should be then spatially distributed within the area. All coral colonies should be recorded to species (or genus at least) level.

Confusion of what an individual colony or ramet is will probably arise when branching, columnar or massive species that have suffered fission (fragmentation and/or partial mortality) are encountered. For the purposes of this protocol, a ramet is any colony that is spatially separated from any other colonies of the same species. On some occasions however, it is clear for some fragmented colonies, that in the near past, there was a physiological and physical connection between two or more ramets (i.e. in massive colonies suffering partial mortality or broken branches nearby the mother ramet of ramose species). In these situations, instead of counting 2000 small fragments (i.e. *P. porites* or *M. mirabilis*) one should use common sense and spatially delimit the spread of the individual genet (in the many ramets) and count it as a single ramet (genet).

The study is aimed at comparing the incidence of coral diseases between different reefs within a given location (Cuba, Panama, Venezuela, etc.) and across locations in the wider Caribbean. It is fundamental to the proposed experimental design to insure the sampling of large reef areas, in order to estimate

current levels of disease incidence at a suitable spatial scale. Time window to conduct the surveys is between July and October. Frequency of surveys will depend on each individual laboratory's logistics, time and personnel. A template (Lotus 123, Quatro or excel) to input data into the computer is available from E.Weil. If you decide to carry on this protocol, please contact Dr. E. Weil at eweil@caribe.net.

Materials and Methods provided by D. Santavy (Santavy et al., 2001) for coral disease surveys conducted in the Florida Keys.

General Approach

The SCUBA-based field study was developed and implemented over a two-year period in the south Florida region. Survey areas were selected in the Lower Florida Keys in the vicinity of Key West, the New Grounds, and the Dry Tortugas (Fig. 1). Zones that contained hard coral bottom were demarcated within each geographic area. These coral reef zones were located using a prototype of the Florida Marine Research Institute (FMRI) Benthic Habitats Map of the Florida Keys. Potential stations for the pilot surveys were selected using a stratified random design, within the three regional areas. Individual stations were chosen by placing a random grid pattern that incorporated a hexagonal overlay over the individual coral reef zones contained within each geographic area. Surveyors went to randomly selected locations and assessed their suitability for sampling. If the location had sufficient coral coverage (>5%), the site was surveyed; if it was not suitable, the next location on the list was assessed for sampling suitability. Twenty-one stations were surveyed in the 1997 spring pilot survey.

Survey periods targeted late spring, the time when coral diseases are believed to emerge, and late summer, the time when coral diseases are believed to be most prevalent. The 1997 spring and summer pilot surveys were conducted during 1 to 8 June 1997 and 6 to 14 September 1997. Stations assessed as suitable during the spring pilot could not be permanently established at that time, but permission was granted for permanent installation of stakes during the summer pilot. In September 1997, stations assessed during the spring pilot were relocated using GPS coordinates, and permanent sites were established by installing stakes to be used for future surveys in 20 of these stations (1 being omitted at New Grounds). At the same time, 6 new stations were added in the Key West and Dry Tortugas areas.

Survey strategy

Based on results from the 1997 pilot survey, reef type was added as a stratum to the 1998 sampling design (see Results). For the 1998 surveys, 6 additional stations were selected to balance the sample design across two strata: (1) three geographic areas established in the pilot surveys and (2) three reef types. The three reef types—fore reef, back reef, and transitional reef, as defined in the Florida Keys National Marine Sanctuary (FKNMS) Management Document—were used for comparison among the areas. However, not all areas contained all reef types; for example, only deep transitional reefs were found in the New Grounds area. The 1998 spring survey was conducted from 25 May to 1 June 1998 and the 1998 summer survey from 2 to 11 September 1998. Thirty-two stations were assessed in the spring and summer of 1998. A power analysis was performed on the data acquired from both spring and summer 1998 surveys to determine the optimal number of sampling stations and the appropriate α level for data analysis.

Survey methodology

All surveys were conducted using a radial arc transect method developed for this study. SCUBA was used on deeper reefs and snorkel was used on shallow back reefs. A stainless steel rod was positioned by driving it into the calcareous substratum for temporary sites (1997 spring pilot) or by permanently affixing a 12" stainless steel pipe (all other surveys) at the designated site with underwater epoxy (Gunnebo® Liquid Roc 500, Gunnebo Fastening Corporation 800-336-1640). Site coordinates were determined by GPS technology during 1997, then by Differential-GPS (D-GPS) during 1998 surveys as it became available in our study region, improving our ability to easily locate stations. Surface maps with triangulation bearings and maps of underwater structures were generated, and a subsurface 3" buoy was used to mark each station to enable us to return to each underwater stake.


The survey procedure required inserting a 6-foot pole into the stainless steel pipe. The pole had 2 adjustable collets with a carabineer and a snap shackle on one end of the pole. A 12-m Kevlar™

fishing line contained within a plastic housing reel (fly-fishing reel) was fastened to the snap shackle on the pole. The line was marked every meter and unreeled to desired lengths during the survey. Small fluorescent tags attached to the line were used to mark the 2-m wide transect area under the line. A line tender held the line taut above the reef structures and slowly moved the line in an arc around the fixed central stake, allowing time for the surveyors to record their data (Fig 2). Two surveyors swam in concentric circles directly over the line, one recording the number of colonies of each coral species and the other recording the number of colonies of each species that displayed signs of a specific disease. The surveyors counted colonies larger than 10 cm that fell directly below each 2-m segment of the line, providing more than half of their area occurred within the segment. The originating point of the arc was marked with a weighted subsurface buoy to alert the line tender when an entire arc segment had been completed.

During 1997, the surveys were conducted within the entire 10-m radius, in multiple radial increments that enlarged the arc by 2 m for each complete circle. For example, the first arc segment included the 0- to 2-m increment, the second arc segment included the 2- to 4-m segment, and so on until the entire 10-m radius of the arc had been completed, circumscribing a total area of 314m². A species area curve was constructed to compare the June 1997 mean percent coral disease (MPCD) for increasing increments of the radial arc areas within the arc transect. Analysis of variance was used to determine whether the MPCD could be estimated using a portion of the arc, instead of the entire 10-m radius. Individual MPCD for each arc segment (0-2, then 2-4, and so on for each segment of 10-m radius) and cumulative MPCD for the cumulative areas of the arc segments (0-2, then 0-4, and so on for the entire 10-m) were used for the analysis, with arc area used as the class variable. This investigation determined that an area of 113m², within the 8- to 10-m segment was sufficient for a reliable estimate of total MPCD (see Results). Therefore, during 1998 surveys only the 8- to 10-m segment of the arc radius was used for assessment.

Coral species and coral disease identification

Ten disease conditions affecting 16 species of scleractinian corals and gorgonian sea fans were enumerated (Table 1). Three species of coral contained within the *Montastraea annularis* complex were combined as a single category for data analysis, because discrepancies in identification were noted among some of the species, *Gorgonia ventalina*, were combined as *Gorgonia* spp. All diseases scored if mortality had occurred recently and the signs used to have been detailed conditions described in disease (Bruckner & Holden, 1996), might be the same disease. We could not distinguish between the two conditions based on Bruckner & Bruckner's (1997) mention of patchy necrosis or Holden's (1996) mention of white pox; therefore we used the term patchy necrosis disease/white pox to describe the lesions found on *Acropora palmata* colonies that were not white-band disease or predation. We did not distinguish the differences between white plague type 1 and 2, since the primary difference in distinguishing them in the literature is dependent on the rate of progression (Dustan, 1977; Richardson et al., 1998a, 1998b). This could not be determined in our surveys; therefore, we identified these conditions only as white plague in our surveys. Finally, there is some uncertainty in identifying aspergillosis (Kim et al., 1997; Nagelkerken et al., 1997a, 1997b; Smith et al., 1996). For our surveys, this disease was scored if one of the following



conditions were met: white fungal-like filaments with active lesions (tissue loss) and major skeletal damage, or white fungal-like filaments with active lesions showing coenenchyme purpling, or white fungal-like filaments with active lesions and purple galls in the vicinity of the diseased area. There are some inconsistencies in the literature concerning the signs of this disease, therefore, one might prefer to refer to the condition simply as sea fan disease.

Quality assurance

A rigorous quality assurance (QA) plan was adopted to quantify surveyor error and minimize data processing errors. The data collection protocols required training to improve identification skills and familiarize participants with the survey procedures, including the completion of standardized data forms. Scientific experts assessed the coral species and diseases in the 1997 surveys. Coral surveyors in 1998 were evaluated by expert coral taxonomists and expert coral pathologists for their ability to identify coral species and to classify coral conditions as either healthy, affected by a specific disease, bleached, or physically damaged. Only those individuals who had successfully passed a test (scoring 90% similarity or greater using the experts as truth) were employed to collect data for the surveys. To evaluate inter-surveyor error, we had multiple surveyors take repeated counts of coral colonies by species and of coral disease types by species. To evaluate intra-surveyor error, we had each surveyor count species and disease types multiple times at a single station. The QA plan included procedures for several levels of data verification, including checks made in the field, duplicate surveys, and independent validation of all electronically entered data.

Data analysis

Data were recorded on standardized data sheets photocopied on Dura/Copy™ (J.L. Darling Corp., Tacoma, WA.) underwater paper. All data were entered into a computerized database using a PerForm Pro™-generated template, exported into MS Excel® worksheets, and used to create SAS data sets. Electronic data quality was confirmed twice by someone other than the original data recorder. The parameter of interest was mean percent coral disease (MPCD) (i.e., number of affected colonies per total number of colonies) per unit area. Data were analyzed using one-way ANOVA for unbalanced design in the 1997 pilot surveys and for balanced design in the 1998 surveys. The assumptions for ANOVA were tested and met, including independence, homogeneity of variances, and normality. The class variables or strata used included geographical areas (Key West, New Grounds, or Dry Tortugas) and reef types (back, fore, or transitional reef). Statistical significance for type I error was designated as $\alpha=0.05$ level. Tukey's Studentized Range Test (HSD) was used for means separation at the $\alpha =0.05$ level. A power analysis was performed to determine appropriate type 1 and 2 errors, and statistical power to be employed for future analyses. All analyses for the study were performed using SAS Version 6.12 (Statistical Analysis System Institute Inc., Cary, N.C., 1989-1996).

Appendix II. Priorities for research on coral diseases linked to management objectives in Florida. Developed by Brian Keller and Billy Causey.

1. Direct and indirect causes of coral decline in the Florida Keys, with emphasis on cause and effect. Process-oriented research is required to separate natural variation in community composition from anthropogenic changes. We know that disease contributes to the decline, but we do not know its relative importance and how much forces such as global warming are adding to the problem.
2. Effects of coral bleaching on coral survival. There is a general relationship between severity of bleaching events and coral mortality. However, more information is needed on intra- and interspecific differences in sensitivity to bleaching, habitat-related differences, and synergistic effects, including possible genetic factors.
3. Coral pathology and epizootiology with an emphasis on cause and effect. More information is needed on coral immune systems and how they may be compromised by environmental stresses, whether a number of purported coral diseases meet Koch's criteria, how diseases spread, and whether there are ways to prevent coral diseases.
4. Relationships between water quality and the incidence of diseases. There are claims that elevated nutrient concentrations are associated with black-band disease.

Coral Disease Literature

- Abbott, R.E. 1979. Ecological processes affecting the reef coral populations at the East Flower Garden Bank, northwest Gulf of Mexico. Ph. D. Dissertation. Texas A&M Univ., College Station, TX. 154 pp.
- Aeby, G.S. 1992. The potential effect the ability of a coral intermediate host to regenerate has had on the evolution of its association with a marine parasite. Proc. Seventh Intern. Coral Reef Symp. 2:809-815.
- Aeby, G.S. 1998. A digenean metacercaria from the reef coral, *Porites compressa*, experimentally identified as *Podocotyloides stenometra*. J. Parasitology. 84(6):1259-1261.
- AGRRA. 1998. Atlantic and Gulf Rapid Reef Assessment Protocol. Available on line at: <http://coral.aoml.noaa.gov/agra/method.htm>
- Alker A.P., G.W. Smith and K. Kim (in press) Characterization of *Aspergillus sydowii* (Thom et Church), a fungal pathogen of Caribbean sea fan corals. Hydrobiologia.
- Anonymous. 1996. Coral disease hot spots in the Florida Keys. Science. 274:2017.
- Anonymous. 1997. New Caribbean coral killer. Science. 276:1979.
- Antonius, A. 1973. New observations on coral destruction in reefs. Assoc. Isl. Mar. Lab. Carib. Abstract. 10:3.
- Antonius, A. 1977. Coral mortality in reefs: A problem for science and management. Proc. Third Intern. Coral Reef Symp. 1:617-623.
- Antonius, A. 1981a. Coral Reef Pathology: A review. Proc. Fourth Intern. Coral Reef Symp. 2:3-6.
- Antonius, A. 1981b. The "band" diseases in coral reefs. Proc. Fourth Intern. Coral Reef Symp. 2:7-14.
- Antonius, A. 1985a. Coral diseases in the Indo-Pacific: A first record. Pubblicazioni Staz. Zool. Napoli [(PSZNI), Mar. Ecol.], 6:197-218.
- Antonius, A. 1985b. Black band disease infection experiments on hexacorals and octacorals. Proc. Fifth Intern. Coral Reef Symp. 6:155-160.
- Antonius, A. 1987. Survey of Red Sea coral reef health. 1- Jeddah to Jizan. Proc. Saudi Biol. Soc. 10:149-163.
- Antonius, A. 1988a. Black band disease behavior on Red Sea reef corals. Proc. Sixth Intern. Coral Reef Symp. 3:145-150.
- Antonius, A. 1988b. Distribution and dynamics of coral disease in the eastern Red Sea. Proc. Sixth Intern. Coral Reef Symp. 2:293-298.
- Antonius, A. 1993. Coral reef health in Mauritius. International Society for Reef Studies. First European Regional Meeting. Dec 16-20.
- Antonius, A. 1995a. Coral diseases as indicators of reef health: Field methods. Proc. Sec. Europ. Meet. I.S.R.S., Publ. Serv. Geol. Lux. 29:231-235.
- Antonius, A. 1995b. Pathologic syndromes on reef corals: a review. Proc. Sec. Europ. Meet. I.S.R.S., Publ. Serv. Geol. Lux. 29:161-169.

- Antonius, A. 1999. *Halofolliculina corallasia*, a new coral-killing ciliate on Indo-Pacific reefs. *Reef Sites. Coral Reefs*. 18:300.
- Antonius, A. and A. Weiner. 1982. Coral reefs under fire. *PSZNI (Mar. Ecol.)*. 3:225-277.
- Antonius, A. and B. Riegl. 1997. A possible link between coral diseases and a corallivorous snail (*Drupella cornus*) outbreak in the Red Sea. *Atoll Res. Bull.* 447:1-9.
- Antonius, A. and B. Riegl. 1998. Coral diseases and *Drupella cornus* invasion in the Red Sea. *Reef Sites. Coral Reefs*. 447:1-9.
- Antonius, A. and E. Ballesteros. 1998. Coral reef health problems in the Caribbean.. *Revista de Biología Tropical* 46 (Suppl. 5):
- Antonius, A. and J. Afonso-Carillo. 2001. *Pneophyllum conicum* killing reef-corals in Mauritius: A new Indo-Pacific syndrome? Proceedings of the Intl. Conf. on Scientific Aspects of Coral Reef Assessment, Monitoring, and Restoration, Rosenstiel School of Marine and Atmospheric Science. *Bull. Mar. Sci.* 69: 613-618.
- Aronson, R.B. and W.F. Precht. 1997. Stasis, biological disturbance, and community structure of a Holocene coral reef. *Paleobiology*. 23(3):326-346.
- Aronson, R.B. and W.F. Precht. 2001. Evolutionary palaeoecology of Caribbean coral reefs. In: W.D. Allmon and D.J. Bottjer (Eds). *Evolutionary paleoecology: The ecological context of macroevolutionary change*. Columbia University Press, NY, pp. 171-233.
- Aronson, R.B., W.F. Precht and I.G. Macintyre. 1998. Extrinsic control of species replacement on a Holocene reef in Belize: the role of coral disease. *Coral Reefs*. 17:223-230.
- Australian Institute of Marine Sciences (AIMS). 2000. Status of coral reefs of the world: 2000. Edited by Clive Wilkinson. AIMS, Western Australia. 363 pp.
- Bak, R.P.M. 1983. Neoplasia, regeneration and growth in the reef-building coral *Acropora palmata*. *Mar. Biol.* 77:221-227.
- Bak, R.P.M. and S.R. Criens. 1981. Survival after fragmentation of the colonies of *Madracis mirabilis*, *Acropora palmata*, and *A. cervicornis* (Scleractinia) and the subsequent impact of a coral disease. *Proc. Fourth Intern. Coral Reef Symp.* 2:221-228.
- Banin E., Y. Ben-Haim, M. Fine, T. Israely, and E. Rosenberg. (in press). Virulence mechanisms of the coral bleaching pathogen *Vibrio shiloi*. *Proc 9th Int Coral Reef Symp*
- Banin, E., T. Israely, A. Kushmaro, Y. Loya, E. Orr, and E. Rosenberg. 2000. Penetration of the coral-bleaching bacterium *Vibrio shiloi* into *Oculina patagonica*. *Appl. Environ. Microbiol.* 66(7):
- Borenman, E. 2001. Coral Health. In: E. Borenman, *Aquarium Corals*. pp. 360-385.
- Briggs, K.M., D.L. Chau, R.B. Jonas, and D.J. Ellis. 1999. Photosynthetic, anoxygenic sulfur oxidizing bacteria in black-band disease on boulder coral, Andros Island, the Bahamas. *Intern. Conf. on Scientific Aspects of Coral Reef Assessment, Monitoring, and Restoration*. National Coral Reef Institute, Florida. Abstract 61.
- Bruckner, A.W. 1999. Black-band disease (BBD) of scleractinian corals: occurrence, impacts and mitigation. Ph.D. Thesis, UMI Dissertation Services, 286 pp.

- Bruckner, A.W. 2000a. The Global Coral Disease Database. Abstract. 25th Eastern Fish Health Workshop. p 54.
- Bruckner, A.W. 2000b. The incidence and effect of yellow-blotch disease in the Southern Caribbean. Abstract. 25th Eastern Fish Health Workshop. p. 51.
- Bruckner, A.W. 2001. Coral health and mortality. Recognizing signs of coral diseases and predators. In: Humann and Deloach, *Reef Coral Identification: Florida Caribbean Bahamas*. New World Publications, Inc. Jacksonville, FL pp. 240-271.
- Bruckner, A.W. and R.J. Bruckner. 1997a. The persistence of black-band disease in Jamaica: Impact on Community Structure. Proc. Eighth Intern. Coral Reef Symp. 1:601-606.
- Bruckner, A.W. and R.J. Bruckner 1997b. Outbreak of coral disease in Puerto Rico. *Reef Sites*. Coral Reefs. 16:260.
- Bruckner, A.W. and R.J. Bruckner 1997c. Emerging infections on the reefs. *Natural History*. 12/97-1/98: 48.
- Bruckner, A.W. and R.J. Bruckner. 1998a. Destruction of coral by *Sparisoma viride*. *Reef Sites*. Coral Reefs. 17:350.
- Bruckner, A.W. and R.J. Bruckner. 1998b. Rapid wasting disease: pathogen or predator? *Science*. 279. 2023-2025.
- Bruckner, A.W. and R.J. Bruckner. 1998c. Rapid-Wasting "Disease": Coral predation by stoplight parrotfish. *Reef Encounters*. 23:18-22.
- Bruckner, A.W. and R.J. Bruckner. 1998d. Treating coral disease. *Coastlines*. 8.3:10-11.
- Bruckner, A.W. and R.J. Bruckner. 1999. Rapid assessment of coral reef condition and short-term changes to corals affected by disease in Curaçao, Netherlands Antilles. Intern. Conf. on Scientific Aspects of Coral Reef Assessment, Monitoring, and Restoration. National Coral Reef Institute, Florida. Abstract 62.
- Bruckner, A.W. and R.J. Bruckner. 2000. The prevalence of coral diseases on reefs surrounding Mona Island. Abstract. Ninth Intern. Coral Reef Symp., Bali Indonesia. p. 281.
- Bruckner, A.W. and R.J. Bruckner. *In press*. Coral predation by *Sparisoma viride* and lack of relationship with coral disease. Proc. Ninth Intern. Coral Reef Symp.
- Bruckner, A.W. and R.J. Bruckner. *In press*. Condition of coral reefs off less developed coastlines of Curaçao (*stony corals and algae*). *Atoll Research Bull.*
- Bruckner A.W., R.J. Bruckner and E.H. Williams Jr. 1997a. Spread of a black-band disease epizootic through the coral reef system in St. Ann's Bay, Jamaica. *Bull. Mar. Sci.* 61:918-928.
- Bruckner, R.J. A.W. Bruckner and E.H. Williams, Jr. 1997b. Life history strategies of *Coralliophila abbreviata* Lamarck (Gastropoda:Coralliophilidae) on the southwest coast of Puerto Rico. Proc. Eighth Intern. Coral Reef Symp 1:627-632.
- Bruckner, A.W., R.J. Bruckner and P. Sollins. 2000. Parrotfish predation on live coral: "spot biting" and "focused biting". *Reef Sites*. Coral Reefs. 19:50.
- Bruckner, R.J. and A.W. Bruckner. 1998c. First aid for fragile coral reefs. *Currents*. Summer 1998:5-9.
- Bush, W.S. 1986. Effects and extent of white-band disease on coral reefs surrounding Lee Stocking Island, Bahamas. Class project, ZOOL 677, Dr. M. Reaka, University of Maryland. pp. 1-21.

- Bythell, J. C. & Bythell, M. 1992. Benthic reef community dynamics at selected sites at Buck Island Reef National Monument, 1988 - 1991. Monitoring fixed linear transects using the chain transect technique. Chapter 1. Ecological studies of Buck Island Reef National Monument, St. Croix, U.S. Virgin Islands: a quantitative assessment of selected components of the coral reef ecosystem and establishment of long term monitoring sites. Part II. NPS Coral Reef Assessment Program: pp. 1 - 24.
- Bythell, J.C., E.H. Gladfelter and M. Bythell. 1993. Chronic and catastrophic natural mortality of three common Caribbean reef corals. *Coral Reefs*. 12:143-152.
- Bythell, J.C. and Sheppard, C. 1993. Mass mortality of Caribbean shallow corals. *Mar. Pol. Bull.* 26:296-297.
- Carlton, R.G. and L.L. Richardson. 1995. Oxygen and sulfide dynamics in a horizontally migrating cyanobacterial mat: black band disease of corals. *Federation of European Microbiological Societies Microbiology Ecology*. 18:155-162.
- Cervino, J.M. 1998. New diseases in coral reefs. *Endangered oceans: Threats to human Health*. Center for Health and the Global Environment. Harvard Medical School. Dec. 7 1998.
- Cervino, J., T. Goreau, G. Smith, K. DeMeyer, I. Nagelkerken and R. Hayes. 1997. Fast spreading new Caribbean coral disease. *Reef Encounter*. 22:16-18.
- Cervino, J., T.J. Goreau, T.J., R. Hayes, I. Nagelkerken, K. Patterson, J.W. Porter, G.W. Smith, and C. Quirolo. 1998. Coral disease. *Science*. 280:499-500.
- Cervino, J.M. and G. Smith. 1997. Corals in peril. *Ocean Realm* 42:33-35.
- Cervino, J.M. and G. Smith. 1997. Coral diseases are on the increase. *Earth Times News Service*. 8/31/97
- Cervino, J.M, G.W. Smith, T.J. Goreau, R. Hayes, S. Polson and S. James. 2000. Decreased zooxanthellae densities and mitotic indices in corals affected by yellow band, dark spot, white band and white pox syndromes. Abstract, Ninth International Coral Reef Symposium, Bali, Indonesia. p.
- Cervino, J, T. J. Goreau, I. Nagelkerken, G. W. Smith, and R. Hayes. 2001. Yellow band and dark spot syndromes in Caribbean corals: Distribution, rate of spread, cytology, and effects on abundance and division rate of zooxanthellae. *Hydrobiologia*. 460:53-63.
- Cheney, D.P. 1975. Hard tissue tumors of scleractinian corals. *Advances in Experimental Medicine and Biology*. 64:77-87.
- Cheng, T.C. and A.K.L. Wong. 1974. Chemical, histochemical, and histopathological studies on corals, *Porites* spp., parasitized by trematode metacercaria. *J. Invertebr. Pathol.* 23:303-317.
- Chesher, 1984. Practical problems in coral reef utilization and management: A Tongan case study. *Proc. Third Intern Coral Reef Symp.* 214-217.
- Coles, S.L. 1994. Extensive disease outbreak at Fahl Island, Gulf of Oman, Indian Ocean. *Reef Sites*. *Coral Reefs*. 13:242
- Coles, S.L., and D.G. Seapy. 1998. Ultra-violet absorbing compounds and tumorous growths on a croporid corals from Bandar Khayran, Gulf of Oman, Indian Ocean. *Coral Reefs*. 17:195-198.
- Davis, M, E. Gladfelter, H. Lund, and M. Anderson. 1986. Geographic range and research plan for monitoring white band disease. *Virgin Islands Resource Management Cooperative*. Biosphere Reserve Res. Rep. 6. pp 28.

- Diáz, J.M., J. Garzón-Ferreira and S. Zea. 1992. Evaluación del estado actual del arrecife coralino de la Isla de San Andrés. Final Rep. Project INVEMAR/CORPES, Santa Marta. 147 pp.
- Dinsdale, E.A. 1994. Coral disease on the Great Barrier Reef. Joint Scientific Conference on Science, Management and Sustainability of Marine Habitats in the 21st Century. Abstract. Australian Marine Sciences Association, Townsville, Australia. p. 25.
- Dinsdale, E.A.. In press. Abundance of black-band disease on corals from one location on the Great Barrier reef: a comparison with abundance in the Caribbean region. Proc. Ninth Intern. Coral Reef Symp.
- Downs, C.A., E. Mueller, S. Phillips, J.E. Fauth, and C.M. Woodley. 2000. A molecular biomarker system for assessing the health of coral (*Montastraea faveolata*) during heat stress. Mar. Biotechnol. 2:533-544.
- Dube, D., K. Kim, A.P. Parker and C. D. Harvell. 2002. Size structure and geographic variation in chemical resistance of sea fan corals *Gorgonia ventalina* to a fungal pathogen. Mar. Ecol. Prog. Ser. 231:139-150.
- Ducklow, H.W. and R. Mitchell. 1979. Observations on naturally and artificially diseased tropical corals: scanning electron microscope study. Microbial Ecology. 5:215-223.
- Ducklow, H.W. and R. Mitchell. 1979b. Bacterial populations and adaptations in the mucus layers on living corals. Limnol. Oceanography. 24:715-725.
- Dustan, P. 1977. Vitality of reef coral populations off Key Largo, Florida: recruitment and mortality. Env. Geol. 2:51-58.
- Dustan, P. 1987. Preliminary observations on the vitality of reef corals in San Salvador, Bahamas. In: Curran, H.A. (ed). Proc. Third Symp. on Geol. of the Bahamas, Fort Lauderdale, FL. pp. 57-65.
- Dustan, P. 1993. Developing methods for assessing coral reef vitality: A tale of two scales. Proc. Colloquium on Global Aspects of Coral Reefs: Health, Hazards and History. University of Miami, RSMAS. pp. 38-44.
- Dustan, P. 1999. Coral reefs under stress: sources of mortality in the Florida Keys. Natural Resources Forum. 23:147-155.
- Dustan, P. and J.C. Halas. 1987. Changes in the reef-coral community of Carysfort Reef, Key Largo, Florida: 1974 to 1982. Coral Reefs. 6:91-106.
- Edmunds, P.J. 1991. Extent and effect of black band disease on a Caribbean reef. Coral Reefs. 10:161-165.
- Edmunds, P.J. 2001. Recruitment of scleractinians onto the skeletons of corals killed by black band disease. Coral Reefs. 19:69-74.
- Epstein, P.R. et al. 1998. Marine ecosystems: Emerging diseases as indicators of change: Health of the oceans from Labrador to Venezuela. Year of the Oceans Special Report. Cent. Health Global Environ, Harvard Medical School. 85 pp.
- Fabricius, K. 1999. Tissue loss and mortality in soft corals following mass-bleaching. Reef Sites. Coral Reefs. 18:54.
- Fenner, 1998. Coral-list posting on the occurrence of BBD. March. p. 18.
- Feingold, J.S. 1988. Ecological studies of a cyanobacterial infection on the Caribbean Sea plume *Pseudopterigorgia acerosa* (Coelenterata: Octocorallia). Proc. Sixth Intern. Coral Reef Symp. 3:157-162.

- Feingold, J.S. and L.L. Richardson. 1999. Impact of plague type II disease on populations of *Dichocoenia stokesii* in southeast Florida. Intern. Conf. on Scientific Aspects of Coral Reef Assessment, Monitoring, and Restoration. National Coral Reef Institute, Florida. Abstract 62.
- Franklin, E.C. 1998. Battling a coral disease in the Florida Keys National Marine Sanctuary. Ocean Resources Foundation. Winter 98-99:4.
- Frias-Lopez, J., A.L. Zerkle, G.T. Bonheyo, and B.W. Fouke. 2002. Partitioning of bacterial communities between seawater and healthy, black band diseased, and dead coral surfaces. Applied and Environmental Microbiology. 68(5): 2214-2228.
- Garrett, P. and H. Ducklow. 1975. Coral Diseases in Bermuda. Nature. 253:349-350.
- Garzón-Ferreira, J. and M. Kielman. 1993. Extensive mortality of corals in the Columbian Caribbean during the last two decades. Global Aspects of Coral Health. 247-253.
- Garzón-Ferreira, J. and S. Zea. 1992. A mass mortality of *Gorgonia ventalina* (Cnidaria:Gorgoniidae) in the Santa Marta area. Caribbean Coast of Columbia. Bull. Mar. Sci. 50:522-526.
- Geiser, D.M., J.W. Taylor, K.B. Ritchie, and G.W. Smith. 1998. Cause of sea fan death in the West Indies. Nature. 394(6689):137-138.
- Gil-Agudelo, D.L. and J. Garzón-Ferreira. 1999. Seasonal variation of dark spots disease in the Columbian Caribbean. Intl. Conf. on Scientific Aspects of Coral Reef Assessment, Monitoring, and Restoration. Ft. Lauderdale, Fl. Abstract 91.
- Gil-Agudelo, D.L. and J. Garzón-Ferreira. 2001. Spatial and seasonal variation of dark spots disease in coral communities of the Santa Marta area (Columbian Caribbean).Proceedings of the Intl. Conf. on Scientific Aspects of Coral Reef Assessment, Monitoring, and Restoration. Bull. Mar. Sci. 69:619-629.
- Gladfelter, W.B. 1982. White-band disease in *Acropora palmata*: implications for the structure and growth of shallow reefs. Bull. Mar. Sci. 32: 639-643.
- Gladfelter, W.B. 1991. Population structure of *Acropora palmata* on the windward fore reef, Buck Island National Monument, St. Croix, U.S. Virgin Islands. U.S. Department of the Interior, National Park Service, U.S. Virgin Islands. 172 pp.
- Gladfelter, W.B., E.H. Gladfelter, R.K. Monahan, J.C. Ogden, and R.F. Dill. 1977. Coral destruction, environmental studies of Buck Island Reef National Monument. Prepared for the US Nat Park Serv. U.S. Department of Interior. Biosphere Reserve Research Report No. 6.
- Glazebrook, J.S. and H.M. Streiner. 1994. Pathology associated with tumours and black band disease in corals from Agincourt Reef. Joint Scientific Conference on Science, Management and Sustainability of Marine Habitats in the 21st Century. Austr. Mar. Sci. Assoc., Townsville, Australia. Abstract p 25.
- Glynn P.W. 1988. Predation on coral reefs: Some key processes, concepts and research directions. Proc. Sixth Intern. Coral Reef Symp. 1:51-62.
- Glynn, P.W. 1990. Feeding ecology of selected coral reef macroconsumers: Patterns and effects on coral community structure. In: Z. Dubinsky (ed.), Coral reefs; ecosystems of the world 25, Elsevier Science Publishers, NY. pp. 365-400.
- Glynn, P.W. 1996. Coral reef bleaching: facts, hypotheses and implications. Global Change Biology. 2:495-509.

- Glynn, P.W., E.C. Peters, and L. Muscatine. 1986. Coral tissue microstructure and necrosis: Relation to catastrophic coral mortality in Panama. *Dis. Aquat. Org.* 1:29-37.
- Goenaga, C. and R.H. Boulon. 1992. The state of Puerto Rican and U.S. Virgin Island corals: An aid to managers. *Carib. Fish. Manage. Council.* 73 pages.
- Goldberg, W.M. and J.C. Makemson. 1981. Description of a tumorous condition in a gorgonian coral associated with a filamentous green algae. *Proc. Fourth Intern. Coral Reef Symp. Manila.* 2:685-697.
- Goldberg, W. M., J. C. Makemson, and S.B. Colley. 1984. *Entocladia endozoica* sp. nov., a pathogenic Chlorophyte: Structure, life history, physiology, and effect on its coral host. *Biol. Bull.* 166: 368-383.
- Goreau, T.J., J. Cervino, M. Goreau, R. Hayes, M. Hayes, L. Richardson, G. Smith, K. DeMeyer, I. Nagelkerken, J. Garzón-Ferreira, D. Gil, G. Garrison, E.H. Williams, L. Bunkley Williams, C. Quirolo, K. Patterson, J.W. Porter and K. Porter. 1998. Rapid spread of diseases in Caribbean coral reefs. *Rev. Biol. Trop.* 46 Suppl. 5:157-171.
- Green, E.P. and A. W. Bruckner. 2000. The significance of coral disease epizootiology for coral reef conservation. *Biological Conservation.* 96:347-361.
- Greenstein, B.J. and J.M. Pandolfi. 1997. Preservation of community structure in modern coral reef life and death assemblages of the Florida Keys: Implications for the Quaternary record of coral reefs. *Bull. Mar. Sci.* 19:39-59.
- Greenstein, B.J., H.A. Curran and J.M. Pandolfi. 1998. Shifting ecological baselines and the demise of *Acropora cervicornis* in the western North Atlantic and Caribbean province: a Pleistocene perspective. *Coral Reefs.* 17:249-261.
- Griffin, S. P. 1998. The effects of sunlight on the progression of black band disease. *Rev. Biol. Trop.* 46:175-179.
- Grosholz, E.D. and G.M. Ruiz. 1997. Evidence for local adaptation of black band disease at Carrie Bow Cay, Belize. *Proc. Eighth Intern. Coral Reef Symp.* 1:579:582.
- Guzmán, H.M. and C. Guevara. 1998. Morbilidad masiva de organismos arrecifales zooxantelados durante el blanqueamiento de 1995 en Cayos Cochinos, Honduras. *Rev. Biol. Trop.* 46 Suppl. 4:165-173.
- Hayes, R.L. and N. I. Goreau. 1998. The significance of emerging diseases in the tropical coral reef ecosystem. *Rev. Biol. Trop.* 46 Suppl. 5: 173-185.
- Harvell, C.D., K. Kim, J.M. Burkholder, R.R. Colwell, P.R. Epstein, D.J. Grimes, E.E. Hofmann, E.K. Lipp, A.D.M.E. Osterhaus, R.M. Overstreet, J.W. Porter, G.W. Smith, and G.R. Vasta. 1999. Emerging marine diseases: Climate links and anthropogenic factors. *Science.* 285:1505-1510.
- Harvell, D., K. Kim, C. Quirolo, J. Weir, and G. Smith. 2001. Coral bleaching and disease: contributors to 1998 mass mortality in *Briareum asbestinum* (Octocorallia, Gorgonacea). *Hydrobiologia.* 460: 97-104.
- Health Ecological and Economic Divisions (HEED). 1998. Marine Ecosystems: emerging diseases as indicators of change. *Health of the oceans from Labrador to Venezuela.* 85 pp.
- Hoetjes, P.C. 1998. Coral-list posting from July 9 on YBD outbreak in Curaçao.

- Hudson, J.H. 2000. First aid for massive corals infected with black band disease, *Phormidium corallyticum*: An underwater aspirator and post-treatment sealant to curtail reinfection. Diving for Science in the 21st Century, American Academy of Underwater Sciences, 430 Nahant Rd Nahant MA 01908 USA, Oct 2000, pp. 10-11.
- Hughes, T.P., 1994. Catastrophes, phase shifts, and large scale degradation of a Caribbean coral reef. *Science*. 265:1547-1551.
- Jaap, W.C., J.C. Halas and R.G. Muller. 1988. Community dynamics of stony corals (Scleractinia and Milleporina) at Key Largo National Marine Sanctuary, Key Largo, Florida during 1981-1986. *Proc. Sixth Intern. Coral Reef Symp.* 2:237-243.
- Jaap, W.C., J.W. Porter, J. Wheaton, K. Hackett, M. Lybolt, M. Callahan, C. Tskos, G. Yanev and P. Dustan. 2000. Coral reef monitoring project executive summary EPA Science Advisory Panel. Key Colony Beach, December 5-6, 2000.
- Kim, K., C.D. Harvell, P.D. Kim, G.W. Smith, and S.M. Merkel. 2000. Fungal disease resistance of Caribbean sea fan corals (*Gorgonia* spp.). *Mar. Biol.* 136:259-267.
- Kinne, O. 1980. Diseases of marine animals. In: O. Kinne (ed.), *Diseases of marine animals. Vol. 1: General aspects, protozoa to gastropoda*. John Wiley and Sons. pp.13-67.
- Knowlton, N., J.C. Lang, M.C. Rooney and P. Clifford. 1981. Evidence for delayed mortality in hurricane-damaged Jamaican staghorn populations. *Nature*. 294:251-252.
- Korrubel, J.L. and B. Riegl. 1998. A new coral disease from the southern Arabian Gulf. *Reef Sites. Coral Reefs*. 17:22.
- Kramer, P.A., R.N. Ginsburg, J.C. Lang, and R.S. Steneck. *In press* The Atlantic and Gulf Rapid Reef Assessment (AGRRA) program: A large-scale comparison of reef health indicators for the western Atlantic. *Atoll Research Bulletin*.
- Kushmaro, A., Y. Loya, M. Fine and E. Rosenberg. 1996. Bacterial infection and coral bleaching. *Science*. 380:396.
- Kuta, K.G. and L.L. Richardson. 1994. Distribution and frequency patterns of black band disease in the northern Florida Keys. *Abstract. Bull. Mar. Science*. 54:1078.
- Kuta, K.G. and L.L. Richardson. 1996. Abundance and distribution of black band disease on coral reefs in the northern Florida Keys. *Coral Reefs*. 15:219-223.
- Kuta, K.G. and L.L. Richardson. 1997. Black band disease and the fate of coral colonies in the Florida Keys. *Proc. Eighth Intern. Coral Reef Symp.* 1:575-578.
- Lauckner, G. 1980. Diseases of Cnidaria. In: O. Kinne (ed.), *Diseases of Marine Animals. Volume 1. General Aspects, Protozoa to Gastropoda*. John Wiley & Sons. Chapter 6.
- Laydo, R. 1984. Inference of a "white band" epidemic in the elkhorn coral *Acropora palmata* populations in Tobago, WI. *Proc. Assoc. Is. Mar. Labs. Carib. Abstract*. 18:12.
- Liddell, W.D. and S.L. Ohlhorst. 1992. Ten years of disturbance and change on a Jamaican fringing reef. *Proc. Seventh Intern. Coral Reef Symp.* 1:144-150.
- Littler, M.M. and D.S. Littler. 1995. Impact of CLOD Pathogen on Pacific Coral Reefs. *Science*. 267: 1356-1360

- Littler, M.M. and D.S. Littler. 1996. Black band disease in the South Pacific. *Reef Sites*. Coral Reefs. 15:20.
- Loya, Y., G. Bull, and M. Pichon. 1984. Tumour formation in scleractinian corals. *Helgol Wiss. Meeresunters.* 37:99-112.
- McField, M.D. 1999. Coral response during and after mass bleaching in Belize. *Bull. Mar. Sci.* 64(1):155-172.
- Miller, I. 1996. Black band disease on the Great Barrier Reef. *Reef Sites*. Coral Reefs. 15:58.
- Miller, J., C. Rogers and R. Waara. *In press*. Monitoring the coral disease, plague type II, on coral reefs in St. John, U.S. Virgin Islands. *Revista de Biología Tropical*.
- Mitchell, R. and I. Chet. 1975. Bacterial attack of corals in polluted seawater. *Microbial Ecol.* 2:227-233.
- Morse, D.E., A.N.C. Morse and H. Duncan. 1977. Algal "tumors" in the Caribbean Sea Fan, *Gorgonia ventalina*. *Proc. Third Intern. Coral Reef Symp.* 1:623-629.
- Morse, D.E., A.N.C. Morse, H. Duncan, and R.K. Trench. 1981. Algal tumours in the Caribbean Octocorallian, *Gorgonia ventalina*: II. Biochemical characterization of the algae, and first epidemiological observations. *Bull. Mar. Sci.* 31:399-409.
- Nagelkerken, I., K. Buchan, G.W. Smith, K. Bonair, P. Bush, J. Garzon-Ferrera, L. Botero, P. Gayle, C. Heberer, C. Petrovic, L. Porta and P. Yoshioka. 1997a. Widespread disease in Caribbean sea fans: I. Spreading and general characteristics. *Proc. Eighth Intern. Coral Reef Symp.* 1:679-682.
- Nagelkerken, I., K. Buchan, G.W. Smith, K. Bonair, P. Bush, J. Garzon-Ferrera, L. Botero, P. Gayle, C. Heberer, C. Petrovic, L. Porta and P. Yoshioka. 1997. Widespread disease in Caribbean sea fans: II. Patterns of infection and tissue loss. *Mar. Ecol. Prog. Ser.* 160:255-263.
- Nagelkerken, I., G.W. Smith, E. Shelden, M. Karel, and S. James. 1999. Sea urchin *Meoma ventricosa* die-off in Curaçao (Netherlands Antilles) associated with a pathogenic bacterium. *Dis. Aquat. Org.* 38:71-74.
- NCORE, 2001. Priorities for Caribbean Coral Research. Results from an International Workshop of the National Center for Caribbean Coral reef Research (NCORE). Edited by J.W. McManus. October 3-5, 2001. Miami, Florida.
- NOAA, 1999. Laminated Coral Disease Cards. 1) Disease and predation on scleractinian corals; 2) disease and predation: staghorn and elkhorn coral; 3) western Atlantic coral disease and predation: star coral; and 4) disease, predation and tissue anomalies affecting stony and soft coral. ©A.W. Bruckner, 1999.
- Nugues, M.M. 2002. Impact of a coral disease outbreak on coral communities in St. Lucia: What and how much has been lost? *Mar. Ecol. Prog. Ser.* 229: 61-71.
- Patterson, K.L., J.W. Porter, K.B. Ritchie, S.W. Polson, E. Mueller, E.C. Peters, D.L. Santavy and G.W. Smith. 2002. The etiology of white pox, a lethal disease of the Caribbean elkhorn coral, *Acropora palmata*. *PNAS* 99:8725-8730.
- Paul, J. H. and E. Vacelet. 1981. Elevated levels of microbial activity in the coral surface monolayer. *Mar. Ecol. Prog. Ser.* 33: 29-40.
- Peters, E.C. 1984. A survey of cellular reactions to environmental stress and disease in Caribbean scleractinian corals. *Helgol. Wiss. Meeresunters.* 37:113-137.

- Peters, E.C. 1993. Diseases of other invertebrate phyla: Porifera, cnidaria, ctenophora, annelida, echinodermata. In: Couch and Fournie (eds.), Adv. Fish. Science, Pathology of Marine and Estuarine Organisms. CRC Press, Florida. pp. 393-449.
- Peters, E.C. 1997. Diseases of coral-reef organisms. In: C. Birkeland (ed.). Life and Death of Coral Reefs. Chapman & Hall, New York. pp. 114-139.
- Peters, E.C. 2002. Coral disease diagnostics: histopathology. Coral Disease and Health Consortium.
- Peters, E.C., J.J. Oprandy and P.P. Yevich. 1983. Possible causal agent of white band disease in Caribbean scleractinian corals. J. Invert. Pathol. 41:394-396.
- Peters, E.C., J.C. Halas, and H.B. McCarty. 1986. Calicoblastic neoplasms in *Acropora palmata*, with a review of reports on anomalies of growth and form in corals. J. Nat. Cancer Inst. (JNCI). 76:895-912.
- Peters, E.C. and H.B. McCarty. 1996. Carbonate Crisis. Geotimes. 4/96:20-23.
- Peters, E.C., N.J. Gassman, J.C. Firman, R.H. Richmond, and E.A. Power. 1997. Ecotoxicology of tropical marine ecosystems. 1997. Annual Review Issue on Tropical Ecotoxicology. Environ. Toxicol. Chem. 16(1):12-40.
- Polson, S. W. and G.W. Smith. Systematic 2000. Characterization and comparison of various bacterial isolates pathogenic or putatively pathogenic to scleractinian coral species. Abstract, Ninth International Coral Reef Symposium, Bali, Indonesia. p.
- Porter, J.W. and O.W. Meier. 1992. Quantification of loss and change in Floridian reef coral populations. Amer. Zool. 23:625-640.
- Porter, J.W. and J. I. Tougas. 2001. Reef ecosystems: threats to their biodiversity. Encyclopedia of Biodiversity. 5:73-95.
- Porter, J.W., P. Dustan, W.C. Jaap, J.L. Wheaton, K.L. Patterson, V. Kosmynin, O. Meier, M.E. Patterson, and M. Parsons 2001. Patterns of spread of coral disease in the Florida Keys. Hydrobiologia. 460:1-24.
- Raghukumar, C. and S. Raghukumar. 1991. Fungal invasion of massive corals. PSZNI Mar. Ecol. 4:329-358.
- Ramos-Flores, T. 1983. Lower marine fungus associated with black line disease in star corals (*Montastrea annularis*). Biol. Bull. 165:429-435.
- Ravindran, J., C. Raghukumar, and S. Raghukumar. 1999. Disease and stress-induced mortality of corals in Indian Reefs and observations on bleaching of corals in the Andamans. Current Science (India Academy of Sciences) 76(2): 233-237.
- Ravindran, J., C. Raghukumar and S. Raghukumar. 2001. Fungi in *Porites lutea*: association with healthy and diseased corals. Dis. Aquat. Org. 47: 219-228.
- Raymundo, L.J.H., and C.D. Harvell, in press. Porites Ulcerative White Spot Disease: A new disease impacting Indo-Pacific Coral Reefs.
- Red Sea Marine Park. 2001. The pink-blue disease in *Acropora* sp. www.iui-eilat.ac.il/peacepark.htm
- Reeves, L. 1994. Newly discovered: Yellow band disease strikes Keys Reefs. Underwater USA. 11(8):16.

- Richardson, L.L. 1992. Red band disease: a new cyanobacterial infestation of corals. Proc. Tenth Ann. Amer. Acad. Underw. Sci. 153-160.
- Richardson, L.L. 1995. Monitoring and Assessment of Coral Reef Health: Coral Disease Incidence and Cyanobacterial Blooms as Reef Health Indicators. A Coral reef Symposium on Practical, reliable, Low Cost Monitoring Methods for Assessing the Biota and Habitat Conditions of Coral Reefs. Jan 26-27 Annapolis, MD. pp. 51-56.
- Richardson, L.L. 1996. Horizontal and vertical migration patterns of *Phormidium corallyticum* and *Beggiatoa* spp. associated with black-band disease of corals. Microbial Ecol. 32:323-325.
- Richardson, L.L. 1997. Occurrence of the black band disease cyanobacterium on healthy corals of the Florida Keys. Bull. Mar. Sci. 61: 485-490.
- Richardson, L.L. 1998. Coral diseases: what is really known. Trends Ecol. Evol. 13 :438-443.
- Richardson, L.L. 1999. Emerging coral diseases. U.S. Global Change research Program Seminar Series. June 22, 1999.
- Richardson, L.L. 2000. Coral epidemiology. In Opportunities for Environmental Applications of Marine Biotechnology. Proc. Oct. 5-6 Workshop. Board on Biology, National research Council, National Academy Press. pp. 85-95.
- Richardson, L.L. and R.G. Carlton. 1993. Behavioral and chemical aspects of Black Band Disease of corals: An *in situ* field and laboratory study. In: John Heine (Ed.) Diving For Science. Proc. Amer. Acad. Underw. Sci. Tenth Ann. Sci. Diving Symp. 2:107-116.
- Richardson, L.L., K.G. Kuta, S. Schnell and R.G. Carlton. 1997. Ecology of the black band disease microbial consortium. Proc. Eighth Intern. Coral Reef Symp. 1:597-600.
- Richardson L.L, W.M. Goldberg, K.G. Kuta, R.B. Aronson, G.W. Smith, K.B. Ritchie, J.C. Halas, J.S. Feingold and S.L. Miller. 1998a. Florida's mystery coral-killer identified. Nature. 392:557-558.
- Richardson L.L, W.M. Goldberg, R.G. Carlton and J.C. Halas. 1998b. Coral disease outbreak in the Florida Keys: Plague type II. Rev. Biol. Trop. 46 Suppl. 5:187-198.
- Richardson L.L., G.W. Smith, K.B. Ritchie, and R.G. Carlton. (in press). Integrating microbiological, microsensor, physiologic and molecular techniques in the study of coral disease pathogenesis. Hydrobiologia
- Richardson, L.L. and R.A. Aronson. In press. Infectious diseases of reef corals. Proc. Ninth Intern. Coral Reef Symp.
- Ritchie, K.B., and G.W. Smith. 1997. Physiological comparison of bacterial communities from various species of scleractinian corals. Proc. Eighth Intern. Coral Reef Symp. 1:521-526.
- Ritchie, K.B. and G.W. Smith. 1998. Type II white-band disease. Rev. Biol. Trop. 46 Suppl. 5:199-203.
- Ritchie, K.B. and G.W. Smith. 1995. Preferential carbon utilization patterns by surface bacterial communities from water mass, normal and white-band diseased *Acropora cervicornis*. Molec. Mar. Biol. Biotechnol. 4:345-352.

- Rogers, C. 1985. Degradation of Caribbean and Western Atlantic coral reefs and decline of associated fisheries. Proc. Fifth Intern. Coral Reef Symp. Tahiti. 6:491-496.
- Rosenberg, E. 1999. Inhibition of photosynthesis and bleaching of zooxanthellae by the coral pathogen *Vibrio shiloi*. Environ. Microbiol. 1(3):223-230.
- Rützler, K. and D.L. Santavy. 1983. The black band disease of Atlantic reef corals. 1. Description of the cyanophyte pathogen. PSZNI (Mar. Ecol.). 4:301-319.
- Rützler, K., D.L. Santavy and A. Antonius. 1983. The black band disease of Atlantic reef corals. 111. Distribution, ecology, and development. PSZNI (Mar. Ecol.). 4:329-358.
- Santavy, D.L., W.D. Jeffrey, R.A. Snyder, J. Campbell, P. Malouin and L. Cole. 1994. Microbial community dynamics of healthy and stressed coral hosts. Bull. Mar. Sci. Abstract. 54:10-1078.
- Santavy, D.L., T. Schmidt, S.S. Wilkinson, D.H. Buckley and A.W. Bruckner. 1996. The phylogeny and alliance of cyanobacteria affiliated with two band diseases in corals from the western Atlantic. Abstract. 96th ASM Meeting.
- Santavy, D.L., D.H. Buckley, A.W. Bruckner and T. Schmidt 1996. Cyanobacteria of black-band and red-band diseases affecting western Atlantic corals. Abstract.
- Santavy, D.L. and E.C. Peters. 1997. Microbial pests: Coral disease research in the western Atlantic. Proc. Eighth Intern. Coral Reef Symp. 1:607-612.
- Santavy, D.L., E.C. Peters, C. Quirolo, J.W. Porter, and C.N. Bianchi. 1999. Yellow-blotch disease outbreak on reefs of the San Blas Islands, Panama. *Reef Sites*. Coral Reefs. 18:97.
- Santavy, D.L., E. Mueller, E.C. Peters, L. MacLaughlin, J.W. Porter, K.L. Patterson and J. Campbell. 2001. Quantitative assessment of coral diseases in the Florida Keys: Strategy and methodology. Hydrobiologia. 460: 39-52.
- Shinn, E.A. 1989. What is really killing the corals. Sea Frontiers. 35: 72-81.
- Shinn, E.A., G.W. Smith, J.M. Prospero, P. Betzer, M.L. Hayes, V. Garrison and R.T. Barber. 2000. African dust and the demise of Caribbean coral reefs, Geophysical Research Letters. 27:3029-3032.
- Smith, G.W. 1999. The Decline of the Coral Reef--Coral Bleaching and Diseases. SciTalk6: Coral Reef Discussion. The National Health Museum.
- Smith, G.W. 2000. Pathogens of Coral Reef Organisms. Ninth International Coral Reef Symposium. abstract
- Smith, G.W., L.D. Ives, I.A. Nagelkerken and K.B. Ritchie. 1996. Caribbean sea fan mortalities. Nature. 383:487.
- Smith, G.W., C.D. Harvell and K. Kim. 1998. Response of sea fans to infection with *Aspergillus* sp. (fungi). Rev. Biol. Trop. 46 Suppl. 5:205-208.
- Smith, G.W., C.D. Harvell, K. Kim, K.B. Ritchie, S.C. James, and K.C. Buchan. 1998 (in press). Cellular events occurring during the pathogenesis of aspergillosis of *Gorgonia* species. Proc. Symp. Natural History of the Bahamas.
- Taylor, D.L. 1983. The black band disease of Atlantic reef corals. 11. Isolation, cultivation and growth of *Phormidium coralyticum*. PSZNI (Mar. Ecol.). 4:321-328.

- TeStrake, D., W.C. Jaap, E. Truby and R. Reese. 1988. Fungal filaments in *Millepora complanata* Lamarck, 1815 (Cnidaria:Hydrozoa) after mass expulsion of zooxanthella. *Biological Sciences*. 5:184-188.
- Toren, A., L. Landau, A. Kushmaro, Y. Loya, and E. Rosenberg. 1998. Effect of temperature on adhesion of *Vibrio* strain AK-1 to *Oculina patagonica* and on coral bleaching. *Appl. Environ. Microbiol.* 64:1379-1384.
- Upton, S.J. and E.C. Peters. 1986. A new and unusual species of *Coccidium* (Apicomplexa: Agamococcidiorida) from Caribbean Scleractinian Corals. *Journal of Invertebrate Pathology*. 47:184-193.
- Viehman, T.S and L.L. Richardson. *In press*. Motility patterns of *Beggiatoa* and *Phormidium coralyticum* in black band disease. *Proc. Ninth Intern. Coral Reef Symp.*
- M.F. Walker, G.W. Smith and L.L. Richardson. 2000. Characterization of a Bacterial Pathogen Associated with Plague Type II: Chemotaxis and Mechanism of Coral Tissue Necrosis, Abstract. 25th Eastern Fish Health Workshop. Plymouth, MA, April, 2000.
- Weil, E., I. Urreiztieta and J. Garzon-Ferreira. *in press*. Geographic variability in the incidence of coral and octocoral diseases in the wider Caribbean. *Proc. Ninth Intern. Coral Reef Symp.*
- Weil, E., G.W. Smith and M. Mills. 2001. Spatial and Temporal Variability of Coral and Octocoral Disease in Bermuda. 30th Scientific Meeting, Assoc. Mar. Labs. Caribbean, La Parguera, June, 2001.
- Weil, E. 2001. Caribbean coral reef diseases, status and research needs. In: J.W. McManus (ed.) *Priorities for Caribbean Coral Research*. Weil 1-8.
- Weil, E and I. Urreiztieta. 2000. Latitudinal variability in coral disease in the wider Caribbean. Abstract, Caribbean Coastal Marine Productivity Network.
- Weir, J.R., V. Garrison, G.W. Smith, and E.A. Shinn. *In press*. The relationship between gorgonian coral (Cnidaria: Gorgonacea) disease and African dust storms. *Proc. Ninth Intern. Coral Reef Symp.*
- Wilkinson, C., O. Linden, H. Cesar, G. Hodgson, J. Rubens, and A.E. Strong. 1999. Ecological and socioeconomic impacts of 1998 coral mortality in the Indian Ocean: An ENSO impact and a warning of future change? *Ambio*. 28:188-196.
- Williams, C. 1996. Diseased reef alarms researchers. *Geotimes*. 3/97 pp. 6.
- Williams, E.H., Jr., P.J. Bartels, and L. Bunkley-Williams. 1999. Predicted disappearance of coral-reef ramparts: A direct result of major ecological disturbances. *Global Change Biol.* 5:839-845.
- Williams, E.H., Jr. and L. Bunkley-Williams. 1990. The world-wide coral reef bleaching cycle and related sources of coral mortality. *Atoll Res. Bull. (Smithsonian)*. 335:1-71.
- Williams, E.H., Jr., and L. Bunkley-Williams. 2000. Major marine ecological disturbances of the Caribbean. *Infect. Dis. Rev.* 2(3):110-127.
- C. M. Woodley, D. L. Santavy, W. H. Walker, A.W. Bruckner, D. W. Howard, and S. M. McLaughlin. 2001. Coral Disease And Health Consortium: Finding Solutions. Abstract.
- Woodley, J.D., E.A. Chomesky, P.A. Clifford, J.B.C. Jackson, L.S. Kaufman, N. Knowlton, J.C. Land, M.P. Pearson, J.W. Porter, M.C. Rooney, K.W. Rylaarsdam, V.J. Tunnicliffe, C.M. Wahle, J.L. Wulff, A.S.C. Curtis, M.D. Dallmeyer, B.P. Jupp, M.A.R. Koehl, J. Niegel and E.M. Sides. 1981. Hurricane Allen's impact on Jamaican coral reefs. *Science*. 214: 749-755.

van Woesik, R. 1998. Lesion healing on massive *Porites* spp. corals. Mar. Ecol. Prog. Ser. 164:213-220.

Yamashiro, H., M. Yamamoto, and R. van Woesik. 2000. Tumor formation on the coral *Montipora informis*. Dis. Aquat. Org. 41:211-217.

Zorpette, G. 1995. More coral trouble. Sci. Amer. 273:36-37.