

# IOM Report, 2002, Tables 4-1, 6-1, 6-2

	<b>Fever</b>	<b>Systemic, Any</b>	<b>Erythema or Swelling</b>	<b>Pain, Any</b>
<b>Acellular pertussis</b>	<b>0 - 7%</b>	<b>17 - 29%</b>	<b>12 - 15%</b>	<b>51 - 77%</b>
<b>Hepatitis A</b>	<b>0 - 3%</b>	<b>4 - 22%</b>	<b>4 - 40%</b>	<b>40 - 52%</b>
<b>Hepatitis B</b>	<b>0 - 4%</b>	<b>10%</b>	<b>1 - 99%</b>	<b>11 - 43%</b>
<b>Influenza</b>	<b>1 - 13%</b>	<b>11 - 34%</b>	<b>11 - 21%</b>	<b>24 - 86%</b>
<b>Rabies</b>	<b>2 - 18%</b>	<b>3%</b>	<b>1 - 18%</b>	<b>4 - 52%</b>
<b>Tetanus - diphtheria (Td)</b>	<b>1 - 9%</b>	<b>17 - 26%</b>	<b>22 - 35%</b>	<b>43 - 85%</b>
<b>Anthrax</b>	<b>1 - 8%</b>	<b>1 - 36%</b>	<b>3 - 42%</b>	<b>Sore: 67 - 83%</b>

[www.nap.edu/catalog/10310.html](http://www.nap.edu/catalog/10310.html)