



October 6, 2009

Jeannine Cody
Permits, Conservation, and Education Division
National Marine Fisheries Service (NMFS)
Office of Protected Resources
1315 East-West Highway
Silver Spring, MD 20910-3226

Dear Ms. Cody,

PRBO Conservation Science (PRBO), along with partners Oikonos Ecosystem Knowledge and Point Reyes National Seashore, is requesting renewal of our Incidental Harassment Authorization (IHA) under the Marine Mammal Protection Act. This authorization allows harassment of small numbers of marine mammals incidental to conducting seabird and pinniped research operations on Southeast Farallon Island, Año Nuevo Island, and Point Reyes National Seashore in central California. Marine mammals species that could be potentially impacted by these activities include the California stocks of northern elephant seal (*Mirounga angustirostris*), harbor seal (*Phoca vitulina richardii*), Steller sea lion (*Eumetopias jubatus*), and California sea lion (*Zalophus californianus*).

Our current IHA expires on December 11, 2009. We would like to renew this authorization for another year. All actions are the same at all sites as documented in our current IHA. We are requesting the same level of take for all species, with the exception of California Sea Lions (CASL) on Southeast Farallon Island (SEFI).

We are requesting an increase in our take of CASL on SEFI from the current level of 1,500 animals to 5000 animals. We base this request on two primary reasons. First, the number of CASL at SEFI has increased dramatically in recent years, with maximum counts exceeding 11,000 animals (Figure 1). These increases may be related to recent shifts in ocean climate. These abundances are unprecedented in our 4 decades of continuous research on SEFI, and current maximum abundances are well in excess of double those observed in the 1990's and early 2000's. It was these earlier abundances which were used to estimate a take of 1,500 animals per year. In addition, the distribution of CASL on SEFI has shifted rapidly in recent years. There has been a large influx of animals hauling up in landing coves and at our primary elephant seal study site during the winter months. This has caused a major reduction in our ability to collect valuable data on the breeding success of this elephant seal population due to take concerns for CASL.

Landings are required for island operations so impacting animals during those events is unavoidable. Therefore, we are requesting a significant increase in our CASL take on SEFI, from 1,500 to 5,000 animals. We see this increase as appropriate, based upon the increasing number and shifting distribution of the animals. As none of the sites we visit on SEFI are breeding areas for CASL, no nursing females or pups will be affected by any potential increase in take. We will continue to take all possible efforts to minimize any potential pinniped take.




Figure 1. Maximum number of California Sea Lions observed on Southeast Farallon Island weekly surveys, by year, from 1970-2008.

Thank you for reviewing this request and we look forward to hearing from you.

Sincerely,

Russell Bradley
Farallon Program Manager
Marine Ecology Division
PRBO Conservation Science
rbradley@prbo.org

Request for Marine Mammal Protection Act
Incidental Harassment Authorization

**Seabird Research on the South Farallon Islands, Año Nuevo Island,
and Point Reyes National Seashore**

Submitted by

PRBO Conservation Science
3820 Cypress Drive #11
Petaluma, CA 94954



Oikonos Ecosystem Knowledge
P.O. Box 1932
Benicia, CA 94510



Point Reyes National Seashore
1 Bear Valley Rd.
Point Reyes Station, CA 94956



To:

Permits, Conservation, and Education Division
National Marine Fisheries Service (NMFS)
Office of Protected Resources
1315 East-West Highway
Silver Spring, MD 20910-3226

I. Locations:

Numerous marine sanctuaries, refuges, and parks are designated within the action area. These include the Monterey Bay and Gulf of the Farallones National Marine Sanctuaries; Point Reyes National Seashore, Farallones National Wildlife Refuge; and Anõ Nuevo State Park and Reserve.

The action area consists of the following 3 locations:

1. **South Farallon Islands (SFI)** ($37^{\circ} 41' 55''$ / $123^{\circ} 00' 10''$) – consists of Southeast Farallon Island (SEFI) and West End Island (WEI). These two islands are directly adjacent to each other and separated by only a ~30ft channel. The SFI have a land area of approximately 120 acres (49 ha) and are part of the Farallon National Wildlife Refuge. The islands are located near the edge of the continental shelf 28 miles west of San Francisco, California. SFI lie within the waters of the Gulf of the Farallones National Marine Sanctuary. These waters have been designated as EFH for numerous groundfish species (Farallon Island /Fanny Shoal area), salmon species (Chinook and Coho Salmon), and pelagic species (Northern Anchovy and Pacific Sardine). These waters constitute foraging areas for all species of pinnipeds listed in this application.
2. **Anõ Nuevo Island (ANI)** is located 1/4 mile offshore of Anõ Nuevo Point in San Mateo County, California ($37^{\circ}06' N$, $122^{\circ}20' W$). This small 25 acre island is part of the 4000 acre Anõ Nuevo State Reserve, all of which is owned and operated by California State Parks. ANI lies within the Monterey Bay National Marine Sanctuary and the newly established Anõ Nuevo State Marine Conservation Area. Like SFI, the waters adjacent to ANI are foraging areas for pinnipeds and represent EFH for many groundfish, salmon, and pelagic forage fish species.
3. **Point Reyes National Seashore (PRNS)** is located 40 miles north of San Francisco Bay. The affected area for the purposes of this document are the headland coastal areas of this large National Park. Waters adjacent to PRNS are foraging areas for pinnipeds and represent EFH for many groundfish, salmon, and pelagic forage fish species. In addition, PRNS lies within close proximity (6 miles) of the Cordell Bank National Marine Sanctuary.

1 and 2. Detailed Description of the Activities and Dates, Duration, and Region of Activity

South Farallon Islands

PRBO Conservation Science has conducted year round wildlife research and monitoring activities at the SFI, part of the Farallon National Wildlife Refuge, since 1968. This work is conducted through a collaborative agreement with the USFWS. While some research focuses on marine mammals, see NMFS Scientific Permit 373-1575, other research focusing on seabirds, and some procedures involved in maintaining the SEFI field station may involve incidental take of marine mammals.

A. Seabird Research and Boat Landings

Seabird research activities involve observational and marking (i.e. netting and banding for capture-mark-recapture) studies of breeding seabirds. Occasionally researchers may travel to coastal areas of the island to conduct observational seabird research where non breeding marine mammals are present. These sorts of tasks include viewing breeding seabirds from an observation blind or censusing shorebirds. This activity usually involves one or two observers. Access to the refuge involves landing in small boats, 14-18ft open motorboats which are hoisted onto the island using a derrick system. These activities can also result in incidental take of pinnipeds.

Research on SEFI is conducted year round. Most intertidal areas of the island, where marine mammals are present, are rarely visited in seabird research. Most potential for incidental take will occur at the island's 2 landings, North Landing and East Landing (Figure 1). These sites are visited ~ 1-3 times per day by researchers for a maximum of 500 visits per year. In both locations researchers are located >50 ft above any pinnipeds which may be present. These pinnipeds are primarily California Sea Lions or Northern Elephant Seals, to a lesser extent Harbor seals, and very rarely Steller Sea Lions. Most visits to these areas are brief (~15 minutes), though seabird observers are present from 2-5 hours daily at North Landing from early April – early August each year to conduct observational studies on breeding Common Murres. Boat landings to re-supply the field station, lasting 1-3 hours, are conducted once every 2 weeks at one of the these locations. Activities involve launching of the boat with one operator, with 2-4 other researchers assisting with the operations from land. At East Landing, the primary landing site, all personnel assisting with the landing stay on the loading platform 30 ft above the water. At North Landing, loading operations occur at the water level in the intertidal.

B. Pinniped Research

The purpose of northern elephant seal research on SFI, which could result in Level B harassment to SSLs, is to 1) determine how northern elephant seal population dynamics and health changes in response to oceanographic events and anthropogenic impacts and 2) monitor population expansion into new areas where there is potential for negative interactions with other wildlife and humans (Allen 2004). This location is one of many sites in CA where pinniped research would take place but is the only location where harassment to SSLs could occur. That is, the objectives of the northern elephant seals population and health assessment study are to monitor northern elephant seal population dynamics and health along north central California from the Russian River (RR) in Sonoma County to Point Reyes (PR) and out to the South Farallon Islands (SFI; Southeast Farallon Island and West End Island)

We will compare the age and sex specific survival and reproductive rates of seals tagged between 1976 and 2011 and describe trends in survival and reproduction relative to intrinsic (e.g., density) and extrinsic (e.g., disease and environmental) factors. Using datasets available from the Marine Mammal Center, we will correlate prevalence of disease or the occurrence of environmental events such as ENSO with changes in population status.

Each year, we flipper tag a maximum of 2,000 weaned pups and conduct resighting surveys of tagged adult animals to determine survival and reproductive rates. Three to 4 researchers are present at SFI for this research annually, conducting work daily between the first week of December and the second week of March. Daily time spent near the proximity of elephant seals is ~4 hours. Seals that are incidentally disturbed during the tagging of pups, but their response is minimal and seals very rarely leave the area; generally within minutes they return to pre-disturbance behavior (S. Allen and W. Sydeman pers. observations). We will mark with dye and flipper tags dominant males in order to track their movements between sites (Farallones, PR, etc.). Monitoring the movement of males between and within seasons assists us in determining where new colonies may form because males use sites prior to females initiating breeding at a new colony site.

In addition to the research described above, there is also the potential for harassment of Stellers Sea Lions during marine mammal research activities. Takes of Stellers may occur during winter surveys (December-February) on West End Island (Figure 1) to survey breeding elephant seals. These surveys involve ground transport and observers moving approximately 1500 ft above pinniped colonies in transit to elephant seal areas. There are approximately 5 surveys per year between Early December and late February, each lasting approximately 2 hours, which could result in harassment to Steller sea lions. However, transit above Steller sea lion areas only last approximately 30 minutes maximum duration. Based on previous trips, it is estimated up to 16 Steller sea lions could potentially be harassed each year. During the research activities every effort will be made to avoid incidental disturbance; however, large gatherings of species co-occur on beaches where research is conducted. Thus the capture of one species may incidentally involve the disturbance of other species. Researchers keep low profiles and move slowly to avoid stampeding the seals. These surveys occur in the non-breeding period where there are no breeding females or dependant young pups present.

Permit No. 373-1868-00, issued to The Point Reyes Bird Observatory (PRBO) Conservation Science (Dr. William J. Sydeman, Responsible Party), Petaluma, CA, issued on March 27, 2007 authorizes PRBO to study and monitor population trends, health, and ecology of pinnipeds in California, specifically around the Farallon Islands, Point Reyes Peninsula, San Francisco Bay, and in Sonoma County near the Russian River. Harbor seals and northern elephant seals are the species primary species of study and that which are involved in capture, handling, and sampling. Incidental harassment of elephant seals, harbor seals, California sea lions, and northern fur seals is authorized during research. Takes would be limited to Level B harassment and would likely occur when we are transiting to and from West End Island by foot. The authorization of up to 16 Steller sea lions annually is requested for coverage of Level B harassment incidental to capture and handling activities on California sea lions (Costa), northern elephant seals, and harbor seals (PRBO).

Anõ Nuevo Island

PRBO has also conducted seabird research and monitoring activities on Año Nuevo Island (ANI), part of the Año Nuevo State Reserve, since 1992. Collaborations with Oikonos - Ecosystem Knowledge began in 2001 to research seabird burrow nesting habitat quality and restoration. All work is conducted through a collaborative agreement with California State Parks. Some procedures involved in accessing the island (by 12' Zodiac boat) or seabird research may involve incidental take of marine mammals. Non-breeding pinnipeds may occasionally be present on the small beach in the center of the island where the boat is landed. California sea lions may also occasionally be present near a small group of subterranean seabird nest boxes on the island terrace. There are usually 2-3 researchers involved in island visits.

Research on ANI is conducted once/week April-August, and occasional intermittent visits are made during the rest of the year. The maximum number of visits per year would be 20. A component of the seabird research involves nesting habitat restoration and monitoring which requires sporadic visits from September-November, between the seabird breeding season and the elephant seal pupping season. Most intertidal areas of the island where marine mammals are present are not ever visited during seabird research, excepting the landing beach. (Figure 2) Most potential for incidental take will occur at this location as well as just north of this beach up on the island's terrace where a small number of seabird nest boxes are located. The landing beach is visited upon arrival and departure during the weekly visit, and the nest boxes are checked one time that day. In both locations researchers are located >50 ft away from any pinnipeds which may be hauled out. Landings and visits to nest boxes are brief (~15 minutes).

Point Reyes

The National Park Service conducts research, resource management and routine maintenance services at Point Reyes National Seashore. This involves both marine mammal research, see NMFS Scientific Permit 373-1575, and seabird research involving maintaining the facilities around the Seashore. Both types of work may involve incidental take of marine mammals. Additionally, habitat restoration of the seashore includes restoration and removal of non-native invasive plants, and coastal dune habitat. Non-native plant removal is timed to avoid the breeding seasons of pinnipeds; however, on occasion non-breeding animals may be present at various beaches throughout the year. Additionally, elephant seals will haul out on human structures and block access to facilities. For example, they haul out on a boat ramp at the Life Boat Station and in various car parking lots around the seashore.

Research along the Seashore includes monitoring seabird breeding and roosting colonies. Seabird monitoring usually involves one or two observers. Surveys are conducted by small boats, 14-22 ft open motorboats that survey along the shoreline. These activities can also result in incidental take of pinnipeds.

Research at PRNS is conducted year round, with an emphasis during the seabird nesting season with occasional intermittent visits the rest of the year. The maximum number of visits per year is 20. A component of the seabird research involves habitat restoration and

monitoring which requires sporadic visits from September-November, between the seabird breeding season and the elephant seal pupping season. Most areas where research occurs and where marine mammals are present are not ever visited, excepting the landing beaches along Point Reyes Headland. In all locations researchers are located >50 ft away from any pinnipeds which may be hauled out. Elephant seals may haul out on boat ramps and parking lots year round. To maintain access of the boat ramps for park enforcement rangers and researchers, we will need to move seals from the boat ramp and adjacent parking lot for safety to seals and people. Removal will use the minimum tool which includes shaking a blue tarp or using plywood boards created by the Marine Mammal Center for moving seals. Both methods are effective in moving all age classes of elephant seals (F. Gulland MMC; R. DeLong, NMFS, pers. com.). Intermittent visits to areas of PRNS where pinniped takes may occur also conducted for 1) research on other species such as seabirds, sharks, and subtidal mapping and 2) resource management activities such as non-native plant management and intertidal monitoring.

3. Species and current population size of affected Marine Mammals in Area

California stock of:

Northern elephant seal (*Mirounga angustirostris*)

Harbor seal (*Phoca vitulina richardii*)

Steller sea lion (*Eumetopias jubatus*)

California sea lion (*Zalophus californianus*)

Farallon Islands populations (approximate annual range, PRBO unpublished data)

Northern elephant seal (*Mirounga angustirostris*) (~50-750)

Harbor seal (*Phoca vitulina richardii*) (~40-120)

Steller sea lion (*Eumetopias jubatus*) (~50-150)

California sea lion (*Zalophus californianus*) (~1,000–7,000)

Año Nuevo Island populations (approximate annual range, UCSC and NOAA- Mark Lowry, Southwest Fisheries Science Center, La Jolla, CA, unpublished data)

Northern elephant seal (*Mirounga angustirostris*) (~900-1000)

Harbor seal (*Phoca vitulina richardii*) (~100-150)

Steller sea lion (*Eumetopias jubatus*) (~400-600)

California sea lion (*Zalophus californianus*) (~4,000–9,500)

Point Reyes populations (approximate annual range, NPS unpublished data)

Northern elephant seal (*Mirounga angustirostris*) (~100-2,000)

Harbor seal (*Phoca vitulina richardii*) (~20-4,000 at nine locations)

California sea lion (*Zalophus californianus*) (~100–500)

4. Status and Distribution of the Affected Species

Northern Elephant Seal: The northern elephant breeding population is distributed from central Baja California, Mexico to the Point Reyes Peninsula in northern California.

Along this coastline there are 13 major breeding colonies. The Northern elephant seal was exploited for its oil during the 18th and 19th centuries and by 1900 the population was reduced to 20-30 individuals on Guadalupe Island (Hoelzel et al. 1993, Hoelzel 1999). As a result of this bottleneck the genetic diversity found in this species is extremely low (Hoelzel 1999). The recent formation of most rookeries indicates that there is no genetic differentiation among populations. Although movement and genetic exchange occurs among colonies, most seals return to their natal site to breed (Huber et al. 1991). Recolonization of their former breeding range progressed north from the San Benito and Guadalupe Islands off Baja California to the most recent northernmost breeding site at Point Reyes Headlands. In the last three decades, annual pup production has increased at the rate of $9.43 \pm 0.51\%$ per year in California and $5.19 \pm 0.33\%$ per year over the entire range (Barlow et al. 1993). A complete population count of elephant seals is not possible because all age classes are not ashore at the same time. Elephant seal population size is usually estimated by counting the number of pups produced and multiplying by the inverse of the expected ratio of pups to total animals (McCann 1985). Stewart et al. (1994) used McCann's multiplier of 4.5 to extrapolate from 28,164 pups to a population estimate of 127,000 elephant seals in the U.S. and Mexico in 1991. The multiplier of 4.5 was based on a stable population. Boveng (1988) and Barlow et al. (1993) argue that a multiplier of 3.5 is more appropriate for a rapidly growing population such as the California stock of elephant seals. Based on the estimated 28,450 pups born in California and this 3.5 multiplier, the California stock was approximately 101,000 in 2001 (Carretta et al. 2002). At Point Reyes, the population grew at 32.8% per year between 1988 and 1997 (Sydeman and Allen 1999) and around 10% per year since 2000 (S. Allen unpubl. data), and in 2006 around 700 pups were born at three primary breeding areas. The population on the Farallon Islands has declined by 3.4% per year since 1983, and in recent years numbers have fluctuated between 100 and 200 pups (W. Sydeman, D. Lee, unpubl. data).

Elephant seals congregate in central California to breed from late November to March. Females typically give birth to a single pup and attend the pup for up to 6 weeks. Breeding occurs after the pup is weaned by attending males. After breeding, seals migrate to the Gulf of Alaska or deeper waters in the eastern Pacific. Adult females and juveniles return to terrestrial colonies to molt in April and May, and males return in June and July to molt, remaining onshore for around 3 weeks.

Pacific Harbor Seal: Harbor seals are one of the most widely distributed northern hemisphere pinnipeds and are found in coastal, estuarine and some times fresh water of both the Atlantic Ocean and Pacific Oceans. There is considerable regional genetic differentiation between harbor seal populations as they are generally limited in migratory movements. Presently, there are three recognized stocks (California, Oregon and Washington Coast, and Inland Washington). There is some question whether the San Francisco Bay population may be a separate stock based on genetic analyses (D. German, Sonoma State University, pers. com.). There is some limited movement among stocks; however, only a small portion of harbor seals move outside state waters (Lamont et al. 1996).

In central California, harbor seals breed annually from March through May and molt in June and July. Females give birth to a single pup and attend the pup for around 30 days, at which time they wean pups. Mating occurs in the water around the time of weaning. Harbor seals are resident year round at terrestrial colonies; however, juveniles may disperse to other colonies ranging up to ~ 500 km. Individual adult seals may also migrate widely from breeding colonies.

Given the wide distribution of harbor seals it is not surprising that their population trends vary widely. Harbor seal populations in the Eastern North Pacific along the West Coast of the United States are all increasing. Along the coast of Washington and Oregon harbor seals increased in number at a rate of between 4 to 7% per annum with an estimated population of over 30,000 (Jefferies et al 1997). Additionally, along the California coast harbor seal numbers have increased at 3.5% per year from 1982 to 1995 with a minimum population of approximately 28,000 (Hanan and Beeson 1994, Carretta et al. 2001). Brown et al. (2005) estimated a population of 10,087 harbor seals in Oregon. There is evidence that the population may have reached equilibrium off Oregon and Washington and central California (Barlow et al. 1998, Sydeman and Allen 1999, NOAA Stock Assessment 2003). The most recent population estimate for California based on mark-recapture analysis 43,449 based on a correction factor of 1.65, and the population at Point Reyes was estimated to be 7,524 for the molt season based on the same correction factor (Lowry et al. 2005, Manna et al. 2006).

California Sea Lion: California sea lions range from southern Mexico up to British Columbia and breed almost entirely on islands in southern California, Western Baja California and the Gulf of California. In recent years, California sea lions have begun to breed annually in small numbers at Año Nuevo Island and South Farallon Island, California. One abandoned pup was found at Point Reyes National Seashore at Wildcat Beach in 2003. This species is separated into three recognized stocks based on three geographic regions (U.S. stock, Western Baja stock, and the Gulf of California stock; Lowry et al. 1992). Some movement has been documented between these geographic stocks, but rookeries in the U.S. are widely separated from major rookeries of western Baja California, Mexico (Barlow et al. 1995). Commercial harvest of the species in southern California and Mexico reduced the population to approximately 1,500 individuals by the 1920s. Since the passage of the Marine Mammal Protection Act in 1972, the California sea lion population has steadily increased along the West Coast of the United States (Carretta et al. 2002). The California sea lion has the largest population of any sea lion species and is the only sea lion whose population is showing a healthy growth rate of 5% to 6.2% per annum. Net production between 1980 and 2001 averaged 15.1%. Annual incidental takes in fisheries is approximately 915 individuals; however, the population is growing by 8.2% per year and fishing mortality is declining (Barlow et al. 1995). Current U.S. population estimates range from 237,000 and 244,000 (NOAA Stock Assessment 2003), and an additional 44,000 to 53,000 animals in Mexico (Aurioles-Gamboa and Zavala-Gonzalez 1994).

California sea lions give birth in May through July and breeding occurs in July and August. Females and pups are resident at breeding colonies year round and males migrate north to feeding areas from central California to British Columbia, Canada.

During years of low food availability (ENSO), females and juveniles may also migrate north in search of prey, and in some particularly poor years (1997-1998), there can be mass mortality of pups at rookeries.

On the Farallon Islands California Sea Lions haul out in many intertidal areas year round, fluctuating from several hundred to several thousand animals. Breeding animals are concentrated in areas where researchers do not visit (PRBO unpublished data).

California sea lions at Point Reyes haul out at only a couple locations, but individuals will haul out through out the area and will occur on human structures such as boat ramps. The annual population averages around 300-500 during the fall through spring months, but on occasion, several thousand sea lions can arrive depending upon local prey resources (S. Allen, unpublished data).

Steller Sea Lion: Steller sea lions breed from the Kuril Islands and Okhotsk Sea through the Aleutian Islands and the Gulf of Alaska, and south to central California (Merrick et al. 1987). The Steller sea lion was hunted during the sealing era for fur, hides, blubber, and other organs. More recently, Steller sea lions were harvested during a modern pup hunt that lasted from 1959-1972 in which approximately 45,000 pups were taken (Pasquel and Adkison 1994). At the cessation of the modern commercial hunting the Steller sea lion was found along the Pacific Rim from California to Japan with approximately 70% of the population in Alaskan waters. Two separate populations are recognized within US waters: an eastern population that includes animals east of Cape Suckling, Alaska (144° W), and a western population that includes animals west of Cape Suckling. Despite the cessation of the commercial hunt, the Steller sea lion population has experienced a rapid decrease since the mid-1980s with the western population declining by >64% in the last 30 years (Loughlin et al. 1992). The number in 1989 was estimated at 68,094 individuals. This total includes 10,000 in Russia, 47,960 in Alaska, 6,109 in British Columbia, 2,261 in Oregon, and 1,764 in California (Loughlin et al 1992). Numbers in Alaska have been declining by 7.8 % since 1994 (National Marine Mammal Laboratory 1995) and have declined by 3% in California (Le Boeuf et al. 1991, Ono 1993).

On Southeast Farallon Island, California, the abundance of females declined an average of 3.6% per year from 1974 to 1997 (Sydeman and Allen 1999). Pup counts at Año Nuevo declined 5% annually through the 1990s (NOAA Stock Assessment 2003), and have apparently stabilized between 2001 and 2005 (M. Lowry, SWFSC unpublished data). In 2000, the combined pup estimate for both islands was 349. In 2005, the pup estimate was 204 on Año Nuevo. Pup counts on the Farallon Islands have generally varied from 5-15 (Hastings and Sydeman 2002, PRBO unpublished data). Pups have not been born at Point Reyes Headland since the 1970s and Steller sea lions are seen in very low numbers there currently (S. Allen, unpubl. data).

Steller sea lions give birth in May through July and breeding occurs a couple of weeks after birth. Non-reproductive animals congregate at a few haul out sites, including at Año Nuevo and Point Reyes Headland. Pups are weaned during the winter and spring of the following year.

In 1990, the Steller sea lion was listed as a threatened species under the ESA, and the western stock was listed as endangered in 1997. In the 1960s and 70s the number of sea lions caught in trawl nets peaked, while present day numbers are low. California fisheries target several of the most important prey items for Steller sea lions and millions of metric tons of prey have been removed by fisheries in recent decades. Incidental mortality of Steller sea lions in fisheries was very low between 1990 and 2001 in California. Shooting of adults during fisheries interactions in central California have been documented by the Marine Mammal Stranding Network and one adult male was found shot at Point Reyes, California in the 1990s. In Alaska, there are also several processes that have been debated as contributing to the decline of the Steller sea lion population, including global climate change and killer whale predation (Springer et al. 2003).

On the Farallon and Año Nuevo Islands, Steller sea lion breeding colonies are located in closed areas where researchers never visited, eliminating any risk of disturbing breeding animals.

5. Type of Incidental Taking Authorization Requested

The proposed research and associated activities may result in “take by incidental harassment only” (Level B Harassment). An Incidental Harassment Authorization (IHA) is requested. All takes will involve incidental human presence near pinniped haulout areas.

6. Number of Marine Mammals that may be Affected

South Farallon Islands

M= Male, F= female, not preg = not pregnant, preg = pregnant

Species	Type of Research Affecting MM	# takes per year	# takes per animal per year	Pups M/F	Subadult M/F	Adult M/F(not preg)/F(preg)
Northern Elephant Seal	Seabird and Landings	100	1	0/0	30/30	20/20/10
Harbor Seal	Seabird and Landings	75	1	5/5	20/20	10/10/5
Steller Sea Lion	Seabird and Landings	25	1	0/0	10/10	3/2/0
Steller Sea Lion (Marine Mammal Research)	Northern Elephant Seal	16	1	0/0	7/7	1/1/0

California Sea Lion	Seabird and Landings	1500	1	0/0	500/500	300/200/0
---------------------	----------------------	------	---	-----	---------	-----------

Anõ Nuevo Island: (All takes at Ano Nuevo are due to seabird research)

Species	# takes per year	# takes per animal per year	Pups M/F	Subadult M/F	Adult M/F(not preg)/F(preg)
Northern Elephant Seal	100	1	0/0	30/30	20/10/10
Harbor Seal	30	1	3/3	5/5	5/5/4
Steller Sea Lion	10	1	0/0	4/2	3/1/0
California Sea Lion	1300	1	0/0	600/400	200/100/0

Point Reyes (All takes at Point Reyes are due to seabird and other non marine mammal research)

M= Male, F= female, not preg = not pregnant, preg = pregnant

Species	# takes per year	# takes per animal per year	Pups M/F	Subadult M/F	Adult M/F(not preg)/F(preg)
Northern Elephant Seal	300	1	0/0	170/100	20/10/0
Harbor Seal	100	1	0/0	25/25	25/25/0
California Sea Lion	500	1	0/0	100/100	250/50/0

7. Anticipated Impact of the Activity upon the Species or Stock

The only anticipated impacts would be temporary disturbances caused by pinnipeds observing humans on land. All measures will be taken to ensure that flushes do not result in a stampede of pinnipeds heading to the sea. From our experience, if extra caution is taken to conduct slow movements and stay close to the ground – a stampede is extremely rare. This might alter behaviors and cause animals to flush from the area. Animals may return to the same site once researchers have left or go to an alternate haul out site, which usually occurs within 30 minutes (Allen et al. 1985). Long term affects of this take are likely minimal as very few breeding animals will be present in areas where takes may occur.

It is expected that any incidental disturbance to pinnipeds from both types of research would have minimal, short-term effects and no long-term effects on the individuals. Incidental disturbance is believed to have minimal impacts because pinnipeds usually

return to a site or a nearby site within 30 minutes upon conclusion of research activities (Allen et al. 1985). Numerous Incidental Harassment Authorizations and Letters of Authorizations under the MMPA, Incidental Take Permits under Section 10(a)(1)(b) of the ESA, issued by NMFS (e.g. 72 FR 124), and reports on more localized areas (e.g., Demarchi and Bentley 2004) have analyzed the potential effects of incidental disturbance to pinnipeds from various sources. Based on these reports, the effects to pinnipeds appear, at the most, to displace the animals temporarily from their haul out sites. Based on previous research reports from PRBO, maximum disturbance to Steller sea lions would result in the animals flushing into the water in response to presence of the researchers. It is not expected that pinnipeds would permanently abandon a haul-out site during PRBO's research, as precautions would be taken to not disturb the same haul-out site on frequent occasions. No research would occur on pinniped rookeries; therefore, mother and pup separation or crushing of pups is not a concern. Harassment may occur as researchers approach the haul out sites with vessels and during capture and sampling activities. In PRBO's final report of activities conducted from 2000-2005 under permit No. 373-1575, they reported disturbing less than 16 Steller sea lions during all elephant seal surveys on West End Island in the Farallones. Reports are not due for bird research until after the current IHA expires but, to date, ~700 seals and sea lions have been harassed from bird research at all sites. In general, the minimization measures taken by the staff are effective. No other disturbance to Stellers was reported from any other location.

8. Anticipated Impact on Subsistence Uses

As there is no subsistence hunting of the stocks in question, no impacts on subsistence uses are expected as the result of the proposed project.

9. Anticipated Impact of the Activity upon the Habitat of the Marine Mammal Populations, and the Likelihood of Restoration of the Affected Habitat

Incidental marine mammal takes will not result in the physical altering of marine mammal habitat. No survey or sampling equipment will be left in habitat areas, no toxic chemicals will be present, and all state and federal marine regulations, including those from National Marine Sanctuaries, will be followed in regards to boat emissions. No major breeding habitat will be affected by this activity.

10. Anticipated Impact of the Loss or Modification of Habitat

There is no anticipated loss or modification of habitat.

11. Impact Minimization/Mitigation Methods

Seabird Research

All possible measures will be taken to reduce marine mammal disturbance for the activities described above. Actions are conducted as quickly and unobtrusively as possible, to reduce the length of time on haul outs and number of animals disturbed.

Researchers keep a low, hunched profiles, speak softly, and generally minimize actions that might startle non-target species. Seabird observations at North Landing on SEFI will be conducted in an observation blind where researchers are shielded from the view of hauled out pinnipeds. Transit adjacent to Stellers Sea Lion colonies on West End island will be done in a low crouch or crawl to avoid detection by animals. Beach landings on ANI will only occur after any pinnipeds that might be present on the landing beach have entered the water. ANI researchers accessing seabird nest boxes will crawl slowly if pinnipeds are within view. These activities and takes are not in breeding areas and reproductive animals will likely not be affected.

If seals need to be moved away from human facilities at Point Reyes, the “minimum tool” will be used. For example, waving a blue tarp at 20-30 feet distance or holding plywood boards and walking towards elephant seals can cause elephant seals to move. Physical touch is not required to cause elephant seals to move away from structures and would not occur.

Pinniped Research

Pinniped researches will keep maximum distance from survey animals in an effort to reduce disturbance. In addition, researchers will keep a low, hunched profiles, speak softly, and generally minimize actions that might startle non-target species. All efforts will be made to have no interaction – and keep out of visual site – of Steller’s Sea Lions when transiting to WEI for elephant seal surveys.

PRBO will suspend the research operations immediately if a dead or injured marine mammal is found in the vicinity of the project area and the death or injury of the animal could be attributable to the research activities.

12. Arctic Subsistence Uses, Plan of Cooperation

Not applicable. No activities will take place in or near a traditional Arctic subsistence hunting area.

13. Monitoring and Reporting

Researchers will collect information on number and species of animals flushed, and number of flushing events. These will be reported annually to NMFS.

Researchers shall take notes of sea lions and seals observed within the proposed research area during studies. The notes shall provide dates, time, tidal height, species, numbers of sea lions and seals present, and any behavior changes.

PRBO shall submit a final report, including these notes, to NMFS within 90 days after the expiration of the IHA.

14. Coordinating Research to Reduce and Evaluate Incidental Take

Visits to intertidal areas of SEFI during research activities will be coordinated to reduce potential take. PRBO is the sole research organization on the SFI, but all efforts will be made to minimize unnecessary trips to areas where pinnipeds may be encountered i.e. completing seabird observations in North Landing in one session and conducting seabird and pinniped research simultaneously. All research goals on ANI will be coordinated internally to accomplish all work goals in a single day and to minimize the necessary number of trips to the island. Once on ANI, researchers will coordinate monitoring schedules so areas near any pinnipeds will be accessed only once per visit. The lead biologist will always serve as an observer to evaluate incidental take and halt any research activities should the potential for incidental take be too great.

15. References

- Allen, S.G., D.G. Ainley, G.W. Page, C.A. Ribic. 1985. The effect of disturbance on harbor seal haul out patterns at Bolinas Lagoon, California. *Fishery Bulletin*. 82: 493-500.
- Auriolos-Gamboa, D., and A. Zavala-Gonzalez. 1994. Ecological factors that determine distribution and abundance of the California sea lion *Zalophus californianus* in the Gulf of California. *Ciencias Marinas* 20:535-553.
- Barlow, J., P. Boveng, M. S. Lowry, B. S. Stewart, B. J. Le Boeuf, W. J. Sydeman, R. J. Jameson, S. G. Allen, and G.W. Oliver. 1993. Status of the northern elephant seal population along the U.S. west coast in 1992. Admin. Rept. LJ-93-01. Southwest Fisheries Science Center, National Marine Fisheries Service, P.O. Box 271, La Jolla, CA. 32 pp.
- Barlow, J., R.L. Brownell, Jr., D.P. DeMaster, K.A. Forney, M.S. Lowry, S. Osmeck, T.J. Ragen, R.R. Reeves, R.J. Small. 1995. U.S. Pacific marine mammal stock assessments: 1995. NOAA Tech. Mem. NMFS 219. 162 pp.
- Barlow, J., P.S. Hill, K.A. Forney, D.P. DeMaster. 1998. U.S. Pacific marine mammal stock assessments: 1998. NOAA Tech. Mem. NMFS 258. 40 pp.
- Boveng, P. 1988. Status of the northern elephant seal population on the U.S. West Coast. Admin. Rep. LJ-88-05 Southwest Fisheries Science Center, National Marine Fisheries Service, P.O. Box 271, La Jolla, CA. 35pp.
- Brown, R.F., B.E. Wright, S.D. Riemer. 2005. Trends in abundance and current status of harbor seals, *Phoca vitulina*, in Oregon 1977-2003. *Marine Mammal Science* 21:657-670.
- Carretta, J. V., J. Barlow, K. A. Forney, M. M. Muto and J. Baker, editors. 2001. U.S. Pacific Marine Mammal Stock Assessments: 2001. NOAA NMFS Dept of Commerce, La Jolla, CA.

Carretta, J. V., M. M. Muto, J. Barlow, J. Baker, K. A. Forney, and M. Lowry, editors. 2002. U.S. Pacific Marine Mammal Stock Assessments: 2001. NOAA NMFS Dept of Commerce, La Jolla, CA.

Hanan, D.A., and M.J. Beeson. 1994. Harbor seal, Phoca vitulina richardsi, census in California, May-June, 1993. Final Rept. to NOAA, NMFS, 501 W. Ocean Blvd., Suite 4200, Long Beach, CA 90802.

Hastings, K.K. and W.J. Sydeman. 2002. Population status, seasonal variation, and long-term population trends of Stellers Sea Lion at the South Farallon Islands, California. *Fisheries Bulletin* 100:51-62.

Hoelzel, A. R. 1999. Impact of population bottlenecks on genetic variation and the importance of Life- history; A case study of the northern elephant seal. *Biological Journal of the Linnaean Society* 68:23-39.

Hoelzel, A. R., J. Halley, S. J. O'Brien, C. Campagna, T. Arnbom, B. Le Boeuf, K. Ralls, and G.A. Dover. 1993. Elephant seal genetic variation and the use of simulation models to investigate historical population bottlenecks. *Journal of Heredity* 84:443-449.

Huber, H.R., A.C. Rovetta, L.A. Fry, and S. Johnston. 1991. Age specific natality of northern elephant seals at the South Farallon Islands, California. *J. Mammalogy*. 72: 525-534.

Jeffries, S.J., R.F. Brown, H.R. Huber, and R.L. DeLong. 1997. Assessment of harbor seals in Washington and Oregon 1996. Annual report to the MMPA Assessment Program, Office of Protected Resources, NMFS, NOAA, 1335 East-West Highway, Silver Spring MD 20910. Available at National Marine Mammal Laboratory, 7600 Sand Point Way NE, Seattle, WA 98115.

Lamont, M.M., Vida, J.T., Harvey, J.T., Jeffries, S., Brown, R., Huber, H.H., DeLong, R., and W.K. Thomas. 1996. Genetic substructure of the Pacific Harbor Seal (Phoca vitulina richardsi) off Washington, Oregon, and California. *Marine Mammal Science* 12:402-413.

Le Boeuf, B.J., K. Ono, and J. Reiter. 1991. History of the Stellar sea lion population at Año Nuevo Island, 1961-1991. Administrative Report LJ-91-45C. National Marine Fisheries Service, Southwest Fisheries Science Center, La Jolla. 9 pp. Available from Southwest Fisheries Science Center, P.O. Box 271, La Jolla, Ca 92038.



Loughlin, T.R., A.S. Perlov, and V.A. Vladimirov. 1992. Range-wide survey and estimation of total number of Stellar sea lions in 1989. *Mar. Mam. Sci.* 8:220-239.

Lowry, M.S., J.V. Carretta, and K.A. Forney. 2005. Pacific harbor seal, Phoca vitulina richardsi, census in California during May - July 2004. Administrative Report LJ-05-06, available from Southwest Fisheries Science Center, 8604 La Jolla Shores Drive, La Jolla, CA 92037. 38 p

- Manna, J., D. Press, D. Roberts and S. Allen. 2006. Harbor Seal Monitoring: Point Reyes National Seashore and Golden Gate National Recreation Area. Annual Report of the National Park Service, San Francisco Bay Area Network. 22pp.
- McCann, T.S. 1985. Size, status and demography of southern elephant seals (Mirounga leonina) populations. In J.K. Ling and M.M. Bryden (eds.), Studies of Sea Mammals in South Latitudes. South Australian Museum. 132 pp.
- Merrick, R.L., Loughlin, T.R., and D.G. Calkins. 1987. Decline in abundance of the northern sea lion, Eumetopias jubatus, in Alaska, 1956-86. Fishery Bulletin 85: 351-365.
- National Marine Mammal Laboratory. 1995. Status review of the United States Steller sea lion (Eumetopias jubatus) population. National Marine Fisheries Service, Seattle, WA. 61 pp. Available at National Marine Mammal Laboratory, 7600 Sand Point Way NE, Seattle.
- Ono, K.A. 1993. Stellar sea lion research at Año Nuevo Island, California, during the 1992 breeding season. Administrative Report LJ-93-21C. National Marine Fisheries Service, Southwest Fisheries Science Center, La Jolla. 9 pp. Available from Southwest Fisheries Science Center, P.O. Box 271, La Jolla, Ca 92038.
- Pasqual, M.A., M.D. Adkison. 1994. The decline of the Steller sea lion in the northeast pacific: demography, harvest or environment. Ecol. Applications, 4(2) 393-403.
- Springer AM, J.A. Estes, G.B. van Vliet, T.M. Williams, D.F. Doak, E.M. Danner, K.A. Forney, and B. Pfister. 2003. Sequential megafaunal collapse in the North Pacific Ocean: An ongoing legacy of industrial whaling? Proceedings of the National Academy of Sciences of the United States of America 100 (21) 12223-12228.
- Stewart, B., B. Le Boeuf, P. Yochem, H. Huber, R. DeLong, R. Jameson, W. Sydeman, and S. Allen. 1994. History and present status of the northern elephant seal population. In: B.J. Le Boeuf and R.M. Laws (eds.) Elephant seals. Univ. Calif. Press, Berkeley. 414 pp.
- Sydeman, W.J., and S.G. Allen. 1999. Pinniped population dynamics in central California: correlations with sea surface temperature and upwelling indices. Marine Mammal Science. 15: 446-461.

16. Figures

Fig 1. Map of SEFI with pinniped haulout areas and location of North and East Landing West End Island







Fig 2. Map of ANI with pinniped haulout areas, Steller sea lion breeding area, and location of researcher trails and landing beach.

