


## BOTTLENOSE DOLPHIN (*Tursiops truncatus*): Western North Atlantic Coastal Stock

### STOCK DEFINITION AND GEOGRAPHIC RANGE

There are two hematologically and morphologically distinct bottlenose dolphin ecotypes (Duffield et al. 1983; Duffield 1986) which correspond to a shallow, warm water ecotype and a deep, cold water ecotype; both ecotypes have been shown to inhabit waters in the western North Atlantic Ocean (Hersh and Duffield 1990). Bottlenose dolphins which had stranded alive in the western North Atlantic in areas with direct access to deep oceanic waters had hemoglobin profiles matching that of the deep, cold water ecotype (Hersh and Duffield 1990). Hersh and Duffield (1990) also described morphological differences between the deep, cold water ecotype dolphins and dolphins with hematological profiles matching the shallow, warm water ecotype which had stranded in the Indian/Banana River in Florida. Because of their occurrence in shallow, relatively warm waters along the U.S. Atlantic coast and because their morphological characteristics are similar to the shallow, warm water ecotype described by Hersh and Duffield (1990), the Atlantic coastal bottlenose dolphin stock is believed to consist of this ecotype. There are currently insufficient data to allow separation of locally resident bottlenose dolphins (such as those from the Indian/Banana River) from the coastal stock in the western North Atlantic.

The structure of the coastal bottlenose dolphin stock in the western North Atlantic is uncertain, but what is known about it suggests that the structure is complex. A portion of the coastal stock migrates north of Cape Hatteras, North Carolina, to New Jersey during the summer (Scott et al. 1988). It has been suggested that this stock is restricted to waters < 25 m in depth within the northern portion of its range (Kenney 1990) because of an apparent disjunct distribution of bottlenose dolphins centered on the 25 m isobath which was observed during surveys of the region (CeTAP 1980). The lowest density of bottlenose dolphins was observed over the continental shelf, with higher densities along the coast and near the continental shelf edge. The coastal stock is believed to reside south of Cape Hatteras in the late winter (Mead 1975; Kenney 1990); however, the depth distribution of the stock south of Cape Hatteras is uncertain and the coastal and offshore stocks may overlap there. There was no apparent longitudinal discontinuity in bottlenose dolphin herd sightings during aerial surveys south of Cape Hatteras in the winter (Blaylock and Hoggard 1994).

Scott et al. (1988) hypothesized a single coastal migratory stock ranging seasonally from as far north as Long Island, NY, to as far south as central Florida, citing stranding patterns during a high mortality event in 1987-88 and observed density patterns along the U.S. Atlantic coast. Figure 1 illustrates the distribution of 584 bottlenose dolphin herd sightings during aerial surveys from shore to approximately 9 km past the Gulf Stream edge south of Cape Hatteras in the winter in 1992 (Blaylock and Hoggard 1994), from shore seaward to the 25 m isobath during the summer north of Cape Hatteras in 1994 (Blaylock 1995), and within one km of the shore from New Jersey to mid-Florida during three coastal surveys conducted during the summer in 1994 (Blaylock 1995). The proportion of the



**Figure 1.** Sightings of bottlenose dolphins during aerial surveys to the 25 m isobath north of Cape Hatteras during summer 1994, 9 km past the eastern Gulf Stream wall south of Cape Hatteras during winter 1991, and three coastal surveys within one km of shore from New Jersey to mid-Florida during the summer in 1994.

sightings illustrated which might be of bottlenose dolphins from other than the coastal stock is unknown; however, it is reasonable to assume that the coastal surveys within one km of shore minimized inclusion of the offshore stock.

A working hypothesis for the coastal bottlenose dolphin stock structure postulates that there are local, resident stocks in certain embayments and that transient stocks migrate seasonally into and out of these embayments (Scott et al. 1988). In the Indian-Banana River, 28 of 36 marked bottlenose dolphins either resided in or returned to the river system for a period of at least ten years (Odell and Asper 1990). Eight of the marked dolphins were never positively resighted. None of the marked dolphins were reported from outside the river system; however, search outside of the river system was limited. If the working hypothesis is correct, exchange between resident and transient components of the coastal stock could be sufficient to mask any genetic indicators of stock distinction, even though the stock components might be sufficiently distinct to respond differently to population pressures.

## **POPULATION SIZE**

Mitchell (1975) estimated that the coastal bottlenose dolphin population which was exploited by a shore-based net fishery until 1925 (Mead 1975) was at least 13,748 bottlenose dolphins in the 1800s. Recent estimates of bottlenose dolphin abundance in the U.S. Atlantic coastal area were made from two types of aerial surveys. The first type was aerial survey using standard line transect sampling with perpendicular distance data analysis (Buckland et al. 1993) and the computer program DISTANCE (Laake et al. 1993). The alternate survey method consisted of a simple count of all bottlenose dolphins seen from aerial surveys within one km of shore.

An aerial line-transect survey conducted during February-March 1991 in the coastal area south of Cape Hatteras. Sampling transects extended orthogonally from shore out to approximately 9 km past the western wall of the Gulf Stream into waters as deep as 140 m, and the area surveyed extended from Cape Hatteras to mid-Florida (Blaylock and Hoggard 1994). Systematic transects were placed randomly with respect to bottlenose dolphin distribution and approximately 3.3% of the total survey area of approximately 89,900 km<sup>2</sup> was visually searched. Survey transects, area, and dates were chosen utilizing the known winter distribution of the stocks in order to sample the entire coastal population; however, the offshore stock may represent some unknown proportion of the resulting population size estimates. Preliminary estimates of abundance were derived through the application of distance sampling analysis (Buckland et al. 1993) and the computer program DISTANCE (Laake et al. 1993) to the perpendicular distance sighting data. Bottlenose dolphin abundance was estimated to be 12,435 dolphins with coefficient of variation (CV) = 0.18 and the log-normal 95% confidence interval was 9,684-15,967 (Blaylock and Hoggard 1994).

Perpendicular sighting distance analysis (Buckland et al. 1983) of line transect data from an aerial survey throughout the northern portion of the range in July 1994, from Cape Hatteras to Sandy Hook, New Jersey, and from shore to the 25 m isobath, resulted in an abundance estimate of 25,841 bottlenose dolphins (CV = 0.40) (Blaylock 1995) within the approximately 25,600 km<sup>2</sup> area. These data were collected during a pilot study for designing future surveys and are considered to be preliminary in nature.

Either of the aerial line transect surveys and the resulting abundance estimates may have included dolphins from the offshore stock. It is not currently possible to distinguish the two bottlenose dolphin ecotypes during visual aerial surveys and the distribution of the two ecotypes in U.S. Atlantic EEZ waters is uncertain. Additional research is needed to interpret the significance of the line transect survey results.

An aerial survey of the coastal waters within a one km strip along the shore from Sandy Hook to approximately Vero Beach, Florida, was also conducted during July 1994 (Blaylock 1995). Dolphins from the offshore stock are believed unlikely to occur in this area. Observers counted all bottlenose dolphins seen within the one km strip alongshore from Cape Hatteras to Sandy Hook (northern area) and within the one km strip alongshore south of Cape Hatteras to approximately Vero Beach (southern area). The average of three counts of bottlenose dolphins in the northern area was 927 dolphins (range = 303-1,667) and the average of three counts of bottlenose dolphins in the southern area was 630 dolphins (range = 497-815). The sum of the highest counts in both areas was 2,482 dolphins.

### **Minimum Population Estimate**

Reasonable assurance of a minimum population estimate was not provided by line transect surveys because the proportion of dolphins from the offshore stock which might have been observed is unknown. The minimum population size was therefore taken as the highest count of bottlenose dolphins within the one km strip from shore between Sandy Hook and Vero Beach obtained during the July 1994 survey. The maximum count within one km of

shore between Sandy Hook and Cape Hatteras was 1,667 bottlenose dolphins and it was 815 bottlenose dolphins within one km of shore between Cape Hatteras and Vero Beach. The resulting minimum population size estimate for the western North Atlantic coastal bottlenose dolphin stock is 2,482 dolphins.

### **Current Population Trend**

Kenney (1990) reported an estimated 400-700 bottlenose dolphins from the inshore strata of aerial surveys conducted along the U.S. Atlantic coast north of Cape Hatteras in the summer during 1979-1981. These estimates resulted from line transect analyses; thus, they cannot be used in comparison with the direct count data collected in 1994 to assess population trends.

There was no significant difference in bottlenose dolphin abundance estimated from aerial line transect surveys conducted south of Cape Hatteras in the winter of 1983 and the winter of 1991 using comparable survey designs (NMFS unpublished data; Blaylock and Hoggard 1994) in spite of the 1987-88 mortality incident during which it was estimated that the coastal migratory population may have been reduced by up to 53% (Scott et al. 1988).

### **CURRENT AND MAXIMUM NET PRODUCTIVITY RATES**

Current and maximum net productivity rates are not known for this stock. The maximum net productivity rate was assumed to be 0.04 for purposes of this assessment. This value is based on theoretical calculations showing that cetacean populations may not generally grow at rates much greater than 4% given the constraints of their reproductive life history (Reilly and Barlow 1986).

### **POTENTIAL BIOLOGICAL REMOVAL**

Potential biological removal (PBR) has been specified as the product of the minimum population size, one-half the maximum productivity rate, and a "recovery" factor for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population size (OSP) (Anon. 1994). The recovery factor was set at 0.50 because of the stock is listed as depleted under the Marine Mammal Protection Act. PBR for the U.S. Atlantic coastal bottlenose dolphin stock is 25 dolphins.

### **ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY**

Four bottlenose dolphins were reported by the Northeast Region Stranding network to have shown signs of entanglement with fishing gear in Virginia in 1992 (K. Thounhurst, NMFS, personal communication, 1994). One of these was associated with pound net gear. In 1993, eight bottlenose dolphins in Virginia and one in Maryland were reported as entangled in fishing gear, but the gear type was not reported (NMFS unpublished data). Signs of interaction with fisheries (entanglement, net marks, missing appendages) were present in 22% of the bottlenose dolphin strandings investigated in North Carolina in 1993 (NMFS unpublished data).

The percentage of total mortality represented by stranded dolphins is unknown, but 20 bottlenose dolphin mortalities which showed signs of fishery interaction were reported in the Atlantic states of the NMFS Southeast Region in 1993 (NMFS unpublished data). A total of 29 bottlenose dolphins from the U.S. Atlantic coastal stock in the combined 1993 stranding records from both of the NMFS regions were reported to have shown indications of some sort of fishery interaction (NMFS unpublished data). It is unclear whether the interactions contributed to the mortalities or occurred post-mortem. Examination of marine mammal stranding records from the NMFS Southeast Region collected during 1988-1993 showed that an average of 21 (CV = 0.30) stranded bottlenose dolphins from the area including North Carolina to the Florida Keys were discovered annually showing signs of human interaction ranging from net marks and entanglement to gunshot wounds and boat propeller strikes (Southeast U.S. Marine Mammal Stranding Network unpublished data).

Stranding data probably underestimate the extent of fishery-related mortality and serious injury because not all of the dolphins which die or are seriously injured may wash ashore, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery-interaction. Another factor complicating the estimation of fishery-related mortality and serious injury to this stock using stranding data is the fact that an unknown proportion of stranded dolphins are from the offshore stock. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interaction.

The total fishery-related mortality and serious injury for this stock is not less than 10% of the calculated PBR and, therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. This

determination cannot be made for specific fisheries until the implementing regulations for Section 118 of the MMPA have been reviewed by the public and finalized.

### Fishery Information

Menhaden purse seiners have reported an annual incidental take of one to five bottlenose dolphins (NMFS 1991, pp. 5-73). Observer data are not available. The Atlantic menhaden purse seine fishery targets the Atlantic menhaden, *Brevortia tyrannus*, in Atlantic coastal waters approximately 3-18 m in depth. Twenty-two vessels operate off northern Florida to New England from April-January (NMFS 1991, pp. 5-73).

Coastal gillnets operate in different seasons targeting different species in different states throughout the range of this stock. Most nets are staked close to shore, but some are allowed to drift, and nets range in length from 91 m to 914 m. A gillnet fishery for American shad, *Alosa sapidissima*, operates seasonally from Connecticut to Georgia, with nets being moved from coastal ocean waters into fresh water with the shad spawning migration (Read 1994). There has been no direct observation by NMFS of this particular fishery, which is conducted in state waters, but it is considered likely that a few bottlenose dolphins are taken in this fishery each year (Read 1994). The North Carolina sink gillnet fishery operates in October-May targeting weakfish, croaker, spot, bluefish, and dogfish. Another gillnet fishery along the North Carolina Outer Banks targets bluefish in January-March. Similar mixed-species gillnet fisheries, under state jurisdiction, operate seasonally along the coast from Florida to New Jersey, with the exclusion of Georgia. There are no estimates of bottlenose dolphin mortality or serious injury available for these fisheries. A rough estimate of the average total annual coastal gillnet fishing effort is given in Table I.

Observer coverage of the U.S. Atlantic coastal gillnet fishery was initiated by the NEFSC Sea Sampling program in July, 1993; and from July to December 1993, 20 trips were observed. From January to April 1994, 71 trips were observed. This fishery, which extends from North Carolina to New York, is actually a combination of small vessel fisheries that target a variety of fish species, some of which operate right off the beach. The number of vessels in this fishery is unknown, because records are held by both state and federal agencies, and have not, as of yet, been centralized and standardized. Percent coverage by the program is unknown, but it is believed to be very low. No bottlenose dolphins were taken in the observed trips.

The shrimp trawl fishery operates from North Carolina through northern Florida virtually year around, moving seasonally up and down the coast. Estimated total fishing effort is given in Table I. One bottlenose dolphin was recovered dead from a shrimp trawl in Georgia in 1995 (Southeast U.S. Marine Mammal Stranding Network unpublished data), but no bottlenose dolphin mortality or serious injury has been previously reported to NMFS.

A haul seine fishery operates along northern North Carolina beaches during the spring and fall targeting mullet, spot, sea trout, and bluefish. There has been no by-catch of marine mammals reported to NMFS.

**Table I.** Roughly estimated average annual fishing effort (number deployed) by gear type for U.S. Atlantic coastal fisheries from New Jersey to Key West, Florida, in 1992-1993, having the potential for causing serious injury or mortality to bottlenose dolphins (NMFS unpublished data).

Gear Type	Effort
Haul seines	222
Purse seines	11,962
Otter trawls, bottom	22,550
Otter trawls, midwater	70
Gillnets, anchored or staked	22,252
Gillnets, drift and runaround	11,792

### Other Mortality

The nearshore habitat occupied by this stock is adjacent to areas of high human population and in the northern portion of its range is highly industrialized. The blubber of stranded dolphins examined during the 1987-88 mortality event contained anthropogenic contaminants in levels among the highest recorded for a cetacean (Geraci 1989). There are no estimates of indirect human-caused mortality resulting from pollution or habitat degradation, but a recent assessment of the health of live-captured bottlenose dolphins from Matagorda Bay, Texas, associated high levels of certain chlorinated hydrocarbons with low health assessment scores (Reif et al., in preparation).

### STATUS OF STOCK

This stock is considered to be depleted relative to OSP and it is listed as depleted under the Marine Mammal Protection Act (MMPA). There are data suggesting that the population was at an historically high level immediately

prior to the 1987-88 mortality event (Keinath and Musick 1988); however, the 1987-88 anomalous mortality event was estimated to have decreased the population by as much as 53% (Scott et al. 1988). A comparison of historical and recent winter aerial survey data in the area south of Cape Hatteras found no statistically significant difference between population size estimates (Student's t-test,  $P > 0.10$ ), but these estimates may have included an unknown proportion of the offshore stock. Population trends cannot be determined due to insufficient data.

There are no observer data directly linking serious injury and mortality to fisheries, but the total number of bottlenose dolphins assumed from this stock which stranded showing signs of fishery or human-related mortality exceeded PBR in 1993.

The species is not listed as threatened or endangered under the Endangered Species Act, but because this stock is listed as depleted under the MMPA it is a strategic stock.

## REFERENCES

- Anon. 1994. Report of the PBR (Potential Biological Removal) workshop. June 27-29, 1994. NOAA, NMFS Southwest Fisheries Science Center, La Jolla, California, 13 pp. + Appendices.
- Blaylock, R. A. 1995. A pilot study to estimate abundance of the U.S. Atlantic coastal migratory bottlenose dolphin stock. NOAA Tech. Mem. NMFS-SEFSC-362, 9 pp.
- Blaylock, R. A. and W. Hoggard. 1994. Preliminary estimates of bottlenose dolphin abundance in southern U.S. Atlantic and Gulf of Mexico continental shelf waters. NOAA Tech. Mem. NMFS-SEFSC-356, 10 pp.
- Buckland, S. T., D. R. Anderson, K. P. Burnham, and J. L. Laake. 1993. Distance sampling: Estimating abundance of biological populations. Chapman & Hall, London, 446 pp.
- CeTAP (Cetacean and Turtle Assessment Program). 1982. "A Characterization of Marine Mammals and Turtles in the Mid- and North Atlantic Areas of the U.S. Outer Continental Shelf, Final Report", Contract AA551-CT8-48, U.S. NTIS PB83-215855, Bureau of Land Management, Washington, DC, 576 pp.
- Duffield, D. A. 1986. Investigation of genetic variability in stocks of the bottlenose dolphin (*Tursiops truncatus*). Final report to the NMFS/SEFC, Contract No. NA83-GA-00036, 53 pp.
- Duffield, D. A., Ridgway, S. H., and Cornell, L. H. 1983. Hematology distinguishes coastal and offshore forms of dolphins (*Tursiops*). Can. J. Zool. 61: 930-933.
- Geraci, J. R. 1989. Clinical investigation of the 1987-88 mass mortality of bottlenose dolphins along the U.S. central and south Atlantic coast. Final Report to National Marine Fisheries Service, U.S. Navy, Office of Naval Research, and Marine Mammal Commission, 63 pp.
- Hersh, S. L. and D. A. Duffield. 1990. Distinction between northwest Atlantic offshore and coastal bottlenose dolphins based on hemoglobin profile and morphometry. Pages 129-139 in S. Leatherwood and R. R. Reeves (editors), The bottlenose dolphin, Academic Press, San Diego, 653 pp.
- Keinath, J. A. and J. A. Musick. 1988. Population trends of the bottlenose dolphin (*Tursiops truncatus*) in Virginia. Final Contract Report, Contract No. 40-GENF-800564, Southeast Fisheries Science Center, Miami, FL, 36 pp.
- Kenney, R. D. 1990. Bottlenose dolphins off the northeastern United States. Pages 369-386 in S. Leatherwood and R. R. Reeves (editors), The bottlenose dolphin, Academic Press, San Diego, 653 pp.
- Laake, J. L., S. T. Buckland, D. R. Anderson, and K. P. Burnham. DISTANCE user's guide, V2.0. Colorado Cooperative Fish & Wildlife Research Unit, Colorado State University, Ft. Collins, Colorado, 72 pp.
- Mead, J. G. 1975. Preliminary report on the former net fisheries for *Tursiops truncatus* in the western North Atlantic. J. Fish. Res. Bd. Can. 32(7): 1155-1162.
- Mitchell, E. 1975. Porpoise, dolphin and small whale fisheries of the world. Status and problems. Int. Union Conserv. Natur. Resour. Monogr. 3: 1-129.
- NMFS. 1991. Proposed regime to govern the interactions between marine mammals and commercial fishing operations after October 1, 1993. Draft Environmental Impact Statement, June 1991.
- Odell, D. K. and E. D. Asper. 1990. Distribution and movements of freeze-branded bottlenose dolphins in the Indian and Banana Rivers, Florida. Pages 515-540 in S. Leatherwood and R. R. Reeves (editors), The bottlenose dolphin, Academic Press, San Diego, 653 pp.
- Read, A. J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the northwest Atlantic. Rep. Int. Whal. Commn. Special Issue 15: 133-147.

- Reif, J. S., L. J. Hansen, S. Galloway, G. Mitchum, T. L. Schmitt. In preparation. The relationship between chlorinated hydrocarbon contaminants and selected health parameters in bottlenose dolphins (*Tursiops truncatus*) from Matagorda Bay, Texas, 1992. Colorado State University, Fort Collins, and NMFS, Southeast Fisheries Science Center, Miami, Florida.
- Reilly, S. B. and J. Barlow. 1986. Rates of increase in dolphin population size. Fish. Bull., U.S. 84(3): 527-533.
- Scott, G. P., D. M. Burn, and L. J. Hansen. 1988. The dolphin dieoff: long term effects and recovery of the population. Proceedings: Oceans '88, IEEE Cat. No. 88-CH2585-8, Vol. 3: 819-823.