

MINKE WHALE (*Balaenoptera acutorostrata*): Canadian East Coast Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Minke whales have a cosmopolitan distribution, being distributed in polar, temperate and tropical waters. In the North Atlantic, there are four recognized populations — Canadian East Coast, west Greenland, central North Atlantic, and northeastern North Atlantic (Donovan 1991). These divisions were defined by examining segregation by sex and length, catch distributions, sightings, marking data and pre-existing ICES boundaries. However, there were very few data from the Canadian East Coast population.

Minke whales off the eastern coast of the United States are considered to be part of the Canadian East Coast stock, which inhabits the area from the western half of the Davis Strait (45°W) to the Gulf of Mexico. The relationship between this stock and the other three stocks is uncertain. It is also uncertain if there are separate stocks within the Canadian East Coast stock.

The minke whale is common and widely distributed within the U.S. Atlantic Exclusive Economic Zone (EEZ) (CETAP 1982). There appears to be a strong seasonal component to minke whale distribution. Spring and summer are times of relatively widespread and common occurrence, and when the whales are most abundant in New England waters. In New England waters during fall there are fewer minke whales, while during winter the species appears to be largely absent. Like most other baleen whales,

minke whales generally occupy the continental shelf proper, rather than the continental shelf edge region. Records summarized by Mitchell (1991) hint at a possible winter distribution in the West Indies, and in the mid-ocean south and east of Bermuda. As with several other cetacean species, the possibility of a deep-ocean component to the distribution of minke whales exists but remains unconfirmed.




Figure 1. Distribution of minke whale sightings from NEFSC and SEFSC shipboard and aerial surveys during the summers of 1998, 1999, 2002, 2004, 2006 and 2007. Isobaths are the 100-m, 1000-m and 4000-m depth contours.

POPULATION SIZE

The total number of minke whales in the Canadian East Coast population is unknown. However, eleven estimates are available for portions of the habitat (see Appendix IV for details on these surveys and estimates). The best available current abundance estimate for minke whales, 3,312 (CV=0.74), is obtained from the 2006 aerial survey because this survey is recent and covered the largest portion of the animal's habitat.

Earlier estimates

For earlier abundance estimates please see Appendix IV.

Recent surveys and abundance estimates

An abundance estimate of 756 (CV=0.90) minke whales was derived from an aerial survey conducted in August

2002 which covered 7,465 km of trackline over waters from the 1000 m depth contour on the southern edge of Georges Bank to Maine (Table 1). The value of $g(0)$ used for this estimation was derived from the pooled data of the 2002, 2004 and 2006 aerial surveys.

An abundance estimate of 600 (CV=0.61) minke whales was obtained from a line-transect sighting survey conducted during 12 June to 4 August 2004 by a ship and plane that surveyed 6,180 km of trackline from the 100 m depth contour on the southern Georges Bank to the lower Bay of Fundy. The Scotian shelf south of Nova Scotia was not surveyed (Table 1; Palka 2006). Shipboard data were collected using the two independent team line transect method and analyzed using the modified direct duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and $g(0)$, the probability of detecting a group on the track line. Aerial data were collected using the Hiby circle-back line transect method (Hiby 1999) and analyzed accounting for $g(0)$ and biases due to school size and other potential covariates (Palka 2005). The value of $g(0)$ used for this estimation was derived from the pooled data of the 2002, 2004 and 2006 aerial surveys.

An abundance estimate of 3,312 (CV=0.74) minke whales was generated from an aerial survey conducted in August 2006 which surveyed 10,676 km of trackline in the region from the 2000 m depth contour on the southern edge of Georges Bank to the upper Bay of Fundy and to the entrance of the Gulf of St. Lawrence. (Table 1; Palka pers. comm.). The value of $g(0)$ used for this estimation was derived from the pooled data of the 2002, 2004 and 2006 aerial surveys.

An abundance estimate of 3,242 (95%CI=2,051-4,845) minke whales was generated from the Canadian Trans North Atlantic Sighting Survey (TNASS) in July-August 2007. This aerial survey covered area from northern Labrador to the Scotian Shelf, providing full coverage of the Atlantic Canadian coast. Estimates from this survey have not yet been corrected for availability and perception biases (Lawson and Gosselin 2009).

Table 1. Summary of abundance estimates for the Canadian east coast stock of minke whales. Month, year, and area covered during each abundance survey, and resulting abundance estimate (N_{best}) and coefficient of variation (CV).			
Month/Year	Area	N_{best}	CV
Aug 2002	S. Gulf of Maine to Maine	756	0.90
Jun-Jul 2004	Gulf of Maine to lower Bay of Fundy	600	0.61
Aug 2006	S. Gulf of Maine to upper Bay of Fundy to Gulf of St. Lawrence	3,312	0.74
Jul-Aug 2007	N. Labrador to Scotian Shelf	3,242	

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for minke whales is 3,312 animals (CV=0.74). The minimum population estimate for the Canadian East Coast minke whale is 1,899 animals.

Current Population Trend

There are insufficient data to determine population trends for this species.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. Life history parameters that could be used to estimate net productivity are that females mature between 6-8 years of age, and pregnancy rates are approximately 0.86 to 0.93. Based on these parameters, the calving interval is between 1 and 2 years. Calves are probably born during October to March after 10 to 11 months gestation and nursing lasts for less than 6 months. Maximum ages are not known, but for Southern Hemisphere minke whales maximum age appears to be about 50 years (IWC 1991; Katona *et al.* 1993).

For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the

constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a “recovery” factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is 1,899. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor, which accounts for endangered, depleted, or threatened stocks, or stocks of unknown status, relative to optimum sustainable population (OSP) is assumed to be 0.5 because this stock is of unknown status. PBR for the Canadian east coast minke whale is 19.

ANNUAL HUMAN-CAUSED MORTALITY AND INJURY

Recent minke whale takes have been observed in—or have been attributed to—the Northeast bottom trawl, Northeast/Mid-Atlantic lobster trap/pot, and unknown fisheries, although not all takes have resulted in mortalities (Tables 2 to 6).

Data to estimate the mortality and serious injury of minke whales come from the Northeast Fisheries Science Center Observer Program and from records of strandings and entanglements in U.S. waters. For the purposes of this report, only those strandings and entanglement records considered confirmed human-caused mortalities or serious injuries are shown in Tables 3 through 5.

During 2003 to 2007, the U.S. total annual estimated average human-caused mortality was 2.4 minke whales per year (CV=unknown). This is derived from three components: 0.2 minke whales per year from U.S. fisheries using observer data, 1.8 minke whales per year (unknown CV) from U.S. fisheries using strandings and entanglement data, and 0.4 minke whales per year from ship strikes (Glass *et al.* 2009). Detected mortalities should not be considered an unbiased representation of human-caused mortality. Detections are haphazard and not the result of a designed sampling scheme. As such they represent a minimum estimate of human-caused mortality which is almost certainly biased low.

Fishery Information

Detailed fishery information is reported in Appendix III.

Earlier Interactions

Little information is available about fishery interactions that took place before the 1990s. Read (1994) reported that a minke whale was found dead in a Rhode Island fish trap in 1976. A minke whale was caught and released alive in the Japanese tuna longline fishery in 3,000 m of water, south of Lydonia Canyon on Georges Bank, in September 1986 (Waring *et al.* 1990).

Two minke whales were observed taken in the Northeast sink gillnet fishery between 1989 and the present. The take in July 1991, south of Penobscot Bay, Maine was a mortality, and the take in October 1992, off the coast of New Hampshire near Jeffreys Ledge, was released alive.

A minke whale was trapped and released alive from a herring weir off northern Maine in 1990.

Four minke whale mortalities were observed in the Atlantic pelagic drift gillnet fishery during 1995.

One minke whale was reported caught in an Atlantic tuna purse seine off Stellwagen Bank in 1991 (D. Beach, NMFS NE Regional Office, pers. comm.) and another in 1996. The minke caught during 1991 was released uninjured after a crew member cut the rope wrapped around the tail. The minke whale caught during 1996 escaped by diving beneath the net.

One minke whale, reported in the strandings and entanglement database maintained by the New England Aquarium and the Northeast Regional Office/NMFS, was taken in a 3.5-inch gill net on 24 June 1998 off Long Island, New York. This take was assigned to the mid-Atlantic gillnet fishery. No minke whales have been taken in this fishery during observed trips in 1993 to 2007.

U.S.

Northeast Bottom Trawl

The fishery is active in New England waters in all seasons. Detailed fishery information is reported in Appendix III. One freshly dead minke whale was caught in 2004 on the northeast tip of Georges Bank in US waters (Tables 2 and 5). An expanded bycatch estimate has not been generated because, with only one observed take, it is not possible to generate an accurate bycatch estimate. Therefore, this catch is reported as 1, with a resulting 5-year mean annual mortality of 0.2.

Northeast/Mid-Atlantic Lobster Trap/Pot Fishery

The strandings and entanglement database, maintained by the New England Aquarium and the Northeast Regional Office/NMFS, reported 7 minke whale mortalities and serious injuries that were attributed to the lobster fishery during 1990 to 1994; 1 in 1990 (may be a serious injury), 2 in 1991 (1 mortality and 1 serious injury), 2 in 1992 (both mortalities), 1 in 1993 (serious injury) and 1 in 1994 (mortality) (1997 List of Fisheries 62FR33, 2 January 1997). The one confirmed minke whale mortality during 1995 was attributed to the lobster fishery. No confirmed mortalities or serious injuries of minke whales occurred in 1996. From the four confirmed 1997 records, one minke whale mortality was attributed to the lobster trap fishery. In 2002, one minke whale mortality and one live release were attributed to this fishery. The 28 June 2003 mortality, while wrapped in lobster gear, cannot be confirmed to have become entangled in the area, and so is not attributed to the fishery (Table 5). Annual mortalities due to this fishery, as determined from strandings and entanglement records that have been audited, were 1 in 1991, 2 in 1992, 1 in 1994, 1 in 1995, 0 in 1996, 1 in 1997, 0 in 1998 to 2001, 1 in 2002, and 0 in 2003 through 2007. Estimated average annual mortality related to this fishery during 2003 to 2007 was 0 minke whales per year).

Unknown Fisheries

The strandings and entanglement database, maintained by the New England Aquarium and the Northeast Regional Office/NMFS, include 36 records of minke whales within U.S. waters for 1975-1992. The gear include unspecified fishing nets, unspecified cables or lines, fish traps, weirs, seines, gillnets, and lobster gear. One confirmed entanglement was an immature female minke whale, entangled with line around the tail stock, which came ashore on the Jacksonville, Florida jetty on 31 January 1990 (R. Bonde, USFWS, Gainesville, FL, pers. comm.).

The audited NE Regional Office/NMFS entanglement/stranding database contains records of minke whales, of which the confirmed mortalities and serious injuries from the last five years are reported in Table 5. Mortalities (and serious injuries) that were likely a result of a fishery interaction with an unknown fishery include 3 (0) in 1997, 3 (0) in 1999, 1 (1) in 2000, 2 (0) in 2001, 1 (0) in 2002, 5 (0) in 2003, 2 (0) in 2004, 0 (0) in 2005, 0 (0) in 2006, and 1 (1) in 2007 (Table 3). Examination of minke entanglement records from 1997 indicates that 4 out of 4 confirmed records of mortality were likely a result of fishery interactions. One was attributed to the lobster pot fishery (see above), and three were not attributed to any particular fishery because the information from the entanglement event did not contain the necessary details. Of the five mortalities in 1999, two were attributed to an unknown trawl fishery and three to some other fishery. Of the two interactions with an unknown fishery in 2000, one was a mortality and one was a serious injury. In 2001, the two confirmed fishery interactions were both from an unknown fishery. In 2002, there was one mortality in an unknown fishery. In 2003, 5 of 5 confirmed mortalities were due to interactions with an unknown fishery (Tables 3 and 5). In 2004, of the four confirmed mortalities, two were due to an interaction with an unknown fishery. In 2005 and 2006 there were no mortalities attributed to fishery interactions. In 2007 there was one mortality and one serious injury, both attributed to unknown fisheries.

In general, an entangled or stranded cetacean could be an animal that is part of an expanded bycatch estimate from an observed fishery and thus it is not possible to know if an entangled or stranded animal is an additional mortality. During 1997 through 2007, no minke whale bycatch estimates were generated from observed takes in any fishery observed by the NEFSC Observer Program, therefore, the strandings from those years in which mortalities were attributable to fishery interactions can be added into the human-caused mortality estimate. During 2003 to 2007, as determined from strandings and entanglement records, the estimated average annual mortality is 1.8 minke whales per year in unknown fisheries (Table 3).

CANADA

Read (1994) reported interactions between minke whales and gillnets in Newfoundland and Labrador, in cod traps in Newfoundland, and in herring weirs in the Bay of Fundy. Hooker *et al.* (1997) summarized bycatch data from a Canadian fisheries observer program that placed observers on all foreign fishing vessels operating in Canadian waters, on between 25% and 40% of large Canadian fishing vessels (greater than 100 feet long), and on approximately 5% of smaller Canadian fishing vessels. During 1991 through 1996, no minke whales were observed taken.

Herring Weirs

During 1980 to 1990, 15 of 17 minke whales were released alive from herring weirs in the Bay of Fundy. During January 1991 to September 2002, 26 minke whales were trapped in herring weirs in the Bay of Fundy. Of these 26, 1 died (H. Koopman, pers. comm.) and several (number unknown) were released alive and unharmed (A.

Westgate, pers. comm.).

Other Fisheries

Six minke whales were reported entangled during 1989 in the now non-operational groundfish gillnet fishery in Newfoundland and Labrador (Read 1994). One of these animals escaped and was still towing gear, the remaining five animals died.

Salmon gillnets in Canada, now no longer used, had taken a few minke whales. In Newfoundland in 1979, one minke whale died in a salmon net. In Newfoundland and Labrador, between 1979 and 1990, it was estimated that 15% of the Canadian minke whale takes were in salmon gillnets. A total of 124 minke whale interactions were documented in cod traps, groundfish gillnets, salmon gillnets, other gillnets, and other traps. The salmon gillnet fishery ended in 1993 as a result of an agreement between the fishermen and North Atlantic Salmon Fund (Read 1994).

Five minke whales were entrapped and died in Newfoundland cod traps during 1989. The cod trap fishery closed in Newfoundland in 1993 due to the depleted groundfish resources (Read 1994).

In 2005, four minke whales were reported entangled in fishing gear in Newfoundland and Labrador. Two (entangled in salmon net and mackerel trap gear) were released alive and two (involved with whelk pot and toad crab pot fisheries) were dead (Ledwell and Huntington 2006). Four minke whales were reported entangled in 2007 in Newfoundland and Labrador. All were released alive (Ledwell and Huntington 2007).

Table 2. Summary of the incidental mortality of minke whales (*Balaenoptera acutorostrata*) by commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the mortalities recorded by on-board observers (Observed Mortality), the estimated annual mortality (Estimated Mortality), the estimated CV of the annual mortality (Estimated CVs) and the mean annual mortality (CV in parentheses).

Fishery	Years	Vessels	Data Type ^a	Observer Coverage ^b	Observed Mortality	Estimated Mortality	Estimated CVs	Mean Annual Mortality
Northeast Bottom Trawl	03-07	unk	Obs. Data	.03, .04, .05, .06, .06	0, 1, 0, 0, 0	1 ^c	unk ^c	0.2 ^c
Total								0.2 ^c

a) Observer data (Obs. Data), used to measure bycatch rates, are collected within the Northeast Fisheries Science Center (NEFSC) Fisheries Observer Program.
 b) Observer coverage for trawl fishery is measured in trips.
 c) Analysis of bycatch mortality attributed to the Northeast bottom trawl fishery has not been generated, due to the small number of observed takes, so the single mortality reported by the fisheries observer program is counted as 1.

Table 3. From strandings and entanglement data, summary of confirmed incidental mortalities and serious injuries of minke whales (*Balaenoptera acutorostrata*) by commercial fishery: includes years sampled (Years), number of vessels active within the fishery (Vessels), type of data used (Data Type), mortalities and serious injuries assigned to this fishery (Assigned Mortality), and mean annual mortality and serious injuries. See Table 5 for details. (NA=Not Available)

Fishery	Years	Vessels	Data Type ^a	Assigned Mortality	Mean Annual Mortality
Unknown Fisheries	03-07	NA	Entanglement & Strandings	5, 2, 0, 0, 2	1.8
TOTAL					1.8 (CV=unk)

a. Data from records in the entanglement and strandings data base maintained by the New England Aquarium and the Northeast Regional Office/NMFS (Entanglement and Strandings).

Table 4. Summary of minke whales (<i>Balaenoptera acutorostrata</i>) released alive, by commercial fishery, years sampled (Years), ratio of observed mortalities recorded by on-board observers to the estimated mortality (Ratio), the number of observed animals released alive and injured (Injured), and the number of observed animals released alive and uninjured (Uninjured). (NA = Not Available)				
Fishery	Years	Ratio	Injured	Uninjured
Pelagic longline	03-07	0	0	1 ^a
a. Minke whale released alive from pelagic longline gear in 2003.				

Table 5. Confirmed U.S. and Canadian human-caused mortality and serious injury records of Canadian East Coast stock of minke whales, January 2003 through December 2007.						
Date ^a	Report Type ^b	Age, Sex, Length	Location ^a	Assigned Cause: P=primary, S=secondary		Notes/Observations
				Ship strike	Entang./ Fsh. Inter.	
05/24/03	mortality	Adult Male 7.6m	Gloucester, MA		P	Unknown fishery; line marks on head and dorsal fin; no line present; cut across back anterior to dorsal fin; no gear recovered
05/31/03	mortality	Juvenile Female 3.6m (est)	Martha's Vineyard, MA		P	Unknown fishery; whale stranded live wrapped in about 15 feet of 5.5 inch mesh netting, probably trawl gear
06/28/03	mortality	Yearling Male 5.1m	Chatham, MA		P	Unknown fishery; wrapped in lobster gear; gear not recovered
08/09/03	mortality	Juvenile Female, 3.5m (est)	Harwich, MA		P	Unknown fishery; hemorrhaging in areas with net marks on whale; no gear recovered
09/13/03	mortality	Juvenile Female 6m (est)	Casco Bay, ME		P	Unknown fishery; freshly dead; external chaffing marks and belly slit open; no gear recovered
05/06/04	mortality	Adult Female 7.7m	Martha's Vineyard, MA		P	Unknown fishery; constricting line marks on peduncle; indications of drowning from internal exam; no gear present
06/01/04	mortality	Juvenile Female 6.5m	Chatham, MA	P		Large area of subdermal hemorrhaging

07/19/04	mortality	Adult Female 7.9m	Eastham, MA		P	Unknown fishery; extensive entanglement markings; no gear recovered
08/04	mortality	age & sex unknown 4m (est)	Georges Bank		P	Northeast Bottom Trawl; fresh dead, rigid, had to cut out of net, rope in mouth
05/23/05	mortality	Juvenile Male 5.9m	Port Elizabeth, NJ	P		Ribs shattered; liver ruptured; evidence of internal hemorrhaging
07/16/07	serious injury	age & sex unknown 10m (est)	Trescott, ME		P	Unknown fishery; wrapped in gear and anchored; no gear recovered
08/05/07	mortality	Juvenile Female 4.3m	Cape Cod Bay, MA		P	Unknown fishery; chronic entanglement with severe emaciation and dehydration and loss of protein; line lacerated blubber layer across back and at flipper insertions; severe hemorrhage and necrosis of blubber at gear entanglement points; gear consists of 11/64" diameter rope
<p>a. The date sighted and location provided in the table are not necessarily when or where the serious injury or mortality occurred; rather, this information indicates when and where the whale was first reported beached, entangled, or injured.</p> <p>b. National guidelines for determining what constitutes a serious injury have not been finalized. Interim criteria as established by NERO/NMFS (Glass <i>et al.</i> 2009) have been used here. Some assignments may change as new information becomes available and/or when national standards are established.</p>						

Other Mortality

Minke whales have been and continue to be hunted in the North Atlantic. From the Canadian East Coast population, documented whaling occurred from 1948 to 1972 with a total kill of 1,103 animals (IWC 1992). Animals from other North Atlantic minke populations are presently still being harvested at low levels.

U.S.

Minke whales inhabit coastal waters during much of the year and are thus subject to collision with vessels. According to the NMFS/NER marine mammal entanglement and stranding database, on 7 July 1974, a necropsy of a minke whale suggested a vessel collision; on 15 March 1992, a juvenile female minke whale with propeller scars was found floating east of the St. Johns Channel entrance (R. Bonde, USFWS, Gainesville, FL, pers. comm.); and on 15 July 1996 the captain of a vessel reported hitting a minke whale offshore of Massachusetts. After reviewing this record, it was concluded the animal struck was not a serious injury or mortality. On 12 December 1998, a minke whale was struck and presumed killed by a whale watching vessel in Cape Cod Bay off Massachusetts.

During 1999 to 2003, no minke whale was confirmed struck by a ship. During 2004 and 2005, one minke whale mortality was attributed to ship strike in each year (Table 5). During 2006 and 2007, no minke whale was confirmed struck by a ship. Thus, during 2003 to 2007, as determined from stranding and entanglement records, the estimated annual average was 0.4 minke whales per year struck by ships.

In October 2003, an Unusual Mortality Event was declared involving minke whales and harbor seals along the coast of Maine. Two of the seven criteria established to designate such an event were met by these species. Specifically, there was a marked increase in mortalities when compared with historical records, and the mortalities

were occurring in a localized area of the Maine coast. From 11-30 September 2003, nine minke whales were reported along the mid-coast to southern Maine. Results from analyses for biotoxins failed to show the presence of either saxitoxin or domoic acid (by ELISA and Receptor Binding Assay). Most whale carcasses that were examined appeared to be in good body condition immediately prior to death. Since October 2003, the number of minke whale stranding reports has returned to normal. There were two minke whale stranding mortalities in NC in 2005 but in neither case could cause of death be attributed to human causes (Glass *et al.* 2008). There were 7 minke whale stranding mortalities reported along the US Atlantic coast in 2006. Three were in New Jersey, one in Massachusetts, one in Rhode Island, and two in the EEZ. One of the stranding mortalities from New Jersey was reported with signs of human interaction due to pieces of plastic found in the stomach.

CANADA

The Nova Scotia Stranding Network documented whales and dolphins stranded on the coast of Nova Scotia between 1991 and 1996 (Hooker *et al.* 1997). Researchers with the Department of Fisheries and Oceans, Canada documented strandings on the beaches of Sable Island (Lucas and Hooker 2000). Sable Island is approximately 170 km southeast of mainland Nova Scotia. Lucas and Hooker (2000) reported 4 minke whales stranded on Sable Island between 1970 and 1998, 1 in spring 1982, 1 in January 1992, and a mother/calf in December 1998. On the mainland of Nova Scotia, a total of 7 reported minke whales stranded during 1991 to 1996. The 1996 stranded minke whale was released alive off Cape Breton on the Atlantic Ocean side, the rest were found dead. All the minke whales stranded between July and October. One was from the Atlantic Ocean side of Cape Breton, 1 from Minas Basin, 1 was at an unknown location, and the rest stranded in the vicinity of Halifax, Nova Scotia. It is unknown how many of the strandings resulted from fishery interactions.

Whales and dolphins stranded between 1997 and 2007 on the coast of Nova Scotia as recorded by the Marine Animal Response Society (MARS) and the Nova Scotia Stranding Network are as follows (Table 6): 4 minke whales stranded in 1997 (1 in June and 3 in July), 0 documented strandings in 1998 to 2000, 1 in September 2001, 4 in 2002 (1 in July, 1 in August, and 2 in November), 2 in 2003 (1 in August and 1 in October), 0 in 2004, 3 in 2005 (1 in June and 2 in August), 8 in 2006 (1 in January, 2 in May, 1 in July, 1 in August, 1 in November (live) and 2 in December), and 1 in 2007 (October) (Table 6).

The Whale Release and Strandings program has reported seven minke whale stranding mortalities in Newfoundland and Labrador between 2003 and 2007 (Ledwell and Huntington 2001; 2002; 2003; 2004; 2006; 2007) (Table 6).

Area	YEAR					Total
	2003	2004	2005	2006	2007	
Nova Scotia	2	0	3	8	1	14
Newfoundland and Labrador	1	2	1	1	2	7
Total	3	2	4	9	3	21

STATUS OF STOCK

The status of minke whales, relative to OSP, in the U.S. Atlantic EEZ is unknown. The minke whale is not listed as endangered under the Endangered Species Act (ESA). The total U.S. fishery-related mortality and serious injury for this stock is not less than 10% of the calculated PBR and, therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. This is not a strategic stock because estimated human-related mortality and serious injury does not exceed PBR and the minke whale is not listed as a threatened or endangered species under the ESA.

REFERENCES CITED

- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.
- CETAP 1982. A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the U.S. outer continental shelf, final report. Washington, DC, Bureau of Land Management. #AA551-CT8-48 538 pp.
- Donovan, G.P. 1991. A review of IWC stock boundaries. Rep. Int. Whal. Comm. (Special Issue) 13: 39-68.
- Glass, A.H., T.V.N. Cole and M. Garron 2009. Mortality and serious injury determinations for baleen whale stocks along the United States eastern seaboard and adjacent Canadian Maritimes, 2003-2007. Northeast Fish Sci Cent Ref Doc. 09-04. 19 pp.
- Glass, A.H., T.V.N. Cole, M. Garron, R.L. Merrick and R.M. Pace, III 2008. Mortality and serious injury determinations for baleen whale stocks along the United States Eastern Seaboard and adjacent Canadian Maritimes, 2002-2006. Northeast Fish. Sci. Cent. Ref. Doc. 08-04. 18 pp.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. Pages 179-189 in: G.W. Garner, S.C. Amstrup, J.L. Laake *et al.*, (eds.) Marine Mammal Survey and Assessment Methods. Balkema, Rotterdam.
- Hooker, S.K., R.W. Baird and M.A. Showell 1997. Cetacean strandings and bycatches in Nova Scotia, Eastern Canada, 1991-1996. Meeting document SC/49/O5 submitted to the 1997 International Whaling Commission Scientific Committee meeting in Bournemouth, UK.
- IWC [International Whaling Commission] 1991. Appendix 11. Biological parameters of North Atlantic minke whales. In Annex F, report of the sub-committee on North Atlantic minke whales. Rep. Int. Whal. Comm. 41: 160.
- IWC [International Whaling Commission] 1992. Annex K. Report of the working group on North Atlantic minke whales. Rep. Int. Whal. Comm. 42: 246-251.
- Katona, S.K., V. Rough and D.T. Richardson 1993. A field guide to whales, porpoises, and seals from Cape Cod to Newfoundland. Smithsonian Institution Press, Washington, DC. 316 pp.
- Lawson, J.W. and J.-F. Gosselin 2009. Distribution and preliminary abundance estimates for cetaceans seen during Canada's Marine Megafauna Survey - A component of the 2007 TNASS. Can. Sci. Advisory Sec. Res. Doc. 2009/031. 28 pp.
- Ledwell, W. and J. Huntington 2001. Whale entrapments in fishing gear, strandings and sightings of marine animals and summary of the Entrapment Program during 2001 - Newfoundland region. Report to the Habitat Stewardship Program of Environment Canada. 3 pp.
- Ledwell, W. and J. Huntington 2002. Whale entrapments in fishing gear and summary of the Marine Animal Disentanglement Program in Newfoundland and Labrador during 2002. Report to the Habitat Stewardship Program of Environment Canada. 4 pp.
- Ledwell, W. and J. Huntington 2003. Whale entrapments in fishing gear reported during 2003 and a summary of the Whale Release and Strandings Program in Newfoundland and Labrador. Report to the Department of Fisheries and Oceans. St. John's, Newfoundland, Canada 3 pp.
- Ledwell, W. and J. Huntington 2004. Marine animal entrapments in fishing gear in Newfoundland and Labrador and a summary of the Whale Release and Strandings Program during 2004. Report to the Department of Fisheries and Oceans. St. John's, Newfoundland, Canada 3 pp.
- Ledwell, W. and J. Huntington 2006. Whale, leatherback sea turtles and basking shark entrapments in fishing gear in Newfoundland and Labrador and a summary of the Whale Release and Strandings Program during 2005. Report to the Department of Fisheries and Oceans. St. John's, Newfoundland, Canada 18 pp.
- Ledwell, W. and J. Huntington 2007. Whale and leatherback sea turtle entrapment in fishing gear in Newfoundland and Labrador and a summary of the Whale Release and Strandings Program during 2006. Report to the Department of Fisheries and Oceans. St. John's, Newfoundland, Canada 3 pp.
- Lucas, Z.N. and S.K. Hooker 2000. Cetacean strandings on Sable Island, Nova Scotia, 1970-1998. Can. Field-Nat. 114(1): 46-61.
- Mitchell, E.D. 1991. Winter records of the minke whale (*Balaenoptera acutorostrata* Lacepede 1804) in the southern North Atlantic. Rep. Int. Whal. Comm. 41: 455-457.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. Rep. Int. Whal. Comm. (Special Issue) 16: 27-50.

- Palka, D.L. 2005. Aerial surveys in the northwest Atlantic: estimation of $g(0)$, Proceedings of a Workshop on Estimation of $g(0)$ in Line-Transect Surveys of Cetaceans. European Cetacean Society's 18th Annual Conference; Kolmården, Sweden; Mar. 28, 2004.
- Palka, D.L. 2006. Summer abundance estimates of cetaceans in US North Atlantic Navy Operating Areas. Northeast Fish. Sci. Cent. Ref. Doc. 06-03. 41 pp.
<http://www.nefsc.noaa.gov/nefsc/publications/crd/crd0603/crd0603.pdf>
- Palka, D.L. and P.S. Hammond 2001. Accounting for responsive movement in line transect estimates of abundance. *Can. J. Fish. Aquat. Sci* 58: 777-787.
- Read, A.J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the northwest Atlantic. Pages 133-147 *in*: W.F. Perrin, G.P. Donovan and J. Barlow, (eds.) Gillnets and cetaceans. Reports of the International Whaling Commission. (Special Issue) 15.
- Wade, P.R. and R.P. Angliss 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Waring, G.T., P. Gerrior, P.M. Payne, B.L. Parry and J.R. Nicolas 1990. Incidental take of marine mammals in foreign fishery activities off the northeast United States, 1977-1988. *Fish. Bull.* 88(2): 347-360.

