

Table 186. Estimated Numbers of AIDS Diagnoses Among Adults and Adolescents, by Sex and Transmission Category: 2006 to 2009

[All data are provisional. Data are included for 50 states, the District of Columbia, Puerto Rico, the U.S. Virgin Islands, Guam, and the U.S. Pacific Islands. Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing risk-factor information, but not for incomplete reporting. Acquired immunodeficiency syndrome (AIDS) is a specific group of diseases or conditions which are indicative of severe immunosuppression related to infection with the human immunodeficiency virus (HIV). Data are subject to retrospective changes and may differ from those data in Table 184. For information on HIV death rates, go to Table 129]

Transmission category					Cumulative
	2006	2007	2008	2009	through 2009 ¹
Persons 13 years old and over, total²	36,987	36,213	35,512	34,980	1,132,836
Males, total	27,067	26,435	26,175	26,102	903,661
Male-to-male sexual contact	16,665	16,680	16,637	17,171	535,570
Injection drug use	4,126	3,744	3,554	3,207	199,565
Male-to-male sexual contact and injection drug use	1,994	1,841	1,729	1,608	79,693
Heterosexual contact ³	4,080	4,004	4,066	3,956	75,901
Other ⁴	202	167	190	159	12,931
Females, total	9,920	9,777	9,337	8,879	229,173
Injection drug use	2,553	2,453	2,192	1,982	90,102
Heterosexual contact ³	7,172	7,139	7,007	6,740	131,886
Other ⁴	195	185	137	157	7,185

¹ From the beginning of the epidemic through 2009. ² Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total. ³ Heterosexual contact with a person known to have, or to be at high risk for, HIV infection. ⁴ Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

Source: U.S. Centers for Disease Control and Prevention, Atlanta, GA, *HIV Surveillance Report, 2009*, Vol. 21, <<http://www.cdc.gov/hiv/surveillance/resources/reports/2009report>>.

Table 187. Estimated Numbers of Persons Living with an AIDS Diagnosis by Selected Characteristics: 2000 to 2008

[See headnote, Table 186]

Age and characteristic	2000	2005	2006	2007	2008
Total^{1, 2}	323,679	432,846	452,235	471,749	490,696
AGE AS OF END OF YEAR					
Less than 13 years old	2,954	1,506	1,211	942	720
13 and 14 years old	528	811	766	700	624
15 to 24 years old	5,158	7,806	8,281	8,873	9,521
25 to 34 years old	56,129	45,923	45,070	44,904	45,509
35 to 44 years old	146,206	164,551	161,547	156,302	149,198
45 to 54 years old	85,474	149,284	162,424	175,766	188,896
55 to 64 years old	21,717	50,376	58,267	67,048	76,362
65 years old and over	5,513	12,589	14,669	17,213	19,865
RACE/ETHNICITY					
American Indian/Alaska Native	1,035	1,506	1,571	1,635	1,733
Asian ³	2,293	3,777	4,095	4,472	4,883
Black/African American	131,826	183,083	191,618	200,281	209,175
Hispanic/Latino ⁴	66,271	90,836	95,532	100,344	104,791
Native Hawaiian/Other Pacific Islander	174	331	374	418	449
White	118,189	147,585	153,055	158,383	163,286
Multiple races	3,719	5,566	5,831	6,056	6,222
Transmission category:					
MALE ADULT/ADOLESCENT					
Males 13 years old and over, total	251,286	330,239	344,327	358,519	372,528
Male-to-male sexual contact	143,594	196,973	207,449	218,049	228,727
Injection drug use	58,437	65,366	66,042	66,741	67,287
Male-to-male sexual contact and injection drug use	25,635	30,289	30,831	31,368	31,822
Heterosexual contact ⁵	20,922	34,563	36,898	39,209	41,467
Other ⁶	2,698	3,048	3,106	3,151	3,224
FEMALE ADULT/ADOLESCENT					
Females 13 years old and over, total	68,423	98,500	103,819	109,164	114,123
Injection drug use	28,644	34,207	34,872	35,547	35,981
Heterosexual contact ⁵	38,269	62,252	66,803	71,364	75,831
Other ⁶	1,510	2,041	2,144	2,252	2,311
(CHILD LESS THAN 13 YEARS OLD AT DIAGNOSIS)					
Total	3,971	4,107	4,088	4,064	4,043
Perinatal	3,756	3,889	3,875	3,857	3,835
Other ⁶	215	218	213	207	208

¹ Total numbers include persons of unknown race/ethnicity. ² Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total. ³ Includes Asian/Pacific Islander legacy cases. ⁴ Hispanics/Latinos can be of any race. ⁵ Heterosexual contact with a person known to have, or to be at high risk for, HIV infection. ⁶ Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

Source: U.S. Centers for Disease Control and Prevention, Atlanta, GA, *HIV Surveillance Report, 2009*, Vol. 21, <<http://www.cdc.gov/hiv/surveillance/resources/reports/2009report>>.