

APPENDIX 4: Bacterial Pathogen Growth and Inactivation

This guidance represents the Food and Drug Administration's (FDA's) current thinking on this topic. It does not create or confer any rights for or on any person and does not operate to bind FDA or the public. You can use an alternative approach if the approach satisfies the requirements of the applicable statutes and regulations. If you want to discuss an alternative approach, contact the FDA staff responsible for implementing this guidance. If you cannot identify the appropriate FDA staff, call the telephone number listed on the title page of this guidance.

This appendix contains information on the growth and inactivation of bacterial pathogens.

Table A-1 contains information on the minimum water activity (a_w), acidity (pH), and temperature; the maximum, pH, water phase salt, and temperature; and oxygen requirements that will sustain growth for the bacterial pathogens that are of greatest concern in seafood processing. Data shown are the minimum or maximum values, the extreme limits reported among the references cited. These values may not apply to your processing conditions.

Table A-2 contains information on maximum, cumulative time and internal temperature combinations for exposure of fish and fishery products that, under ordinary circumstances, will be safe for the bacterial pathogens that are of greatest concern in seafood processing. These maximum, cumulative exposure times are derived from published scientific information.

Because the nature of bacterial growth is logarithmic, linear interpolation using the time and temperature guidance may not be appropriate. Furthermore, the food matrix effects bacterial growth (e.g., presence of competing microorganisms, available nutrients, growth restrictive agents). Consideration of such attributes is needed when using the information in Tables A-1 and A-2.

In summary, Table A-2 indicates that:

For raw, ready-to-eat products:

- If at any time the product is held at internal temperatures above 70°F (21.1°C), exposure time (i.e., time at internal temperatures

above 50°F (10°C) but below 135°F (57.2°C)) should be limited to 2 hours (3 hours if *Staphylococcus aureus* (*S. aureus*) is the only pathogen of concern),

OR

Alternatively, exposure time (i.e., time at internal temperatures above 50°F (10°C) but below 135°F (57.2°C)) should be limited to 4 hours, as long as no more than 2 of those hours are between 70°F (21.1°C) and 135°F (57.2°C);

OR

- If at any time the product is held at internal temperatures above 50°F (10°C) but never above 70°F (21.1°C), exposure time at internal temperatures above 50°F (10°C) should be limited to 5 hours (12 hours if *S. aureus* is the only pathogen of concern);

OR

- The product is held at internal temperatures below 50°F (10°C) throughout processing,

OR

Alternatively, the product is held at ambient air temperatures below 50°F (10°C) throughout processing.

For cooked, ready-to-eat products:

- If at any time the product is held at internal temperatures above 80°F (26.7°C), exposure time (i.e., time at internal temperatures above 50°F (10°C) but below 135°F (57.2°C)) should be limited to 1 hour (3 hours if *S. aureus* is the only pathogen of concern),

OR

Alternatively, if at any time the product is held at internal temperatures above 80°F (26.7°C), exposure time (i.e., time at internal temperatures above 50°F (10°C) but below 135°F (57.2°C)) should be limited to 4 hours, as long as no more than 1 of those hours is above 70°F (21.1°C);

OR

- If at any time the product is held at internal temperatures above 70°F (21.1°C) but never above 80°F (26.7°C), exposure time at internal temperatures above 50°F (10°C) should be limited to 2 hours (3 hours if *S. aureus* is the only pathogen of concern),

OR

Alternatively, if the product is never held at internal temperatures above 80°F (26.7°C), exposure times at internal temperatures above 50°F (10°C) should be limited to 4 hours, as long as no more than 2 of those hours are above 70°F (21.1°C);

OR

- If at any time the product is held at internal temperatures above 50°F (10°C) but never above 70°F (21.1°C), exposure time at internal temperatures above 50°F (10°C) should be limited to 5 hours (12 hours if *S. aureus* is the only pathogen of concern);

OR

- The product is held at internal temperatures below 50°F (10°C) throughout processing,

OR

Alternatively, the product is held at ambient air temperatures below 50°F (10°C) throughout processing.

Note that the preceding recommended critical limits do not address internal product temperatures between 40°F (4.4°C), the recommended maximum storage temperature for refrigerated fish and fishery products, and 50°F (10°C). That is because growth of foodborne pathogenic bacteria is very slow

at these temperatures and the time necessary for significant growth is longer than would be reasonably likely to occur in most fish and fishery product processing steps. However, if you have processing steps that occur at these temperatures that approach the maximum cumulative exposure times listed in Table A-2 below for the pathogenic bacteria of concern in your product, you should consider development of a critical limit for control at these temperatures.

It is not possible to furnish recommendations for each pathogenic bacteria, process, type of fish and fishery product, and temperature or combination of temperatures. Programmable models to predict growth rates for certain pathogens associated with various foods under differing conditions have been developed by the U.S. Department of Agriculture's (Pathogen Modeling Program (PMP)) and the United Kingdom's (Food MicroModel (FMM) program). These programs can provide growth curves for selected pathogens. You indicate the conditions, such as pH, temperature, and salt concentration that you are interested in and the models provide pathogen growth predictions (e.g., growth curve, time of doubling, time of lag phase, and generation time). FDA does not endorse or require the use of such modeling programs, but recognizes that the predictive growth information they provide may be of assistance to some processors. However, you are cautioned that significant deviations between actual microbiological data in specific products and the predictions do occur, including those for the lag phase of growth. Therefore, you should validate the time and temperature limits derived from such predictive models.

Table A-3 contains information on the destruction of *Listeria monocytogenes* (*L. monocytogenes*). Lethal rate, as used in this table, is the relative lethality of 1 minute at the designated internal product temperature as compared with the lethality of 1 minute at the reference internal product temperature of 158°F (70°C) (i.e., $z = 13.5^\circ\text{F}$ (7.5°C)). For example, 1

minute at 145°F (63°C) is 0.117 times as lethal as 1 minute at 158°F (70°C). The times provided are the length of time at the designated internal product temperature necessary to deliver a 6D process for *L. monocytogenes*. The length of time at a particular internal product temperature needed to accomplish a six logarithm reduction in the number of *L. monocytogenes* (6D) is, in part, dependent upon the food in which it is being heated. The values in the table are generally conservative and apply to all foods. You may be able to establish a shorter process time for your food by conducting scientific thermal death time studies. Additionally, lower degrees of destruction may be acceptable in your food if supported by a scientific study of the normal initial levels in the food. It is also possible that higher levels of destruction may be necessary in some foods, if especially high initial levels are anticipated.

Table A-4 contains information on the destruction of *Clostridium botulinum* (*C. botulinum*) type B (the most heat-resistant form of non-proteolytic *C. botulinum*). Lethal rate, as used in this table, is the relative lethality of 1 minute at the designated internal product temperature as compared with the lethality of 1 minute at the reference product internal temperature of 194°F (90°C) (i.e., for temperatures less than 194°F (90°C), $z = 12.6^\circ\text{F}$ (7.0°C); for temperatures above 194°F (90°C), $z = 18^\circ\text{F}$ (10°C)). The times provided are the length of time at the designated internal product temperature necessary to deliver a 6D process for *C. botulinum*. The values in the table are generally conservative. However, these values may not be sufficient for the destruction of non-proteolytic *C. botulinum* in dungeness crabmeat because of the potential protective effect of lysozyme. You may be able to establish a shorter process time for your food by conducting scientific thermal death time studies. Additionally, lower degrees of destruction may be acceptable in your food if supported by a scientific study of the normal inoculum in the food.

TABLE A-1
LIMITING CONDITIONS FOR PATHOGEN GROWTH

PATHOGEN	MIN. A _w (USING SALT)	MIN. pH	MAX. pH	MAX. % WATER PHASE SALT	MIN. TEMP.	MAX. TEMP.	OXYGEN REQUIREMENT
BACILLUS CEREUS	0.92	4.3	9.3	10	39.2°F 4°C	131°F ¹ 55°C	facultative anaerobe ⁴
CAMPYLOBACTER JEJUNI	0.987	4.9	9.5	1.7	86°F 30°C	113°F 45°C	micro-aerophile ²
CLOSTRIDIUM BOTULINUM, TYPE A, AND PROTEOLYTIC TYPES B AND F	0.935	4.6	9	10	50°F 10°C	118.4°F 48°C	anaerobe ³
CLOSTRIDIUM BOTULINUM, TYPE E, AND NON-PROTEOLYTIC TYPES B AND F	0.97	5	9	5	37.9°F 3.3°C	113°F 45°C	anaerobe ³
CLOSTRIDIUM PERRINGENS	0.93	5	9	7	50°F 10°C	125.6°F 52°C	anaerobe ³
PATHOGENIC STRAINS OF ESCHERICHIA COLI	0.95	4	10	6.5	43.7°F 6.5°C	120.9°F 49.4°C	facultative anaerobe ⁴
LISTERIA MONOCYTOGENES	0.92	4.4	9.4	10	31.3°F -0.4°C	113°F 45°C	facultative anaerobe ⁴
SALMONELLA SPP.	0.94	3.7	9.5	8	41.4°F 5.2°C	115.2°F 46.2°C	facultative anaerobe ⁴
SHIGELLA SPP.	0.96	4.8	9.3	5.2	43°F 6.1°C	116.8°F 47.1°C	facultative anaerobe ⁴
STAPHYLOCOCCUS AUREUS GROWTH	0.83	4	10	20	44.6°F 7°C	122°F 50°C	facultative anaerobe ⁴
STAPHYLOCOCCUS AUREUS TOXIN FORMATION	0.85	4	9.8	10	50°F 10°C	118°F 48°C	facultative anaerobe ⁴
VIBRIO CHOLERAE	0.97	5	10	6	50°F 10°C	109.4°F 43°C	facultative anaerobe ⁴
VIBRIO PARAHAEMOLYTICUS	0.94	4.8	11	10	41°F 5°C	113.5°F 45.3°C	facultative anaerobe ⁴
VIBRIO VULNIFICUS	0.96	5	10	5	46.4°F 8°C	109.4°F 43°C	facultative anaerobe ⁴
YERSINIA ENTEROCOLITICA	0.945	4.2	10	7	29.7°F -1.3°C	107.6°F 42°C	facultative anaerobe ⁴

1. Has significantly delayed growth (>24 hours) at 131°F (55°C).

2. Requires limited levels of oxygen.

3. Requires the absence of oxygen.

4. Grows either with or without oxygen.

TABLE A-2
TIME AND TEMPERATURE GUIDANCE FOR
CONTROLLING PATHOGEN GROWTH AND TOXIN FORMATION IN FISH AND FISHERY PRODUCTS

POTENTIALLY HAZARDOUS CONDITION	PRODUCT TEMPERATURE	MAXIMUM CUMULATIVE EXPOSURE TIME
GROWTH AND TOXIN FORMATION BY <i>BACILLUS CEREUS</i>	39.2-43°F (4-6°C) 44-59°F (7-15°C) 60-70°F (16-21°C) Above 70°F (21°C)	5 days 1 day 6 hours 3 hours
GROWTH OF <i>CAMPYLOBACTER JEJUNI</i>	86-93°F (30-34°C) Above 93°F (34°C)	48 hours 12 hours
GERMINATION, GROWTH, AND TOXIN FORMATION BY <i>CLOSTRIDIUM BOTULINUM</i> TYPE A, AND PROTEOLYTIC TYPES B AND F	50-70°F (10-21°C) Above 70°F (21°C)	11 hours 2 hours
GERMINATION, GROWTH, AND TOXIN FORMATION BY <i>CLOSTRIDIUM BOTULINUM</i> TYPE E, AND NON-PROTEOLYTIC TYPES B AND F	37.9-41°F (3.3-5°C) 42-50°F (6-10°C) 51-70°F (11-21°C) Above 70°F (21°C)	7 days 2 days 11 hours 6 hours
GROWTH OF <i>CLOSTRIDIUM PERFRINGENS</i>	50-54°F (10-12°C) 55-57°F (13-14 °C) 58-70°F (15-21°C) Above 70°F (21°C)	21 days 1 day 6 hours ¹ 2 hours
GROWTH OF PATHOGENIC STRAINS OF <i>ESCHERICHIA COLI</i>	43.7-50°F (6.6-10°C) 51-70°F (11-21°C) Above 70°F (21°C)	2 days 5 hours 2 hours
GROWTH OF <i>LISTERIA MONOCYTOGENES</i>	31.3-41°F (-0.4-5°C) 42-50°F (6-10°C) 51-70°F (11-21°C) 71-86°F (22-30°C) Above 86°F (30°C)	7 days 1 day 7 hours 3 hours 1 hour
GROWTH OF <i>SALMONELLA</i> SPECIES	41.4-50°F (5.2-10°C) 51-70°F (11-21°C) Above 70°F (21°C)	2 days 5 hours 2 hours
GROWTH OF <i>SHIGELLA</i> SPECIES	43-50°F (6.1-10°C) 51-70°F (11-21°C) Above 70°F (21°C)	2 days 5 hours 2 hours
GROWTH AND TOXIN FORMATION BY <i>STAPHYLOCOCCUS AUREUS</i>	50°F (7-10°C) 51-70°F (11-21°C) Above 70°F (21°C)	14 days 12 hours ¹ 3 hours
GROWTH OF <i>VIBRIO CHOLERA</i>	50°F (10°C) 51-70°F (11-21°C) 71-80°F (22-27°C) Above 80°F (27°C)	21 days 6 hours 2 hours 1 hour ²
GROWTH OF <i>VIBRIO PARAHAEMOLYTICUS</i>	41-50°F (5-10°C) 51-70°F (11-21°C) 71-80°F (22-27°C) Above 80°F (27°C)	21 days 6 hours 2 hours 1 hour ²
GROWTH OF <i>VIBRIO VULNIFICUS</i>	46.4-50°F (8-10°C) 51-70°F (11-21°C) 71-80°F (22-27°C) Above 80°F (27°C)	21 days 6 hours 2 hours 1 hour ²
GROWTH OF <i>YERSINIA ENTEROCOLITICA</i>	29.7-50°F (-1.3-10°C) 51-70°F (11-21°C) Above 70°F (21°C)	1 day 6 hours 2.5 hours

1. Additional data needed.
2. Applies to cooked, ready-to-eat foods only.

TABLE A-3
INACTIVATION OF LISTERIA MONOCYTOGENES

INTERNAL PRODUCT TEMPERATURE (°F)	INTERNAL PRODUCT TEMPERATURE (°C)	LETHAL RATE	TIME FOR 6D PROCESS (MINUTES)
145	63	0.117	17.0
147	64	0.158	12.7
149	65	0.215	9.3
151	66	0.293	6.8
153	67	0.398	5.0
154	68	0.541	3.7
156	69	0.736	2.7
158	70	1.000	2.0
160	71	1.359	1.5
162	72	1.848	1.0
163	73	2.512	0.8
165	74	3.415	0.6
167	75	4.642	0.4
169	76	6.310	0.3
171	77	8.577	0.2
172	78	11.659	0.2
174	79	15.849	0.1
176	80	21.544	0.09
178	81	29.286	0.07
180	82	39.810	0.05
182	83	54.116	0.03
183	84	73.564	0.03
185	85	100.000	0.02

Note: z = 13.5°F (7.5°C).

TABLE A-4
INACTIVATION OF NON-PROTEOLYTIC CLOSTRIDIUM BOTULINUM TYPE B

INTERNAL PRODUCT TEMPERATURE (°F)	INTERNAL PRODUCT TEMPERATURE (°C)	LETHAL RATE*	TIME FOR 6D PROCESS (MINUTES)
185	85	0.193	51.8
187	86	0.270	37.0
189	87	0.370	27.0
190	88	0.520	19.2
192	89	0.720	13.9
194	90	1.000	10.0
196	91	1.260	7.9
198	92	1.600	6.3
199	93	2.000	5.0
201	94	2.510	4.0
203	95	3.160	3.2
205	96	3.980	2.5
207	97	5.010	2.0
208	98	6.310	1.6
210	99	7.940	1.3
212	100	10.000	1.0

Note: For temperatures less than 194°F (90°C), z = 12.6°F (7.0°C); for temperatures above 194°F (90°C), z = 18°F (10°C).

*Note: These lethal rates and process times may not be sufficient for the destruction of non-proteolytic C. botulinum in dungeness crabmeat because of the potential that substances that may be naturally present, such as lysozyme, may enable the pathogen to more easily recover from heat damage.

BIBLIOGRAPHY.

We have placed the following references on display in the Division of Dockets Management, Food and Drug Administration, 5630 Fishers Lane, rm. 1061, Rockville, MD 20852. You may see them at that location between 9 a.m. and 4 p.m., Monday through Friday. As of March 29, 2011, FDA had verified the Web site address for the references it makes available as hyperlinks from the Internet copy of this guidance, but FDA is not responsible for any subsequent changes to Non-FDA Web site references after March 29, 2011.

- Abrahamsson, K., B. Gullmar, and N. Molin. 1966. The effect of temperature on toxin formation and toxin stability of *Clostridium botulinum* type E in different environments. *Can. J. Microbiol.* 12:385-394.
- Adams, M. R., C. L. Little, and M. C. Easter. 1991. Modeling the effect of pH, acidulant, and temperature on the growth of *Yersinia enterocolitica*. *J. Appl. Bacteriol.* 71:65-71.
- Adesiyun, A. A. 1984. Enterotoxigenicity of *Staphylococcus aureus* strains isolated from Nigerian ready-to-eat foods. *J. Food Protect.* 47:438-440.
- Ajmal, M. 1968. Growth and toxin production of *Clostridium botulinum* type E. *J. Appl. Bacteriol.* 31:120-123.
- Ando, Y., and H. Iida. 1970. Factors affecting the germination of spores of *Clostridium botulinum* type E. *Japan. J. Microbiol.* 14:361-370.
- Ando, Y. 1971. The germination requirements of spores of *Clostridium botulinum* type E. *Japan. J. Microbiol.* 15:515-525.
- Aryanta, R. W., G. H. Fleet, and K. A. Buckle. 1991. The occurrence and growth of microorganisms during the fermentation of fish sausage. *Int. J. Food Microbiol.* 13:143-156.
- Augustin, J. C., L. Rosso, and V. Carlier. June 15, 2000. A model describing the effect of temperature history on lag time for *Listeria monocytogenes*. *Int. J. Food Microbiol.* 57(3):169-181.
- Augustin, J. C., V. Zuliani, M. Cornu, and L. Guillier. 2005. Growth rate and growth probability of *Listeria monocytogenes* in dairy, meat and seafood products in suboptimal conditions. *J. Appl. Microbiol.* 99(5):1019-1042.
- Badhey, H., D. J. Cleri, R. F. D'Amato, J. R. Vernaleo, V. Veinni, J. Tessler, A. A. Wallman, A. J. Mastellone, M. Giuliani, and L. Hochstein. 1986. Two fatal cases of type E adult food-borne botulism with early symptoms and terminal neurologic signs. *J. Can. Microbiol.* 23:616-618.
- Baird-Parker, A. C. 1971. Factors affecting the production of bacterial food poisoning toxins. *J. Appl. Bacteriol.* 34:181-197.
- Baird-Parker, A. C., and B. Freame. 1967. Combined effect of water activity, pH and temperature on the growth of *Clostridium botulinum* from spore and vegetative cell inocula. *J. Appl. Bacteriol.* 30:420-429.
- Baker, D. A., C. Genigeorgis, and G. Garcia. 1990. Prevalence of *Clostridium botulinum* in seafood and significance of multiple incubation temperatures for determination of its presence and type in fresh retail fish. *J. Food Protect.* 53:668-673.
- Baker, D. A., C. Genigeorgis, J. Glover, and V. Razavilar. 1990. Growth and toxigenesis of *Clostridium botulinum* type E in fishes packaged under modified atmospheres. *Int. J. Food Microbiol.* 10:269-290.
- Baynes, N. C., J. Comrie, and J. H. Prain. 1983. Detection of bacterial growth by the Malthus conductance meter. *Med. Lab. Sci.* 40:149-158.
- Beckers, H. J., F. M. van Leusden, and P. D. Tips. 1985. Growth and enterotoxin production of *Staphylococcus aureus* in shrimp. *J. Hyg., Camb.* 95:685-693.
- Beltran, A., C. Pelaez, and A. Moral. 1989. Keeping quality of vacuum-packed smoked sardine fillets: microbiological aspects. *Z. Lebensm Unters Forsch.* 188:232-236.

- Benedict, R. C., T. Partridge, D. Wells, and R. L. Buchanan. 1993. *Bacillus cereus*: aerobic growth kinetics. *J. Food Prot.* 56(3): 211-214.
- Ben Embarek, P. K. 1994. Presence, detection, and growth of *Listeria monocytogenes* in seafoods: a review. *Int. J. Food Microbiol.* 23:17-34.
- Ben Embarek, P. K., and H. H. Huss. 1992. Growth of *Listeria monocytogenes* in lightly preserved fish products, p. 293-303. *In* H. H. Huss, et al. (ed.), *Quality Assurance in the Fish Industry, Proceedings of an International Conference, Copenhagen, Denmark, August 26-30, 1991.* Elsevier Sci. Publ. B.V., Amsterdam.
- Ben Embarek, P. K., and H. H. Huss. 1993. Heat resistance of *Listeria monocytogenes* in vacuum packaged pasteurized fish fillets. *Int. J. Food Microbiol.* 20:85-95.
- Bergdoll, M. S. 1989. *Staphylococcus aureus*, p. 463-523. *In* M. P. Doyle (ed.), *Foodborne microbial pathogens.* Marcel Dekker, Inc., New York, NY.
- Beuchat, L. R. 1973. Interacting effects of pH, temperature, and salt concentration on growth and survival of *Vibrio parahaemolyticus*. *Appl. Microbiol.* 25:844-846.
- Beuchat, L. R. 1974. Combined effects of water activity, solute, and temperature on the growth of *Vibrio parahaemolyticus*. *Appl. Microbiol.* 27:1075-1080.
- Beuchat, L. R., M. R. Clavero, and C. B. Jaquette. 1997. Effects of nisin and temperature on survival, growth, and enterotoxin production characteristics of psychrotrophic *Bacillus cereus* in beef gravy. *Appl. Environ. Microbiol.* 63(5):1953-1958.
- Boutin, B. K., J. G. Bradshaw, and W. H. Stroup. 1982. Heat processing of oysters naturally contaminated with *Vibrio cholerae* Serotype O1. *J. Food Prot.* 45:169-171.
- Boyd, J. W., and B. A. Southcott. 1971. Effects of sodium chloride on outgrowth and toxin production of *Clostridium botulinum* Type E in cod homogenates. *J. Fish. Res. Bd. Can.* 28:1071-1075.
- Brocklehurst, T. F., and B. M. Lund. 1990. The influence of pH, temperature and organic acids on the initiation of growth of *Yersinia enterocolitica*. *J. Appl. Bacteriol.* 69:390-397.
- Bryan, F. L. 1979. *Staphylococcus aureus.* *In* *Food microbiology: public health and spoilage aspects.* The Avi Publishing Co., Inc., Westport, CT.
- Buchanan, R. L. 1991. Microbiological criteria for cooked, ready-to-eat shrimp and crabmeat. *Food Technol.* 45:157-160.
- Buchanan, R. L. 1991. Using spreadsheet software for predictive microbiology applications. *J. Food Safety.* 11:123-134.
- Buchanan, R. L. 1993. Predictive food microbiology. *Trends Food Sci. Technol.* 4:6-11.
- Buchanan, R. L., and J. G. Phillips. 1990. Response surface model for predicting the effects of temperature, pH, sodium chloride content, sodium nitrite concentration and atmosphere on the growth of *Listeria monocytogenes*. *J. Food Prot.* 53:370-376, 381.
- Buchanan, R. L., and L. A. Klawitter. 1992. The effect of incubation temperature, initial pH, and sodium chloride on the growth kinetics of *Escherichia coli*. *Food Microbiol.* 9:185-196.
- Buchanan, R. L., and L. K. Bagi. 1997. Effect of water activity and humectant identity on the growth kinetics of *Escherichia coli* O157:H7. *Food Microbiol.* 14:413-423.
- Buchanan, R. L., and M. L. Cygnarowicz. 1990. A mathematical approach toward defining and calculating the duration of the lag phase. *Food Microbiol.* 7:237-240.
- Campanini, M., A. Casolari, and S. Gola. 1977. Bacterial growth and limiting pH. *Industria Conserve.* 52:326-331.
- Carlin, F., C. Nguyen-the, and A. Abreu da Silva. 1995. Factors affecting the growth of *Listeria monocytogenes* on minimally processed fresh endive. *J. Appl. Bacteriol.* 78:636-646.

- Carlson, V. L., and G. H. Snoeyenbos. 1972. Relationship of population kinetics of *Salmonella typhimurium* and cultural methodology. *Am. J. Vet. Res.* 33:177-184.
- Casales, M. R., C. E. Del Valle, and C. L. Soule. 1985. Critical heating point for thermal processing of mussels in cans. *J. Food Sci.* 50:836-837.
- Catsaras, M., and D. Grebot. 1984. Multiplication des *Salmonella* dans la viande hachée. *Bull. Acad. Vet. France.* 57:501-512.
- Chitchester, C. O., and H. D. Graham (ed.). 1973. *Microbial safety of fishery products.* Academic Press, New York, NY.
- Christiansen, L. N., J. Deffner, E. M. Foster, and H. Sugiyama. 1968. Survival and outgrowth of *Clostridium botulinum* type E spores in smoked fish. *Appl. Microbiol.* 16:553-557.
- Cole, M. B., M. V. Jones, and C. Holyoak. 1990. The effect of pH, salt concentration and temperature on the survival and growth of *Listeria monocytogenes*. *J. Appl. Bacteriol.* 69:63-72.
- Colwell, R. R. (ed.), 1984. *Vibrios in the environment.* John Wiley Interscience, New York, NY.
- Connor, D. E., V. N. Scott, D. T. Bernard, and D. A. Kautter. 1989. Potential *Clostridium botulinum* hazards associated with extended shelf-life refrigerated foods; a review. *J. Food Safety.* 10:131-153.
- Cook, D. W. 1994. Effect of time and temperature on multiplication of *Vibrio vulnificus* in postharvest Gulf Coast shellstock oysters. *Appl. Environ. Microbiol.* 60:3483-3484.
- Cook, D. W., and A. D. Ruple. 1989. Indicator bacteria and *Vibrionaceae* multiplication in post-harvest shellstock oysters. *J. Food Protect.* 52:343-349.
- Cook, D. W., and A. D. Ruple. 1992. Cold storage and mild heat treatment as processing aids to reduce the number of *Vibrio vulnificus* in raw oysters. *J. Food Protect.* 55:985-989.
- Cortesi, M. L., T. Sarli, A. Santoro, N. Murru, and T. Pepe. 1997. Distribution and behavior of *Listeria monocytogenes* in three lots of naturally-contaminated vacuum-packed smoked salmon stored at 2 and 10°C. *Int. J. Food Microbiol.* 37:209-214.
- Craven, S. E. 1980. Growth and sporulation of *Clostridium perfringens* in foods. *Food Technol.* 34:80-87.
- Curtis, L. M., M. Patrick, and C. de W. Blackburn. 1995. Survival of *Campylobacter jejuni* in foods and comparison with a predictive model. *Lett. Appl. Microbiol.* 21:194-197.
- Dahl Sawyer, C. A., and J. J. Pestka. 1985. Foodservice systems: presence of injured bacteria in foods during food product flow. *Ann. Rev. Microbiol.* 39:51-67.
- Dalgaard, P., and L. V. Jorgensen. 1998. Predicted and observed growth of *Listeria monocytogenes* in seafood challenge tests and in naturally contaminated cold-smoked salmon. *Int. J. Food Microbiol.* 40:105-115.
- Datz, M., C. Janetzki-Mittmann, S. Franke, F. Gunzer, H. Schmidt, and H. Karch. 1996. Analysis of the enterohemorrhagic *Escherichia coli* O157 DNA region containing lambdoid phage gene p and shiga-like toxin structural genes. *Appl. Environ. Microbiol.* 62:791-797.
- Davies, A. R., and A. Slade. 1995. Fate of *Aeromonas* and *Yersinia* on modified-atmosphere packaged (MAP) cod and trout. *Lett. Appl. Microbiol.* 21:354-358.
- Deibel, K. E. 1995. Potential of *Staphylococcus aureus* to produce enterotoxin in fish batter at various temperatures, p. 33. *In* Medallion Lab (ed.), *Proceedings of the IFT Annual Meeting.* Medallion Lab, Minneapolis, MN.
- Dengremont, E., and J. M. Membre. 1995. Statistical approach for comparison of the growth rates of five strains of *Staphylococcus aureus*. *Appl. Environ. Microbiol.* 61:4389-4395.

- DePaola, A., G. M. Capers, and D. Alexander. 1994. Densities of *Vibrio vulnificus* in the intestines of fish from the U.S. Gulf Coast. *Appl. Environ. Microbiol.* 60:984-988.
- Destro, M. T., M. F. F. Leitao, and J. M. Farber. 1996. Use of molecular typing methods to trace the dissemination of *Listeria monocytogenes* in a shrimp processing plant. *Appl. Environ. Microbiol.* 62:705-711.
- Dickerson, R. W., and M. R. Berry, Jr. 1976. Heating curves during commercial cooking of the blue crab. *J. Milk Food Technol.* 39:258-262.
- Dickerson, R. W., Jr., and R. B. Read, Jr. 1973. Cooling rates of foods. *J. Milk Food Technol.* 36:167-171.
- Dickson, J. S., G. R. Siragusa, and J. E. Wray, Jr. 1992. Predicting the growth of *Salmonella typhimurium* on beef by using the temperature function integration technique. *Appl. Environ. Microbiol.* 58:3482-3487.
- Diez de Medina, D., G. Flick, R. Whittman, A. Diallo, and R. Croonenberghs. 1995. Growth of *Listeria monocytogenes* in fresh blue crab (*Callinectes sapidus*) meat in the presence of naturally occurring microflora during refrigerated storage, p. 178. *In* Institute of Food Technologists (ed.), Abstracts of IFT Annual Meeting 1995. Virginia Polytechnic Institute, Blacksburg, VA.
- Dillon, R., and T. Patel. 1992. *Listeria* in seafoods: a review. *J. Food Prot.* 55:1009-1015.
- Dillon, R., and T. Patel. 1993. Effect of cold smoking and storage temperature on *Listeria monocytogenes* in inoculated cod fillets (*Gadus morbus*). *Food Res. Int.* 26:97-101.
- Dillon, R., T. Patel, and S. Ratnam. 1994. Occurrence of *Listeria* in hot and cold smoked seafood products. *Int. J. Food Microbiol.* 22:73-77.
- Doyle, M. P. (ed.), 1989. Foodborne bacterial pathogens. Marcel-Dekker, New York, NY.
- Doyle, M. P., and D. J. Roman. 1982. Prevalence and survival of *Campylobacter jejuni* in unpasteurized milk. *Appl. Environ. Microbiol.* 44(5):1154-1158.
- Doyle, M. P., and D. J. Roman. 1981. Growth and survival of *Campylobacter fetus* subsp. *jejuni* as a function of temperature and pH. *J. Food Prot.* 44:596-601.
- Duran, A. P., B. A. Wentz, J. M. Lanier, F. D. McClure, A. H. Schwab, A. Swartzentruber, R. J. Barnard, and R. B. Read. 1983. Microbiological quality of breaded shrimp during processing. *J. Food Prot.* 46:974-977.
- Eklund, M. W., D. I. Weiler, and F. T. Poysky. 1967. Outgrowth and toxin production of non-proteolytic type B *Clostridium botulinum* at 3.3 to 5.6°C. *J. Bacteriol.* 93:1461-1462.
- Eklund, M. W., F. T. Poysky, R. N. Paranjpye, L. C. Lashbrook, M. E. Peterson, and G. A. Pelroy. 1995. Incidence and sources of *Listeria monocytogenes* in cold-smoked fishery products and processing plants. *J. Food Prot.* 58:502-508.
- Eklund, M. W., G. A. Pelroy, R. Paranjpye, M. E. Peterson, and F. M. Teeny. 1982. Inhibition of *Clostridium botulinum* types A and E toxin production by liquid smoke and NaCl in hot-process smoke-flavored fish. *J. Food Prot.* 45:935-941.
- Elsea, W. R., W. E. Mosher, R. G. Lennon, V. Markellis, and P. F. Hoffman. 1971. An epidemic of food-associated pharyngitis and diarrhea. *Arch. Environ. Health.* 23:48-56.
- El-Shenawy, M. A., and E. H. Marth. 1988. Inhibition and inactivation of *Listeria monocytogenes* by sorbic acid. *J. Food Prot.* 51:842-847.
- El-Shenawy, M. A., and E. H. Marth. 1988. Sodium benzoate inhibits growth of or inactivates *Listeria monocytogenes*. *J. Food Prot.* 51:525-530.
- European Chilled Food Federation. 1997. Guidelines for good hygienic practice in the manufacture of chilled foods.

- Fantasia, L. D., L. Mestrandrea, J. P. Schrade, and J. Yager. 1975. Detection and growth of enteropathogenic *Escherichia coli* in soft ripened cheese. *Appl. Microbiol.* 29(2):179-185.
- Fapohunda, A. O., K. W. McMillen, D. L. Marshall, and W. M. Waites. 1994. Growth of selected cross-contaminating bacterial pathogens on beef and fish at 15° and 35°C. *J. Food Prot.* 57:337-340.
- Farber, J. M. 1991. *Listeria monocytogenes* in fish products. *J. Food Prot.* 54:922-924.
- Farber, J. M. and J. Z. Losos. 1988. *Listeria monocytogenes*: a foodborne pathogen. *Can. Med. Assoc. J.* 138(5):413-418.
- Fernandes, C. F., G. J. Flick, and T. B. Thomas. 1998. Growth of inoculated psychrotrophic pathogens on refrigerated fillets of aquacultured rainbow trout and channel catfish. *J. Food Protect.* 61:313-317.
- Fernandez, P. S., S. M. George, C. C. Stills, and M. W. Peck. 1997. Predictive model of the effect of CO₂, pH, temperature, and NaCl on the growth of *Listeria monocytogenes*. *Int. J. Food Microbiol.* 37:37-45.
- Fluer, F. S., and Y. V. Ezepchuk. 1970. *Microbiologiya* 39:396-410 (Source: Rhodehammel, E. J., and S. M. Harmon. Chapter 14. *Bacillus cereus*. In *FDA bacteriological analytical manual*, 8th ed. (1995). AOAC International, McLean, VA).
- Food and Agriculture Organization of the United Nations. 1999. *FAO Fisheries Report. No. 604, Report of the FAO expert consultation on the trade impact of Listeria in fish products*. Rome, Italy.
- Food and Agriculture Organization of the United Nations and World Health Organization. 2005. *Microbiological Risk Assessment Series No. 8, Risk assessment of Vibrio vulnificus in raw oysters: interpretative summary and technical report*. Rome, Italy.
- Fuchs, R. S., and P. J. A. Reilly. 1992. The incidence and significance of *Listeria monocytogenes* in seafoods, p. 217-229. In H. H. Huss, M. Jakobsen, and J. Liston (ed.), *Quality assurance in the fish industry*. Proc. Int. Conf. August 26-30, 1991, Copenhagen. Elsevier Science Publishers B.V., Amsterdam.
- Fuhs, A., and G. J. Bonde. 1957. The nutritional requirements of *Clostridium perfringens*. *J. Gen. Microbiol.* 16:317-329.
- Fujino, T., G. Sakaguchi, and Y. Takeda (ed.). 1973. *International Symposium on Vibrio parahaemolyticus*. Saikon, Tokyo.
- Garren, D. M., M. A. Harrison, and Y-W. Huang. 1994. *Clostridium botulinum* type E outgrowth and toxin production in vacuum skin packaged shrimp. *Food Microbiol.* 11:467-472.
- Garren, D. M., M. A. Harrison, and Y-W. Huang. 1995. Growth and production of toxin of *Clostridium botulinum* type E in rainbow trout under various storage conditions. *J. Food Prot.* 58:863-866.
- Garthright, W. E. 1991. Refinements in the prediction of microbial growth curves. *Food Microbiol.* 8:239-248.
- Gay, M., O. Cerf, and K. R. Davey. 1996. Significance of pre-incubation temperature and inoculum concentration on subsequent growth of *Listeria monocytogenes* at 14 degrees C. *J. Appl. Bacteriol.* 81(4):433-438.
- George, S. M., B. M. Lund, and T. F. Brocklehurst. 1988. The effect of pH and temperature on initiation of growth of *Listeria monocytogenes*. *Lett. Appl. Microbiol.* 6:153-156.
- George, S. M., L. C. C. Richardson, and M. W. Peck. 1996. Predictive models of the effect of temperature, pH, and acetic and lactic acids on the growth of *Listeria monocytogenes*. *Int. J. Food Microbiol.* 31:73-90.
- Gibbs, P. A., A. R. Davies, and R. S. Fletcher. 1994. Incidence and growth of psychrotrophic *Clostridium botulinum* in food. *Food Control.* 5:5-7.
- Gibson, A. M., N. Bratchell, and T. A. Roberts. 1988. Predicting microbial growth:

- growth responses of *Salmonellae* in a laboratory medium as affected by pH, sodium chloride and storage temperature. *Int. J. Food Microbiol.* 6:155-178.
- Gill, C. O., and L. M. Harris. 1982. Survival and growth of *Campylobacter fetus* subsp. *jejuni* on meat and in cooked foods. *Appl. Environ. Microbiol.* 44:259-263.
 - Godwin, G. J., R. M. Grodner, and A. F. Novak. 1977. Twenty-four hour methods for bacteriological analyses in frozen raw breaded shrimp. *J. Food Science.* 42:750-754.
 - Gooch, J. A., A. DePaola, J. Bowers, and D. L. Marshall. June 2002. Growth and survival of *Vibrio parahaemolyticus* in postharvest American oysters. *J. Food Prot.* 65(6):970-974.
 - Gould, G. W. 1999. Sous vide foods: conclusions of an ECFF botulinum working party. *Food Control.* 10: 47-51.
 - Gourama, H., W. Y. Tsai, and L. B. Bullerman. 1991. Growth and production of enterotoxins A and D by *Staphylococcus aureus* in salad bar ingredients and clam chowder. *J. Food Prot.* 54:844-847.
 - Goverde, R. L. J., J. G. Kusters, and J. H. J. Huls in't Veld. 1994. Growth rate and physiology of *Yersinia enterocolitica*; influence of temperature and presence of the virulence plasmid. *J. Appl. Bacteriol.* 77:96-104.
 - Grecz, N., and L. H. Arvay. 1982. Effect of temperature on spore germination and vegetative cell growth of *Clostridium botulinum*. *Appl. Environ. Microbiol.* 43:331-337.
 - Greenwood, M. H., E. F. C. Coetzee, B. M. Ford, P. Gill, W. L. Hooper, S. C. W. Matthews, S. Patrick, J. V. S. Pether, and R. J. D. Scott. 1985. The bacteriological quality of selected retail, ready-to-eat food products. III. Cooked crustaceans and mollusks. *Environ. Health.* 93:236-239.
 - Griffin, M. R., E. Dalley, M. Fitzpatrick, and S. H. Austin. 1983. *Campylobacter* gastroenteritis associated with raw clams. *J. Med. Soc. N.J.* 80:607-609.
 - Groubert, T. N., and J. D. Oliver. 1994. Interaction of *Vibrio vulnificus* and the Eastern oyster, *Crassostrea virginica*. *J. Food Prot.* 57:224-228.
 - Guyer, S., and T. Jemmi. 1991. Behavior of *Listeria monocytogenes* during fabrication and storage of experimentally contaminated smoked salmon. *Appl. Environ. Microbiol.* 57:1523-1527.
 - Halpin-Dohnalek, M. I., and E. H. Marth. 1989. *Staphylococcus aureus*: production of extracellular compounds and behavior in foods - a review. *J. Food Prot.* 52:267-282.
 - Hanninen, M. L., H. Korkeala, and P. Pakkala. 1984. Growth and survival characteristics of *Campylobacter jejuni* in liquid egg. *J. Hyg., Camb.* 92:53-58.
 - Hany, O. E., R. Siddiqi, and M. A. Khan. 1993. Growth response of *Listeria monocytogenes* NCTC 7973 in two different media at four incubation temperatures. *Ann. Acad. Med. Singapore, China.* 22:300-302.
 - Harrison, M. A., Y-W. Huang, C-H. Chao, and T. Shineman. 1991. Fate of *Listeria monocytogenes* on packaged, refrigerated, and frozen seafood. *J. Food Prot.* 54:524-527.
 - Hathcox, A. K., L. R. Beuchat, and M. P. Doyle. 1995. Death of enterohemorrhagic *Escherichia coli* O157:H7 in real mayonnaise and reduced-calorie mayonnaise dressing as influenced by initial population and storage temperature. *Appl. Environ. Microbiol.* 61:4172-4177.
 - Hauschild, A. H. W. 1989. *Clostridium botulinum*, p. 111-189. In M. P. Doyle (ed.), *Foodborne bacterial pathogens*. Marcel Dekker, Inc., New York, NY.
 - Helmy, Z. A., A. Abd-El-Bakey, and E. I. Mohamed. 1984. Factors affecting germination and growth of *Bacillus cereus* spores in milk. *Zbl. Mikrobiol.* 139:135-141.
 - Hobbs, G. 1976. *Clostridium botulinum* and its importance in fishery products. *Adv. Food Res.* 22:135-185.

- Holler, C., D. Witthuhn, and B. Janzen-Blunck. 1998. Effect of low temperatures on growth, structure, and metabolism of *Campylobacter coli* SP10. *Appl. Environ. Microbiol.* 64: 581-587.
- Holley, R. A., and M. Proulx. 1986. Use of egg washwater pH to prevent survival of *Salmonella* at moderate temperatures. *Poult. Sci.* 65:922-928.
- Hudson, J. A., and S. J. Mott. 1993. Growth of *Listeria monocytogenes*, *Aeromonas hydrophila*, and *Yersinia enterocolitica* on cold-smoked salmon under refrigeration and mild temperature abuse. *Food Microbiol.* 10:61-68.
- Hudson, J. A., and S. M. Avery. 1994. Growth of *Listeria monocytogenes*, *Aeromonas hydrophila*, and *Yersinia enterocolitica* on cooked mussel tissue under refrigeration and mild temperature abuse. *J. Food Saf.* 14:41-52.
- Hughes, A., and A. Hurst. 1980. The effect of NaCl on the upper temperature limit for growth of and enterotoxin synthesis by *Staphylococcus aureus*. *Can. J. Microbiol.* 26:507-510.
- Huss, H. H. 1992. Development and use of the HACCP concept in fish processing. *Int. J. Food Microbiol.* 15:33-44.
- Huss, H. H. 1997. Control of indigenous pathogenic bacteria in seafood. *Food Control* 8:91-98.
- Huss H. H., I. Shaeffer, E. R. Petersen, and D. C. Cann. February 1979. Toxin production by *Clostridium botulinum* type E in fresh herring in relation to the measured oxidation potential (Eh). *Nord. Vet. Med.* 1(2):81-86.
- Hwang, C. A., and M. L. Tamplin. July 25, 2005. The influence of mayonnaise pH and storage temperature on the growth of *Listeria monocytogenes* in seafood salad. *Int. J. Food Microbiol.* 102(3):277-285.
- Ingham, S. C., R. A. Alford, and A. P. McCown. 1990. Comparative growth rates of *Salmonella typhimurium* and *Pseudomonas fragi* on cooked crab meat stored under air and modified atmosphere. *J. Food Prot.* 53:566-567.
- International Commission on Microbiological Specifications for Foods. 1996. *Microorganisms in foods 5. Characteristics of microbial pathogens.* ICMSF, International Union of Biological Societies. Blackie Academic & Professional, London, England.
- Islam, M. S., M. K. Hasan, and S. I. Khan. 1993. Growth and survival of *Shigella flexneri* in common Bangladeshi foods under various conditions of time and temperature. *Appl. Environ. Microbiol.* 59:652-654.
- James, D. G., and J. Olley. 1971. Spoilage of shark. *Aust. Fish.* 30:11-13.
- Jemmi, T., and A. Keusch. 1992. Behavior of *Listeria monocytogenes* during processing and storage of experimentally contaminated hot-smoked trout. *Int. J. Food Microbiol.* 15:339-346.
- Johnson, K. M., C. L. Nelson, and F. F. Busta. 1983. Influence of temperature on germination and growth of spores of emetic and diarrheal strains of *Bacillus cereus* in a broth medium and in rice. *J. Food Sci.* 48: 286-287. Jones, J. E., S. J. Walker, J. P. Sutherland, M. W. Peck, and C. L. Little. 1994. Mathematical modeling of the growth, survival and death of *Yersinia enterocolitica*. *Int. J. Food Microbiol.* 23:433-447.
- Jorgensen, L. V., and H. H. Huss. 1998. Prevalence and growth of *Listeria monocytogenes* in naturally contaminated seafood. *Int. J. Food Microbiol.* 42:127-131.
- Juneja, V. K., and B. S. Marmer. 1996. Growth of *Clostridium perfringens* from spore inocula in *sous-vide* products. *Int. J. Food Microbiol.* 32:115-123.
- Juneja, V. K., B. S. Marmer, J. G. Phillips, and S. A. Palumbo. 1996. Interactive effects of temperature, initial pH, sodium chloride, and sodium pyrophosphate on the growth kinetics of *Clostridium perfringens*. *J. Food Prot.* 59: 963-968.

- Juneja, V. K., J. E. Call, B. S. Marmer, and A. J. Miller. 1994. The effect of temperature abuse on *Clostridium perfringens* in cooked turkey stored under air and vacuum. *Food Microbiol.* 11:187-193.
- Jyhshium, L., S. L. In, J. L. Slonczewski, and J. W. Foster. 1995. Comparative analysis of extreme acid survival in *Salmonella typhimurium*, *Shigella flexneri*, and *Escherichia coli*. *J. Bacteriol.* 177:4097-4104.
- Kaneko, T., and R. R. Colwell. 1973. Ecology of *Vibrio parahaemolyticus* in Chesapeake Bay. *J. Bacteriol.* 113:24-32.
- Kang, C. K., M. Woodburn, A. Pagenkopf, and R. Cheney. 1969. Growth, sporulation, and germination of *Clostridium perfringens* in media of controlled water activity. *Appl. Microbiol.* 18:798-805.
- Karim, P., and B. Embarek. 1994. Presence, detection, and growth of *Listeria monocytogenes* in sea-foods: a review. *Int. J. Food Microbiol.* 23:17-34.
- Kaspar, C. W., and M. L. Tamplin. 1993. Effects of temperature and salinity on the survival of *Vibrio vulnificus* in seawater and shellfish. *Appl. Environ. Microbiol.* 59:2425-2429.
- Kauppi, K. L., S. R. Tatini, F. Harrell, and P. Feng. 1996. Influence of substrate and low temperature on growth and survival of verotoxigenic *Escherichia coli*. *Food Microbiol.* 13:397-405.
- Kelana, L. C., and M. W. Griffiths. 2003. Growth of autoluminescent *Campylobacter jejuni* in response to various environmental conditions. *J. Food Prot.* 66:1190-1197.
- Koff, R. S., and H. S. Sear. 1967. Internal temperature of steamed clams. *New Engl. J. Med.* 276:737-739.
- Kramer, J. M., and J. M. Gilbert. Chapter 2, *Bacillus cereus* and other *Bacillus* species. In *Foodborne bacterial pathogens*, M. P. Doyle (ed.), Marcel Dekker, Inc., New York, NY.
- Krieg, N. R. (ed.), 1984. *Bergey's manual of systematic bacteriology*, vol. 1. Williams & Wilkins, Baltimore, MD.
- Lachia, R. V., G. J. Silverman, and R. Sharp. 1988. Guide to the salvage of temperature-abused food products in military commissaries. Technical Report on Project No. AAF87-12,II. U.S. Army Natick Research, Development and Engineering Center, Natick, MA.
- Lappi, V. R., A. Ho, K. Gall, and M. Wiedmann. 2004. Prevalence and growth of *Listeria* on naturally contaminated smoked salmon over 28 days of storage at 4 degrees C. *J. Food Prot.* 67(5):1022-1026.
- Leung, C-K., Y-W. Huang, and O. C. Pancorbo. 1992. Bacteria pathogens and indicators in catfish and pond environments. *J. Food Prot.* 55:424-427.
- Lindberg, C. W., and E. Borch. 1994. Predicting the aerobic growth of *Yersinia enterocolitica* O:3 at different pH-values, temperatures and L-lactate concentrations using conductance measurements. *Int. J. Food Microbiol.* 22:141-153.
- Little, C. L., and S. Knochel. 1994. Growth and survival of *Yersinia enterocolitica*, *Salmonella*, and *Bacillus cereus* in brie stored at 4, 8, and 20°C. *Int. J. Food Microbiol.* 24:137-145.
- Lotter, L. P., and L. Leistner. 1978. Minimal water activity for enterotoxin A production and growth of *Staphylococcus aureus*. *Appl. Environ. Microbiol.* 36:377-380.
- Lu, J. Y., R. D. Pace, and W. D. Plahar. 1991. Storage conditions and microbial quality of smoked dry herring in Ghana. *J. Food Prot.* 54:557-559.
- Lund, B. M., A. F. Graham, S. M. George, and D. Brown. 1990. The combined effect of incubation temperature, pH, and sorbic acid on the probability of growth of non-proteolytic, type B *Clostridium botulinum*. *J. Appl. Bacteriol.* 69:481-492.

- Mackey, B. M., and N. Bratchell. 1989. A review: the heat resistance of *Listeria monocytogenes*. Lett. Appl. Microbiol. 9:89-94.
- Mackey, B. M., T. A. Roberts, J. Mansfield, and G. Farkas. 1980. Growth of *Salmonella* on chilled meat. J. Hyg., Camb. 85:115-124.
- Matches, J. R. 1982. Microbial changes in packages, p. 46-70. In R. Martin (ed.), Proceedings of the First National Conference on Seafood Packaging and Shipping. National Fisheries Institute, Seattle, WA.
- Matches, J. R., and J. Liston. 1968. Low temperature growth of *Salmonella*. J. Food Sci. 33:641-645.
- Matches, J. R., and J. Liston. 1972. Effect of pH on low temperature growth of *Salmonella*. J. Milk Food Technol. 35:49-52.
- Matches, J. R., J. Liston, and L. P. Daneault. 1971. Survival of *Vibrio parahaemolyticus* in fish homogenate during storage at low temperature. Appl. Microbiol. 21:951-952.
- Maurelli, A. T., B. Blackmon, and R. Curtiss, III. 1984. Temperature-dependent expression of virulence genes in *Shigella* species. Infect. Immun. 43:195-201.
- McClure, P. J., A. L. Beaumont, J. P. Sutherland, and T. A. Roberts. March 3, 1997. Predictive modeling of growth of *Listeria monocytogenes*. The effects on growth of NaCl, pH, storage temperature and NaNO₂. Int. J. Food Microbiol. 34(3):221-232.
- McElroy, D. M., L.-A. Jaykus, and P. M. Foegeding. February 2000. Validation and analysis of modeled predictions of growth of *Bacillus cereus* spores in boiled rice. J. Food Prot. 63(2):268-272.
- Microbial Food Safety Research Unit. December 12, 2005. Pathogen Modeling Program 7.0 version 1.1.1433.15425. U.S. Department of Agriculture, Agriculture Research Service. <http://www.ars.usda.gov/Services/docs.htm?docid=6788>.
- Miles, D. W., T. Ross, J. Olley, and T. A. McMeekin. 1997. Development and evaluation of a predictive model for the effect of temperature and water activity on the growth rate of *Vibrio parahaemolyticus*. Int. J. Food Microbiol. 38:133-142.
- Molin, G., and I-M. Stenstroem. 1984. Effect of temperature on the microbial flora of herring fillets stored in air of carbon dioxide. J. Appl. Bacteriol. 56:275-282.
- Molin, G., I-M. Stenstroem, and A. Ternstroem. 1983. The microbial flora of herring fillets after storage in carbon dioxide, nitrogen, or air at 2 degrees C. J. Appl. Bacteriol. 55:49-56.
- Motes, M. L., Jr. 1991. Incidence of *Listeria* spp. in shrimp, oysters, and estuarine waters. J. Food Prot. 54:170-173.
- Murphree, R. L., and M. L. Tamplin. 1995. Uptake and retention of *Vibrio cholerae* O1 in the eastern oyster, *Crassostrea virginica*. Appl. Environ. Microbiol. 61:3656-3660.
- Murphy, S. K., and J. D. Oliver. 1992. Effects of temperature abuse on survival of *Vibrio vulnificus* in oysters. Appl. Environ. Microbiol. 58(9):2771-2775.
- National Advisory Committee on Microbiological Criteria for Foods. 1991. *Listeria monocytogenes*: Recommendations of the National Advisory Committee on Microbiological Criteria for Foods. Int. J. Food Microbiol. 14:185-246.
- National Advisory Committee on Microbiological Criteria for Foods. 1992. Microbiological criteria for raw molluscan shellfish. J. Food Prot. 55:463-480.
- National Advisory Committee on Microbiological Criteria for Foods. 1992. Vacuum or modified atmosphere packaging for refrigerated raw fishery products. U.S. Department of Agriculture, Food Safety and Inspection Service, Executive Secretariat, Washington, DC.
- Olmez, H. K., and N. Aran. 2005. Modeling the growth kinetics of *Bacillus cereus* as a function of temperature, pH, sodium lactate

- and sodium chloride concentrations. *Int. J. Food Microbiol.* 98(2):135-143.
- Ostovar, K., and M. J. Bremier. 1975. Effect of thawing on growth of *Staphylococcus aureus* in frozen convenience food items. *J. Milk Food Technol.* 38:337-339.
 - Palumbo, S. A., J. E. Call, F. J. Schultz, and A. C. Williams. 1995. Minimum and maximum temperatures for growth and enterotoxin production by hemorrhagic strains of *Escherichia coli*. *J. Food Protect.* 58:352-356.
 - Pedrosa-Menabrito, A., and J. M. Regenstein. 1988. Shelf-life extension of fresh fish - a review. Part I - Spoilage of fish. *J. Food Qual.* 11:117-127.
 - Pedrosa-Menabrito, A., and J. M. Regenstein. 1990. Shelf-life extension of fresh fish - a review. Part II - Preservation of fish. *J. Food Qual.* 13:129-146.
 - Pelroy, G., M. Peterson, R. Paranjpye, J. Almond, and M. Eklund. 1994. Inhibition of *Listeria monocytogenes* in cold-process (smoked) salmon by sodium nitrite and packaging method. *J. Food Prot.* 57:114-119.
 - Pelroy, G. A., M. W. Eklund, R. N. Paranjpye, E. M. Suzuki, and M. E. Peterson. 1982. Inhibition of *Clostridium botulinum* types A and E toxin formation by sodium nitrite and sodium chloride in hot-process (smoked) salmon. *J. Food Prot.* 45:833-841.
 - Peters, A. C., L. Thomas, and J. W. T. Wimpenny. 1991. Effects of salt concentration on bacterial growth on plates with gradients of pH and temperature. *FEMS Microbiol. Lett.* 77:309-314.
 - Peterson, M. E., G. A. Pelroy, F. T. Poysky, R. N. Paranjpye, R. M. Dong, G. M. Pigott, and M. W. Eklund. 1997. Heat-pasteurization process for inactivation of nonproteolytic types of *Clostridium botulinum* in pickled dungeness crabmeat. *J. Food Prot.* 60:928-934.
 - Peterson, M. E., G. A. Pelroy, R. N. Paranjpye, F. T. Poysky, J. S. Almond, and M. W. Eklund. 1993. Parameters for control of *Listeria monocytogenes* in smoked fishery products: sodium chloride and packaging method. *J. Food Prot.* 56:938-943.
 - Potter, L., and L. Leistner. 1978. Minimal water activity for enterotoxin A production and growth of *Staphylococcus aureus*. *Appl. Environ. Microbiol.* 36:377-380.
 - Pradhan, L., P. Kanekar, and S. H. Godbole. 1985. Microbiology of spoiled mango pickles. Tolerance to salt, acidity, and oil of the microbes isolated from spoiled mango pickles. *J. Food Sci. Technol. India* 22:339-341.
 - Presser, K. A., T. Ross, and D. A. Ratkowsky. 1998. Modeling the growth limits (growth/no growth interface) of *Escherichia coli* as a function of temperature, pH, lactic acid concentration, and water activity. *Appl. Environ. Microbiol.* 64: 1773-1779.
 - Prost, E., and H. Riemann. 1967. Food-borne salmonellosis. *Ann. Rev. Microbiol.* 21:495-528.
 - Raj, H. D. 1970. Public health bacteriology of processed frozen foods. *Lab. Pract.* 19:374-377, 394.
 - Ratkowsky, D. A., J. Olley, T. A. McMeekin, and A. Ball. 1982. Relationship between temperature and growth rate of bacterial cultures. *J. Bacteriol.* 149:1-5.
 - Ratkowsky, D. A., R. K. Lowry, T. A. McMeekin, A. N. Stokes, and R. E. Chandler. 1983. Model for bacterial culture growth rate throughout the entire biokinetic temperature range. *J. Bacteriol.* 154:1222-1226.
 - Reddy, N. R., A. Paradis, M. G. Roman, H. M. Solomon, and E. J. Rhodehamel. (1996). Toxin development by *Clostridium botulinum* in modified atmosphere-packaged fresh tilapia fillets during storage. *J. Food Sci.* 61:632-635.
 - Reed, G. H. 1993. Foodborne illness (Part 2): Salmonellosis. *Dairy, Food, Environ. San.* 13:706.
 - Reed, G. H. 1994. Foodborne illness (Part 3): *Clostridium perfringens* gastroenteritis. *Dairy, Food, Environ. San.* 14:16-17.

- Reed, G. H. 1994. Foodborne illness (Part 4): *Bacillus cereus* gastroenteritis. Dairy, Food, Environ. San. 14:87.
- Reed, G. H. 1994. Foodborne illness (Part 8): *Escherichia coli* Dairy, Food, Environ. San. 14:329-330.
- Reed, G. H. 1994. Foodborne illness (Part 11): Yersinosis. Dairy, Food, Environ. San. 14:536.
- Rey, C. R., H. W. Walker, and P. L. Rohrbaugh. 1975. The influence of temperature on growth, sporulation, and heat resistance of spores of six strains of *Clostridium perfringens*. J. Milk Food Technol. 38:461-465.
- Richards, J. C. S., A. C. Jason, G. Hobbs, D. M. Gibson, and R. H. Christie. 1978. Electronic measurement of bacterial growth. J. Phys. E.: Sci. Instrum. 11:560-568.
- Roberts, D., and R. J. Gilbert. 1979. Survival and growth of non-cholera vibrios in various foods. J. Hyg., Camb. 82:123-131.
- Roberts, T. A., and A. M. Gibson. 1983. Chemical methods for controlling *Clostridium botulinum* in processed meats. Food Technol. 40:163-171, 176.
- Roberts, T. A., and C. M. Derrick. 1978. The effect of curing salts on the growth of *Clostridium perfringens (welchii)* in a laboratory medium. J. Food Technol. 13:394-353.
- Roberts, T. A., and G. Hobbs. 1968. Low temperature growth characteristics of clostridia. J. Appl. Bacteriol. 31:75-88.
- Romick, T. L., H. P. Fleming, and R. F. McFeeters. 1996. Aerobic and anaerobic metabolism of *Listeria monocytogenes* in defined glucose medium. Appl. Environ. Microbiol. 62:304-307.
- Rørvik, L. M., and M. Yndestad. 1991. *Listeria monocytogenes* in foods in Norway. Int. J. Food Microbiol. 13:97-104.
- Rørvik, L. M., M. Yndestad, and E. Skjerve. 1991. Growth of *Listeria monocytogenes* in vacuum-packed, smoked salmon during storage at 4°C. Int. J. Food Microbiol. 14:111-118.
- Rusul, G., and N. H. Yaacob. 1995. Prevalence of *Bacillus cereus* in selected foods and detection of enterotoxin using TECRA-VIA and BCET-RPLA. Int. J. Food Microbiol. 25:131-139.
- Schiemann, D. A. 1988. Examination of enterotoxin production at low temperature by *Yersinia enterocolitica* in culture media and foods. J. Food Prot. 51:571-573.
- Segner, W. P., C. F. Schmidt, and J. K. Boltz. 1971. Minimal growth temperature, sodium chloride tolerance, pH sensitivity, and toxin production of marine and terrestrial strains of *Clostridium botulinum* type C. Appl. Microbiol. 22:1025-1029.
- Shaw, M. K., A. G. Marr, and J. L. Ingraham. 1971. Determination of the minimal temperature for growth of *Escherichia coli*. J. Bacteriol. 105:683-684.
- Skinner, G. E., and J. W. Larkin. 1998. Conservative prediction of time to *Clostridium botulinum* toxin formation for use with time-temperature indicators to ensure the safety of foods. J. Food Protect. 61:1154-1160.
- Smith, G. R., and A. Turner. 1989. The production of *Clostridium botulinum* toxin in mammalian, avian, and piscine carrion. Epidemiol. Infect. 102:467-471.
- Smith, M. G. 1985. The generation time, lag time, and minimum temperature of growth of coliform organisms on meat, and the implications for codes of practice in abattoirs. J. Hyg., Camb. 94:289-300.
- Sneath, P. H. A. (ed.), 1986. Bergey's manual of systematic bacteriology, vol. 2. Williams & Wilkins, Baltimore, MD.
- Sokari, T. 1991. Distribution of enterotoxigenic *Staphylococcus aureus* in ready-to-eat foods in eastern Nigeria. Int. J. Food Microbiol. 12:275-279.
- Stern, N. J., and A. W. Kotula. 1982. Survival of *Campylobacter jejuni* inoculated into ground beef. Appl. Environ. Microbiol. 44(5):1150-1153.

- Subcommittee on Microbiological Criteria. 1985. An evaluation of the role of microbiological criteria for foods and food ingredients. Committee on Food Protection, National Research Council. National Academy Press, Washington, DC. 256 p.
- Sutherland, A. D. 1993. Toxin production by *Bacillus cereus* in dairy products. J. Dairy Res. 60:569-574.
- Sutherland, J. P., and A. J. Bayliss. 1994. Predictive modeling of growth of *Yersinia enterocolitica*: the effects of temperature, pH, and sodium chloride. Int. J. Food Microbiol. 21:197-215.
- Sutherland, J. P., A. Aherne, and A. L. Beaumont. 1996. Preparation and validation of a growth model for *Bacillus cereus*: the effects of temperature, pH, sodium chloride and carbon dioxide. Int. J. Food Microbiol. 30:359-372.
- Sutherland, J. P., A. J. Bayliss, and D. S. Braxton. 1995. Predictive modeling of growth of *Escherichia coli* O157:H7: the effects of temperature, pH and sodium chloride. Int. J. Food Microbiol. 25:29-49.
- Sutherland, J. P., A. J. Bayliss, and T. A. Roberts. 1994. Predictive modeling of growth of *Staphylococcus aureus*: the effects of temperature, pH and sodium chloride. Int. J. Food Microbiol. 21:217-236.
- Swartzentruber, A., A. H. Schwab, A. P. Duran, B. A. Wents, and R. B. Read, Jr. 1980. Microbiological quality of frozen shrimp and lobster tail in the retail market. Appl. Environ. Microbiol. 40:765-769.
- Syndicat National des Fabricants de Plats Prepares. 1995. Aide a la maitrise de l'hygiene alimentaire. Collection Alesial, Alesial Services, Paris, France.
- Taormina, P. J., G. W. Bartholomew, and W. J. Dorsa. 2003. Incidence of *Clostridium perfringens* in commercially produced cured raw meat product mixtures and behavior in cooked products during chilling and refrigerated storage. J Food Prot. 66(1):72-81.
- Tatani, S. R. 1973. Influence of food environments on growth of *Staphylococcus aureus* and production of various enterotoxins. J. Milk Food Technol. 36:559.
- Telzak, E. E., E. P. Bell, D. A. Kautter, L. Crowell, L. D. Budnick, D. L. Morse, and S. Schultz. 1990. International outbreak of type E botulism due to uneviscerated fish. J. Infect. Dis. 161:340-342.
- Thayer, D. W., W. S. Muller, R. L. Buchanan, and J. G. Phillips. 1987. Effect of NaCl, pH, temperature, and atmosphere on growth of *Salmonella typhimurium* in glucose-mineral salts medium. Appl. Environ. Microbiol. 53:1311-1315.
- Thomas, L. V., J. W. T. Wimpenny, and A. C. Peters. 1991. An investigation of the effects of four variables in the growth of *Salmonella typhimurium* using two types of gradient gel plates. Int. J. Food Microbiol. 14:261-275.
- Thomas, L. V., J. W. T. Wimpenny, and A. C. Peters. 1992. Testing multiple variables on the growth of a mixed inoculum of *Salmonella* strains using gradient plates. Int. J. Food Microbiol. 15:165-175.
- Tilton, R. C., and R. W. Ryan. 1987. Clinical and ecological characteristics of *Vibrio vulnificus* in the northeastern United States. Diagn. Microbiol. Infect. Dis. 6:109-117.
- Tipparaju, S., S. Ravishankar, and P. J. Slade. February 2004. Survival of *Listeria monocytogenes* in vanilla-flavored soy and dairy products stored at 8 degrees C. J. Food Prot. 67(2):378-382.
- Twedt, R. M., P. L. Spaulding, and H. E. Hall. 1969. *Vibrio parahaemolyticus*. J. Bacteriol. 98:511-518.
- U.S. Food and Drug Administration and Food Safety and Inspection Service. 1992. Preventing foodborne listeriosis. Department of Health and Human Services, Food and Drug Administration, and U.S. Department of Agriculture, Food Safety and Inspection Service, Washington, DC.

- U.S. Food and Drug Administration and U.S. Department of Agriculture. 2003. Quantitative assessment of relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods. <http://www.fda.gov/downloads/food/scienceresearch/researchareas/riskassessmentsafetyassessment/ucm197330.pdf>.
- U.S. Food and Drug Administration. Foodborne pathogenic microorganisms and natural toxins handbook. In The bad bug book. <http://www.fda.gov/Food/FoodSafety/FoodborneIllness/FoodborneIllnessFoodbornePathogensNaturalToxins/default.htm>.
- U.S. Food and Drug Administration. 2007. National Shellfish Sanitation Program guide for the control of molluscan shellfish 2007 Revision. Department of Health and Human Services, Public Health Service, Food and Drug Administration, Center for Food Safety and Applied Nutrition, College Park, MD. <http://www.fda.gov/Food/FoodSafety/Product-SpecificInformation/Seafood/FederalStatePrograms/NationalShellfishSanitationProgram/ucm046353.htm>.
- U.S. Food and Drug Administration. 2005. Quantitative risk assessment on the public health impact of pathogenic *Vibrio parahaemolyticus* in raw oysters. Department of Health and Human Services, Public Health Service, Food and Drug Administration, Center for Food Safety and Applied Nutrition, College Park, MD.
- Venkataramaiah, N., and A. G. Kempton. 1975. Bacterial growth in seafood on restaurant premises. *Can. J. Microbiol.* 21(11):1788-1797.
- Walker, S. J., P. Archer, and J. G. Banks. 1990. Growth of *Listeria monocytogenes* at refrigeration temperatures. *J. Appl. Bacteriol.* 68:157-162.
- Wang, C., and L. A. Shelef. 1992. Behavior of *Listeria monocytogenes* and the spoilage microflora in fresh cod fish treated with lysozyme and EDTA. *Food Microbiol.* 9:207-213.
- Ward, D. R. and C. R. Hackney (ed.). 1991. Microbiology of marine food products. Van Nostrand Reinhold, New York, NY.
- Watkins, W., and S. McCarthy (ed.). 1995. Proceedings of the 1994 *Vibrio vulnificus* Workshop, second printing. Department of Health and Human Services, Public Health Service, Food and Drug Administration, Center for Food Safety and Applied Nutrition, College Park, MD.
- Weagant, S. D., P. N. Sado, K. G. Colburn, J. D. Torkelson, F. A. Stanley, M. H. Krane, S. E. Shields, and C. F. Thayer. 1988. The incidence of *Listeria* species in frozen seafood products. *J. Food Prot.* 51:655-657.
- Weber, J. T., R. G. Hibbs, Jr., A. Darwish, B. Mishu, A. L. Corwin, M. Rakha, C. L. Hatheway, S. el Sharkawy, S. A. el-Rahim, and M. F. al-Hamd, et al. 1993. A massive outbreak of type E botulism associated with traditional salted fish in Cairo. *J. Infect. Dis.* 167:451-454.
- Weichart, D., and J. D. Oliver. 1992. Low temperature induced non-culturability and killing of *Vibrio vulnificus*. *FEMS Microbiol. Lett.* 79:205-210.
- West, P. A. 1989. The human pathogenic vibrios - a public health update with environmental perspectives. *Epidemiol. Infect.* 103:1-34.
- Whiting, R. C., and K. A. Naftulin. 1992. Effect of headspace oxygen concentration on growth and toxin production by proteolytic strains of *Clostridium botulinum*. *J. Food Prot.* 55:23-17.
- Wright, A. C., R. T. Hill, J. A. Johnson, M. C. Roghman, R. R. Colwell, and J. G. Morris, Jr. 1996. Distribution of *Vibrio vulnificus* in the Chesapeake Bay. *Appl. Environ. Microbiol.* 62:717-724.

- Yang, S. E., and C. C. Chou. July 2000. Growth and survival of *Escherichia coli* O157:H7 and *Listeria monocytogenes* in egg products held at different temperatures. *J. Food Prot.* 63(7):907-911.
- Zaika, L. L., A. H. Kim, and L. Ford, 1991. Effect of sodium nitrite on growth of *Shigella flexneri*. *J. Food Prot.* 54:424-428.
- Zaika, L. L., E. Moulden, L. Weimer, J. G. Phillips, and R. L. Buchanan. 1994. Model for the combined effects of temperature, initial pH, sodium chloride and sodium nitrite concentrations on anaerobic growth of *Shigella flexneri*. *Int. J. Food Microbiol.* 23:345-358.
- Zaika, L. L., J. G. Phillips, and R. L. Buchanan. 1992. Model for aerobic growth of *Shigella flexneri* under various conditions of temperature, pH, sodium chloride and sodium nitrite concentrations. *J. Food Prot.* 55:509-513.
- Zaika, L. L., J. G. Phillips, J. S. Fanelli, and O. J. Scullen. 1998. Revised model for aerobic growth of *Shigella flexneri* to extend the validity of predictions at temperatures between 10 and 19°C. *Int. J. Food Microbiol.* 41:9-19.
- Zwietering, M. H., I. Jongenburger, F. M. Rombouts, and K. van't Riet. 1990. Modeling of the bacterial growth curve. *Appl. Environ. Microbiol.* 56:1875-1881.

NOTES: