


# HIV Surveillance in Adolescents and Young Adults

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention  
Division of HIV/AIDS Prevention



# Diagnoses of HIV Infection among Adolescents and Young Adults Aged 13–24 Years, by Race/Ethnicity, 2007–2010—46 States and 5 U.S. Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.



# Diagnoses of HIV Infection among Adolescents and Young Adults Aged 13–24 Years, by Transmission Category, 2007–2010—46 States and 5 U.S. Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# Diagnoses of HIV Infection and Population among Adolescents Aged 13–19 Years, by Race/Ethnicity, 2010—46 States


**Diagnoses of HIV Infection**

N=2,234



**U.S. Population aged 13–19 years**

N=28,264,651



 American Indian/Alaska Native

 Asian

 Black/African American

 Multiple races

 Hispanic/Latino<sup>a</sup>

 Native Hawaiian/Other Pacific Islander

 White


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.




# Diagnoses of HIV Infection and Population among Young Adults Aged 20–24 Years, by Race/Ethnicity, 2010—46 States

**Diagnoses of HIV Infection**



**U.S. Population aged 20–24 years**



■ American Indian/Alaska Native  
■ Asian  
■ Black/African American

■ Hispanic/Latino<sup>a</sup>  
■ Native Hawaiian/Other Pacific Islander  
■ White

■ Multiple races


*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.




# Diagnoses of HIV Infection among Persons Aged 13 Years and Older, by Sex and Age Group, 2010—46 States and 5 U.S Dependent Areas


**13–19 years**  
N=2,266



**20–24 years**  
N=7,675



**≥ 25 years**  
N=38,137



■ Males ■ Females

*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



## Diagnoses of HIV Infection among Adolescent and Young Adult Males, by Age Group and Transmission Category, 2010—46 States and 5 U.S. Dependent Areas

Transmission category	13–19 years		20–24 years	
	No.	%	No.	%
Male-to-male sexual contact	1,634	91.2	5,898	89.8
Injection drug use (IDU)	23	1.3	143	2.2
Male-to-male sexual contact and IDU	51	2.9	235	3.6
Heterosexual contact <sup>a</sup>	84	4.7	289	4.4
Other <sup>b</sup>	0	0.0	0	0.0
<b>Total</b>	<b>1,792</b>	<b>100</b>	<b>6,565</b>	<b>100</b>

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



## Diagnoses of HIV Infection among Adolescent and Young Adult Females, by Age Group and Transmission Category, 2010—46 States and 5 U.S. Dependent Areas

Transmission category	13–19 years		20–24 years	
	No.	%	No.	%
Injection drug use	38	8.0	107	9.7
Heterosexual contact <sup>a</sup>	436	92.0	1,001	90.2
Other <sup>b</sup>	0	0.0	1	0.1
<b>Total</b>	<b>474</b>	<b>100</b>	<b>1,110</b>	<b>100</b>

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


<sup>b</sup> Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.





# Rates of Diagnoses of HIV Infection among Adolescents Aged 13–19 Years, 2010—46 States and 5 U.S. Dependent Areas


N=2,266



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

# Rates of Diagnoses of HIV Infection among Young Adults Aged 20–24 Years, 2010—46 States and 5 U.S. Dependent Areas



N=7,675



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



# Adolescents and Young Adults Aged 13–24 Years Living with a Diagnosis of HIV Infection, by Sex and Race/Ethnicity, Year-end 2009—46 States and 5 U.S. Dependent Areas



■ American Indian/Alaska Native  
■ Asian<sup>a</sup>  
■ Black/African American  
■ Multiple races

■ Hispanic/Latino<sup>b</sup>  
■ Native Hawaiian/Other Pacific Islander  
■ White  
■ Unknown races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


\* Includes 6 persons of unknown race/ethnicity.

<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.



# Adolescents and Young Adults Aged 13–24 Years Living with a Diagnosis of HIV Infection, by Sex and Transmission Category, Year–end 2009—46 States and 5 U.S. Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting.


<sup>a</sup> Heterosexual contact with a person known to have, or at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# Rates of Adolescents Aged 13–19 Years Living with a Diagnosis of HIV Infection, Year–end 2009— 46 States and 5 U.S. Dependent Areas

N=8,404




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



# Rates of Young Adults Aged 20–24 Years Living with a Diagnosis of HIV Infection, Year–end 2009— 46 States and 5 U.S. Dependent Areas


N=26,329



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.




# AIDS Diagnoses among Adolescents Aged 13–19 Years, by Sex, 1985–2010—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



# AIDS Diagnoses among Young Adults Aged 20–24 Years, by Sex, 1985–2010—United States and 6 U.S. Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.





# AIDS Diagnoses among Persons Aged 13 Years and Older, by Race/Ethnicity and Age Group, 2010—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.



# Rates of AIDS Diagnoses among Adolescents Aged 13–19 Years, 2010—United States and 6 U.S. Dependent Areas


N=582



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



# Rates of AIDS Diagnoses among Young Adults Aged 20–24 Years, 2010—United States and 6 U.S. Dependent Areas N=2,280




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



# Rates of Adolescents Aged 13–19 Years Living with an AIDS Diagnosis, Year-end 2009—United States and 6 U.S. Dependent Areas

N=3,282




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



# Rates of Young Adults Aged 20–24 Years Living with an AIDS Diagnosis, Year-end 2009—United States and 6 U.S. Dependent Areas

N=7,258



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

