

search

 Contacts: [Karen McNulty Walsh](#), (631) 344-8350 or [Mona S. Rowe](#), (631) 344-5056

 [Print-friendly](#)
 [E-mail](#)
[Article](#)
[News Archives](#)[BNL](#)**News Resources**[Newsroom Home](#)[News Archives](#)[Photo Archive](#)[Streaming Video](#)[Fact Sheets](#)[Science Magazine](#)[Management Bios](#)[About Brookhaven](#)**:: Addiction News**[High Dopamine Transporter Levels Not Correlated with ADHD](#)[Study Offers New Clues to Brain-Stomach Interaction in Overeating](#)[Developing Radiotracers for Imaging Studies in Addiction](#)[Study Offers Clues to Brain's Protective Mechanisms Against Alcoholism](#)[Study Reveals Biochemical Signature of Cocaine Craving in Humans](#)

Altered Perception of Reward in Human Cocaine Addiction

Brain-mapping, psychological studies support mechanism for compromised control in drug addiction

October 15, 2006

ATLANTA, GA -- People addicted to cocaine have an impaired ability to perceive rewards and exercise control due to disruptions in the brain's reward and control circuits, according to a series of brain-mapping studies and neuropsychological tests conducted at the U.S. Department of Energy's Brookhaven National Laboratory.

"Our findings provide the first evidence that the brain's threshold for responding to monetary rewards is modified in drug-addicted people, and is directly linked to changes in the responsiveness of the prefrontal cortex, a part of the brain essential for monitoring and controlling behavior," said [Rita Goldstein](#), a psychologist at Brookhaven Lab. "These results also attest to the benefit of using sophisticated brain-imaging tools combined with sensitive behavioral, cognitive, and emotional probes to optimize the study of drug addiction, a psychopathology that these tools have helped to identify as a disorder of the brain."

Goldstein will present details of these studies at a press conference on neuroscience and addiction at the Society for Neuroscience (SfN) annual meeting in Atlanta, Georgia, on Sunday, October 15, 2006, 2 to 3 p.m., and at a SfN symposium on Wednesday, October 18, 8:30 a.m.*

Goldstein's experiments were designed to test a theoretical model, called the Impaired Response Inhibition and Salience Attribution (I-RISA) model, which postulates that drug-addicted individuals disproportionately attribute salience, or value, to their drug of choice at the expense of other potentially but no-longer-rewarding stimuli — with a concomitant decrease in the ability to inhibit maladaptive drug use. In the experiments, the scientists subjected cocaine-addicted and non-drug-addicted individuals to a range of tests of behavior, cognition/thought, and emotion, while simultaneously monitoring their brain activity using functional magnetic resonance imaging (fMRI) and/or recordings of event-related potentials (ERP).



[Rita Goldstein](#) (click image to download hi-res version)


**Brookhaven Lab
 Addiction Symposium**

Feb. 16, 2007, San Francisco
 [Get BNL News via RSS](#)

In one study, subjects were given a monetary reward for their performance on an attention task. Subjects were given one of three amounts (no money, one cent, or 45 cents) for each correct response, up to a total reward of \$50 for their performance. The researchers also asked the subjects how much they valued different amounts of monetary reward, ranging from \$10 to \$1000.

More than half of the cocaine abusers rated \$10 as equally valuable as \$1000, “demonstrating a reduced subjective sensitivity to relative monetary reward,” Goldstein said.

“Such a ‘flattened’ sensitivity to gradients in reward may play a role in the inability of drug-addicted individuals to use internal cues and feedback from the environment to inhibit inappropriate behavior, and may also predispose these individuals to disadvantageous decisions — for example, trading a car for a couple of cocaine hits. Without a relative context, drug use and its intense effects — craving, anticipation, and high — could become all the more overpowering,” she said.



In control subjects (left), brain regions that play a part in experiencing reward were activated in a graded fashion in response to increasing monetary rewards. These regions were not activated in cocaine-addicted subjects offered the same rewards (right). This indicates that cocaine-addicted subjects' ability to respond to non-drug rewards is compromised.

The behavioral data collected during fMRI further suggested that, in the cocaine abusers, there was a “disconnect” between subjective measures of motivation (how much they said they were engaged in the task) and the objective measures of motivation (how fast and accurately they performed on the task).

“These behavioral data implicate a disruption in the ability to perceive inner motivational drives in cocaine addiction,” Goldstein said.

The fMRI results also revealed that non-addicted subjects responded to the different monetary amounts in a graded fashion: the higher the potential reward, the greater the response in the prefrontal cortex. In cocaine-addicted subjects, however, this region did not demonstrate a graded pattern of response to the monetary reward offered. Furthermore, within the cocaine-addicted group, the higher the sensitivity to money in the prefrontal cortex, the higher was the motivation and the self-reported ability to control behavior.

The ERP results showed a similarly graded brain response to monetary reward in healthy control subjects, but not in cocaine-addicted individuals.

“The dysfunctional interplay between reward processing and control of behavior observed in these studies could help to explain the chronically relapsing nature of drug addiction,” Goldstein said. “Our results also suggest the need for new clinical interventions aimed at helping drug abusers manage these symptoms as part of an effective treatment strategy.”

This work was funded by the Office of Biological and Environmental Research within the U.S. Department of Energy’s Office of Science; the National Institute on Drug Abuse; the National Alliance for Research on Schizophrenia and Depression (NARSAD) Young Investigator Award; the National Institute on Alcohol Abuse and Alcoholism; and the General Clinical Research Center at University Hospital Stony Brook. DOE has a long-standing interest in brain-imaging studies on addiction. Brain-imaging techniques such as MRI are a direct outgrowth of DOE’s support of basic physics and chemistry research.

Coauthors on the study included: Nelly Alia-Klein, Dardo Tomasi, Lisa A. Cottone, Thomas Maloney, Frank Telang, and Elisabeth C. Caparelli of Brookhaven Lab; Lei Zhang, Dimitris Samaras, and Nancy K. Squires of Stony Brook University; Linda Chang and Thomas Ernst of the University of Hawaii; and Nora D. Volkow of the National Institute on Drug Abuse.

*For more information about the Society for Neuroscience annual meeting, go to: <http://www.sfn.org/am2006/?CFID=833504&CFTOKEN=58003625>

Number: 06-110 | [BNL Media & Communications Office](#)



One of ten national laboratories overseen and primarily funded by the Office of Science of the U. S. Department of Energy (DOE), Brookhaven National Laboratory conducts research in the physical, biomedical, and environmental sciences, as well as in energy technologies and national security. Brookhaven Lab also builds and operates major scientific facilities available to university, industry and government researchers. Brookhaven is operated and managed for DOE’s Office of Science by Brookhaven Science Associates, a limited-liability company founded by the Research Foundation of the State University of New York on behalf of Stony Brook University, the largest academic user of Laboratory facilities, and Battelle, a nonprofit, applied science and technology organization.

[Privacy and Security Notice](#) | [Contact Web Services for help](#)