

Table 35. Tuberculosis Cases and Percentages by Initial Drug Regimen: Reporting Areas, 2010

Reporting Area	Total Cases	Cases in Persons Alive at Diagnosis	Cases with Information on Initial Drug Regimen		Percentage of Cases in Persons with Initial Drug Regimen ^{1,2}		
			No.	(%)	IR	IRZ	IRZE ³
United States	11,182	10,919	10,769	98.6	0.9	2.9	(85.0)
Alabama	146	142	142	(100.0)	(0.0)	(3.5)	(84.5)
Alaska	57	55	53	(96.4)	(0.0)	(9.4)	(81.1)
Arizona	283	280	266	(95.0)	(0.4)	(1.9)	(89.5)
Arkansas	78	78	78	(100.0)	(0.0)	(3.8)	(14.1)
California	2,327	2,267	2,240	(98.8)	(0.3)	(1.7)	(88.4)
Colorado	71	71	71	(100.0)	(0.0)	(0.0)	(80.3)
Connecticut	85	84	84	(100.0)	(0.0)	(3.6)	(85.7)
Delaware	20	20	20	(100.0)	(0.0)	(0.0)	(75.0)
District of Columbia	44	42	41	(97.6)	(0.0)	(0.0)	(100.0)
Florida	835	814	802	(98.5)	(0.5)	(4.5)	(85.3)
Georgia	411	399	392	(98.2)	(2.0)	(0.0)	(90.8)
Hawaii	115	114	112	(98.2)	(0.0)	(4.5)	(91.1)
Idaho	15	14	14	(100.0)	(0.0)	(21.4)	(78.6)
Illinois	372	364	355	(97.5)	(1.7)	(1.1)	(87.0)
Indiana	90	89	89	(100.0)	(1.1)	(6.7)	(88.8)
Iowa	48	47	47	(100.0)	(0.0)	(6.4)	(83.0)
Kansas	46	45	44	(97.8)	(0.0)	(2.3)	(93.2)
Kentucky	90	89	88	(98.9)	(0.0)	(1.1)	(88.6)
Louisiana	200	198	198	(100.0)	(1.0)	(5.6)	(87.9)
Maine	8	8	8	(100.0)	(0.0)	(0.0)	(62.5)
Maryland	220	216	216	(100.0)	(1.4)	(2.8)	(85.6)
Massachusetts	222	219	206	(94.1)	(1.0)	(1.9)	(73.3)
Michigan	184	173	167	(96.5)	(0.6)	(7.2)	(71.3)
Minnesota	135	133	133	(100.0)	(0.0)	(4.5)	(90.2)
Mississippi	116	115	114	(99.1)	(2.6)	(5.3)	(76.3)
Missouri	107	104	102	(98.1)	(0.0)	(6.9)	(76.5)
Montana	6	6	6	(100.0)	(0.0)	(0.0)	(100.0)
Nebraska	27	27	27	(100.0)	(0.0)	(0.0)	(85.2)
Nevada	114	114	114	(100.0)	(0.0)	(0.0)	(93.9)
New Hampshire	10	9	9	(100.0)	(0.0)	(0.0)	(88.9)
New Jersey	405	398	396	(99.5)	(1.3)	(2.8)	(87.1)
New Mexico	51	46	46	(100.0)	(2.2)	(0.0)	(93.5)
New York State ³	243	239	239	(100.0)	(1.3)	(1.7)	(83.7)
New York City	711	697	678	(97.3)	(0.3)	(2.9)	(73.5)
North Carolina	296	292	292	(100.0)	(0.0)	(2.1)	(83.9)
North Dakota	12	12	12	(100.0)	(0.0)	(0.0)	(100.0)
Ohio	190	183	182	(99.5)	(1.1)	(2.2)	(91.2)
Oklahoma	86	85	78	(91.8)	(10.3)	(15.4)	(57.7)
Oregon	87	83	83	(100.0)	(1.2)	(7.2)	(90.4)
Pennsylvania	238	227	226	(99.6)	(0.4)	(0.9)	(57.5)
Rhode Island	26	25	25	(100.0)	(0.0)	(0.0)	(96.0)
South Carolina	153	148	147	(99.3)	(0.0)	(6.8)	(88.4)
South Dakota	15	14	14	(100.0)	(0.0)	(14.3)	(85.7)
Tennessee	193	185	184	(99.5)	(0.0)	(4.9)	(88.6)
Texas	1,385	1,350	1,337	(99.0)	(2.1)	(3.2)	(89.0)
Utah	20	20	20	(100.0)	(0.0)	(10.0)	(90.0)
Vermont	5	5	5	(100.0)	(0.0)	(20.0)	(80.0)
Virginia	268	262	259	(98.9)	(0.0)	(0.4)	(97.7)
Washington	239	236	234	(99.2)	(0.4)	(2.6)	(86.3)
West Virginia	15	14	14	(100.0)	(0.0)	(0.0)	(100.0)
Wisconsin	55	55	53	(96.4)	(3.8)	(7.5)	(83.0)
Wyoming	7	7	7	(100.0)	(0.0)	(0.0)	(85.7)
American Samoa ⁴	3	3	3	(100.0)	(0.0)	(0.0)	(100.0)
Fed. States of Micronesia ⁴	171	171	171	(100.0)	(0.0)	(1.2)	(92.4)
Guam ⁴	100	97	97	(100.0)	(1.0)	(7.2)	(83.5)
Marshall Islands ⁴	196	193	192	(99.5)	(0.0)	(0.5)	(67.2)
N. Mariana Islands ⁴	32	30	30	(100.0)	(0.0)	(0.0)	(100.0)
Puerto Rico ⁴	80	78	78	(100.0)	(0.0)	(2.6)	(94.9)
Republic of Palau ⁴	17	17	17	(100.0)	(0.0)	(0.0)	(100.0)
U.S. Virgin Islands ⁴

¹Overall U.S. percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥75% of cases.

²I=isoniazid; R=rifampin; Z=pyrazinamide; E=ethambutol.

³Excludes New York City.

⁴Not included in U.S. totals.

Note: Excluding cases with no information on drug regimen, 109 (1.00%) persons were not started on any drugs, 15 (0.14%) were started on one drug, and 1,192 (10.96%) had an initial multidrug regimen other than IR, IRZ, or IRZE.