



Why We Screen Before Vaccination

December 2011

www.vaccines.mil





Across

- 1 Persons with this disease history should not receive smallpox (7)
- 5 Component added to vaccine to enhance immune response (8)
- 8 Common injection site reaction (8)
- 10 TST placed soon after this live vaccine may cause a false-negative (7)
- 13 Flu vaccine contraindicated in immunosuppressed persons (7)
- 14 Vaccine recommended by ACIP for individuals 60 years and older (6)
- 17 Children on this long term therapy should not receive FluMist (7)
- 18 Additional dose of immunizing agent weeks or years after initial dose (7)

Down

- 2 Recommends immunizations requirements, grp (4)
- 3 Prevents infections of common preventable diseases (13)
- 4 Suboptimal antibody response after vaccination may occur in persons undergoing this therapy (12)
- 6 Indicated for individuals traveling to rural areas in Asia during mosquito season (3)
- 7 Vaccine Adverse Event Reporting System; abbr. (5)
- 9 Influenza vaccine recommended for immunocompromised persons; abbr (3)
- 11 This type of medication to treat infection is not a contraindication for vaccination (11)

- 21** Indicated for children to prevent viral paralytic disease (5)
- 23** Recommended vaccine for smokers 18 yrs and older (12)
- 24** Do not vaccinate persons severely allergic to this YF vaccine component (3)
- 25** Vaccine stabilizer in MMR , an excipient (7)
- 27** Vaccine type contraindicated for pregnant women (4)
- 29** Vaccine contraindicated in breastfeeding mothers (8)
- 32** Vaccine component that may cause allergic reaction, i.e. MSG (10)
- 33** Recommended age for first dose of Hep B (5)
- 35** Vaccine recommended for female adolescents, abbr. (3)
- 36** Brief but severe symptoms; synm. (5)
- 38** Encephalopathy after this vaccine contraindicates a second dose (9)
- 12** Thrombocytopenia is a precaution for administration of this vaccine (3)
- 15** Persons with this disease should receive pneumococcal vaccine every five yrs (10)
- 16** Anaphylaxis antidote; abbr. (3)
- 17** Common word for hypersensitivity (7)
- 19** Organization to report vaccine preventable disease; abbr. (3)
- 20** Chronic gastrointestinal disease in children is a precaution for which vaccine (9)
- 22** Healthy nonpregnant persons under age 50 can receive this influenza vaccine (4)
- 26** Earliest age influenza vaccine may be administered (9)
- 28** Vaccine is recommended for asplenic adolescents (3)
- 30** Lethargy after vaccination (7)
- 31** Vaccine indicated for individuals who work with animal population (6)
- 32** Check for contraindications (6)
- 34** Group of individuals who may transmit influenza to high risk persons, abbr. (3)
- 35** A large vaccinated group results in this type of immunity (4)
- 37** Indicated for new parents to prevent spread of whooping cough (4)