

Table 128. Death Rates From Suicide by Selected Characteristics: 1990 to 2007

[Rates per 100,000 population. Excludes deaths of nonresidents of the United States. Beginning 2000, deaths classified according to tenth revision of International Classification of Diseases. See Appendix III]

| Characteristic | 1990 | 2000 | 2003 | 2004 | 2005 | 2006 | 2007 |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| All ages, age adjusted ¹ | 12.5 | 10.4 | 10.8 | 10.9 | 10.9 | 10.9 | 11.3 |
| All ages, crude rate ² | 12.4 | 10.4 | 10.8 | 11.0 | 11.0 | 11.1 | 11.5 |
| Under 1 year | (X) | (X) | (X) | (X) | (X) | (X) | (X) |
| 1 to 4 years | (X) | (X) | (X) | (X) | (X) | (X) | (X) |
| 5 to 14 years | 0.8 | 0.7 | 0.6 | 0.7 | 0.7 | 0.5 | 0.5 |
| 15 to 24 years | 13.2 | 10.2 | 9.7 | 10.3 | 10.0 | 9.9 | 9.7 |
| 25 to 44 years | 15.2 | 13.4 | 13.8 | 13.9 | 13.7 | 13.8 | 14.3 |
| 45 to 64 years | 15.3 | 13.5 | 15.0 | 15.4 | 15.4 | 16.0 | 16.8 |
| 65 to 74 years | 17.9 | 12.5 | 12.7 | 12.3 | 12.6 | 12.6 | 12.6 |
| 75 to 84 years | 24.9 | 17.6 | 16.4 | 16.3 | 16.9 | 15.9 | 16.3 |
| 85 years and over | 22.2 | 19.6 | 16.9 | 16.4 | 16.9 | 15.9 | 15.6 |
| AGE-ADJUSTED RATES | | | | | | | |
| Male | 21.5 | 17.7 | 18.0 | 18.0 | 18.0 | 18.0 | 18.4 |
| Female | 4.8 | 4.0 | 4.2 | 4.5 | 4.4 | 4.5 | 4.7 |
| White male | 22.8 | 19.1 | 19.6 | 19.6 | 19.6 | 19.6 | 20.2 |
| Black male | 12.8 | 10.0 | 9.2 | 9.6 | 9.2 | 9.4 | 8.8 |
| American Indian, Alaska Native male | 20.1 | 16.0 | 16.6 | 18.7 | 18.9 | 18.3 | 18.1 |
| Asian, Pacific Islander male | 9.6 | 8.6 | 8.5 | 8.4 | 7.3 | 7.9 | 9.0 |
| Hispanic male ³ | 13.7 | 10.3 | 9.7 | 9.8 | 9.4 | 8.8 | 10.1 |
| Non-Hispanic, White male ³ | 23.5 | 20.2 | 21.0 | 21.0 | 21.2 | 21.4 | 21.9 |
| White female | 5.2 | 4.3 | 4.6 | 5.0 | 4.9 | 5.1 | 5.2 |
| Black female | 2.4 | 1.8 | 1.9 | 1.8 | 1.9 | 1.4 | 1.7 |
| American Indian, Alaska Native female | 3.6 | 3.8 | 3.5 | 5.9 | 4.6 | 5.1 | 4.9 |
| Asian, Pacific Islander female | 4.1 | 2.8 | 3.1 | 3.5 | 3.3 | 3.4 | 3.5 |
| Hispanic female ³ | 2.3 | 1.7 | 1.7 | 2.0 | 1.8 | 1.8 | 1.9 |
| Non-Hispanic, White female ³ | 5.4 | 4.7 | 5.0 | 5.4 | 5.3 | 5.6 | 5.7 |

X Not applicable. ¹ Age-adjusted death rates were prepared using the direct method, in which age-specific death rates for a population of interest are applied to a standard population distributed by age. Age adjustment eliminates the differences in observed rates between points in time or among compared population groups that result from age differences in population composition. ² The total number of deaths in a given time period divided by the total resident population as of July 1.

³ Persons of Hispanic origin may be any race. Excludes data from states lacking an Hispanic-origin item on their death certificates.

Source: U.S. National Center for Health Statistics, *Health, United States, 2009*. See also <<http://www.cdc.gov/nchs/hus.htm>>.

Table 129. Death Rates From Human Immunodeficiency Virus (HIV) disease by Selected Characteristics

[Rates per 100,000 population. Excludes deaths of nonresidents of the United States. Beginning 2000, deaths classified according to tenth revision of International Classification of Diseases. See Appendix III]

| Characteristic | 1990 | 2000 | 2003 | 2004 | 2005 | 2006 | 2007 |
|--|-------------|------------|------------|------------|------------|------------|------------|
| All ages, age adjusted ¹ | 10.2 | 5.2 | 4.7 | 4.5 | 4.2 | 4.0 | 3.7 |
| All ages, crude ² | 10.1 | 5.1 | 4.7 | 4.4 | 4.2 | 4.0 | 3.7 |
| Under 1 year | 2.7 | (B) | (B) | (B) | (B) | (B) | (B) |
| 1 to 4 years | 0.8 | (B) | (B) | (B) | (B) | (B) | (B) |
| 5 to 14 years | 0.2 | 0.1 | 0.1 | 0.1 | (B) | (B) | (B) |
| 15 to 24 years | 1.5 | 0.5 | 0.4 | 0.5 | 0.4 | 0.5 | 0.4 |
| 25 to 34 years | 19.7 | 6.1 | 4.0 | 3.7 | 3.3 | 2.9 | 2.7 |
| 35 to 44 years | 27.4 | 13.1 | 12.0 | 10.9 | 9.9 | 9.2 | 8.3 |
| 45 to 54 years | 15.2 | 11.0 | 10.9 | 10.6 | 10.6 | 10.1 | 9.5 |
| 55 to 64 years | 6.2 | 5.1 | 5.4 | 5.4 | 5.3 | 5.5 | 5.3 |
| 65 to 74 years | 2.0 | 2.2 | 2.4 | 2.4 | 2.3 | 2.5 | 2.3 |
| 75 to 84 years | 0.7 | 0.7 | 0.7 | 0.8 | 0.8 | 0.8 | 0.8 |
| 85 years and over | (B) | (B) | (B) | (B) | (B) | (B) | (B) |
| AGE-ADJUSTED RATES | | | | | | | |
| Male | 18.5 | 7.9 | 7.1 | 6.6 | 6.2 | 5.9 | 5.4 |
| Female | 2.2 | 2.5 | 2.4 | 2.4 | 2.3 | 2.2 | 2.1 |
| White male | 15.7 | 4.6 | 4.2 | 3.8 | 3.6 | 3.4 | 3.1 |
| Black male | 46.3 | 35.1 | 31.3 | 29.2 | 28.2 | 26.3 | 24.5 |
| American Indian, Alaska Native male | 3.3 | 3.5 | 3.5 | 4.3 | 4.0 | 3.3 | 3.6 |
| Asian, Pacific Islander male | 4.3 | 1.2 | 1.1 | 1.2 | 1.0 | 1.1 | 0.8 |
| Hispanic male ³ | 28.8 | 10.6 | 9.2 | 8.2 | 7.5 | 7.0 | 6.3 |
| Non-Hispanic, White male ³ | 14.1 | 3.8 | 3.4 | 3.1 | 3.0 | 2.8 | 2.5 |
| White female | 1.1 | 1.0 | 0.9 | 0.9 | 0.8 | 0.7 | 0.7 |
| Black female | 10.1 | 13.2 | 12.8 | 13.0 | 12.0 | 12.2 | 11.3 |
| American Indian, Alaska Native female | (B) | 1.0 | 1.5 | 1.5 | 1.5 | 1.5 | 1.7 |
| Asian, Pacific Islander female | (B) | 0.2 | (B) | (B) | (B) | (B) | (B) |
| Hispanic female ³ | 3.8 | 2.9 | 2.7 | 2.4 | 1.9 | 1.9 | 1.8 |
| Non-Hispanic, White female ³ | 0.7 | 0.7 | 0.6 | 0.6 | 0.6 | 0.6 | 0.5 |

B Base figure too small to meet statistical standards. ¹ Age-adjusted death rates were prepared using the direct method, in which age-specific death rates for a population of interest are applied to a standard population distributed by age. Age adjustment eliminates the differences in observed rates between points in time or among compared population groups that result from age differences in population composition. ² The total number of deaths in a given time period divided by the total resident population as of July 1. ³ Persons of Hispanic origin may be any race. Excludes data from states lacking an Hispanic-origin item on their death certificates.

Source: U.S. National Center for Health Statistics, *Health, United States, 2009*. See also <<http://www.cdc.gov/nchs/hus.htm>>.