

## PSYCHIATRIC FOLLOW-UP

Since overdose is often deliberate, patients may attempt suicide by other means during the recovery phase. Psychiatric referral may be appropriate.

## PEDIATRIC MANAGEMENT

The principles of management of child and adult overdoses are similar. It is strongly recommended that the physician contact the local poison control center for specific pediatric treatment.

## DOSAGE AND ADMINISTRATION

Dosage should be initiated at a low level and increased gradually, noting carefully the clinical response and any evidence of intolerance.

### Usual Adult Dosage

Fifteen to 40 mg a day divided into 3 or 4 doses. If necessary, dosage may be increased to 60 mg a day. Dosages above this amount are not recommended. Increases should be made in the morning dose.

### Adolescent and Elderly Patients

In general, lower dosages are recommended for these patients. Five mg 3 times a day may be given initially, and increased gradually if necessary. In elderly patients, the cardiovascular system must be monitored closely if the daily dose exceeds 20 mg.

When satisfactory improvement has been reached, dosage should be reduced to the smallest amount that will maintain relief of symptoms.

Minor adverse reactions require reduction in dosage. Major adverse reactions or evidence of hypersensitivity require prompt discontinuation of the drug.

The safety and effectiveness of protriptyline in pediatric patients have not been established.

## HOW SUPPLIED

Vivactil<sup>®</sup> Tablets:

5 mg	Orange, oval, film-coated tablets in bottles of 100. Debossed OP 701
10 mg	Yellow, oval, film-coated tablets in bottles of 100. Debossed OP 702

Dispense in a tight container as defined in the USP.

Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature].

## METABOLISM

Metabolic studies indicate that protriptyline is well absorbed from the gastrointestinal tract and is rapidly sequestered in tissues.

Relatively low plasma levels are found after administration, and only a small amount of unchanged drug is excreted in the urine of dogs and rabbits. Preliminary studies indicate that demethylation of the secondary amine moiety occurs to a significant extent, and that metabolic transformation probably takes place in the liver. It penetrates the brain rapidly in mice and rats, and moreover that which is present in the brain is almost all unchanged drug.

Studies on the disposition of radioactive protriptyline in human test subjects showed significant plasma levels within 2 hours, peaking at 8 to 12 hours, then declining gradually.

Urinary excretion studies in the same subjects showed significant amounts of radioactivity in 2 hours. The rate of excretion was slow. Cumulative urinary excretion during 16 days accounted for approximately 50% of the drug. The fecal route of excretion did not seem to be important.

## MEDICATION GUIDE

### Antidepressant Medicines, Depression and other Serious Mental Illnesses, and Suicidal Thoughts or Actions

Read the Medication Guide that comes with you or your family member's antidepressant medicine. This Medication Guide is only about the risk of suicidal thoughts and actions with antidepressant medicines. **Talk to your, or your family member's, healthcare provider about:**

- all risks and benefits of treatment with antidepressant medicines
- all treatment choices for depression or other serious mental illness

**What is the most important information I should know about antidepressant medicines, depression and other serious mental illnesses, and suicidal thoughts or actions?**

1. **Antidepressant medicines may increase suicidal thoughts or actions in some children, teenagers, and young adults when the medicine is first started.**
2. **Depression and other serious mental illnesses are the most important causes of suicidal thoughts and actions. Some people may have a particularly high risk of having suicidal thoughts or actions.** These include people who have (or have a family history of) bipolar illness (also called manic-depressive illness) or suicidal thoughts or actions.
3. **How can I watch for and try to prevent suicidal thoughts and actions in myself or a family member?**
  - Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is very important when an antidepressant medicine is first started or when the dose is changed.
  - Call the healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings.
  - Keep all follow-up visits with the healthcare provider as scheduled. Call the healthcare provider between visits as needed, especially if you have concerns about symptoms.

**Call a healthcare provider right away if you or your family member has any of the following symptoms, especially if they are new, worse, or worry you:**

- thoughts about suicide or dying
- attempts to commit suicide
- new or worse depression
- new or worse anxiety
- feeling very agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

**What else do I need to know about antidepressant medicines?**

- **Never stop an antidepressant medicine without first talking to a healthcare provider.** Stopping an antidepressant medicine suddenly can cause other symptoms.
- **Antidepressants are medicines used to treat depression and other illnesses.** It is important to discuss all the risks of treating depression and also the risks of not treating it. Patients and their families or other caregivers should discuss all treatment choices with the healthcare provider, not just the use of antidepressants.
- **Antidepressant medicines have other side effects.** Talk to the healthcare provider about the side effects of the medicine prescribed for you or your family member.
- **Antidepressant medicines can interact with other medicines.** Know all of the medicines that you or your family member takes. Keep a list of all medicines to show the healthcare provider. Do not start new medicines without first checking with your healthcare provider.
- **Not all antidepressant medicines prescribed for children are FDA approved for use in children.** Talk to your child's healthcare provider for more information.

This Medication Guide has been approved by the U.S. Food and Drug Administration for all antidepressants.  
DURAMED PHARMACEUTICALS, INC.  
Subsidiary of Barr Pharmaceuticals, Inc.  
Pomona, New York 10970

