## Narrative Summary – September 2009

September 2009 was warmer than normal, averaging  $68.3^{\circ}F$ ,  $2.4^{\circ}$  above normal ( $65.9^{\circ}F$ ). The hottest September (1990) averaged 72.4°F, while the coolest (1985) averaged  $58.8^{\circ}F$ . There were 8 days in September with maximum temperatures  $\geq 90^{\circ}F$  compared to a normal of 6. There were 59 days this year with maximum temperatures  $\geq 90^{\circ}F$  compared to a normal of 52 days. The greatest number was 79 days in 1967. There were no days in September with maximum temperatures  $\geq 100^{\circ}F$  compared to a normal of <1, and a record of 3 on numerous occasions. There were 19 days this year with maximum temperatures  $\geq 100^{\circ}F$  compared to a normal of 12. The greatest was 28 days in 1958. The following temperatures records were established during September 2009:

		New	Old	
Date	Category	Record	Record	Year
16	High Minimum	65	65	1979 Tie

Precipitation for September 2009 totaled 0.06 inch, 18% of normal (0.33 inch). The wettest September (1947) received 1.34 inches, while the driest (1991 and 1999) received no precipitation. Total precipitation for 2009 (through September) is 3.42 inches, 78% of normal (4.40 inches).

The average wind speed for September 2009 was 8.4 miles per hour (mph), slightly above normal (7.9). The windiest September (1961) averaged 9.2 mph, while the September with the lightest winds (1957) averaged 5.4 mph. The peak gust for September 2009 was 46 mph on September 21. The record wind gust for September was 65 mph in 1953.

The monthly climatological data summaries, as well as other information, are available on the Internet. Address: <u>http://hms.pnl.gov/</u>

 Ken Burk
 373-3215

 HMS Staff
 373-2716

**Note:** The data in this summary pertain specifically to the Hanford Meteorology Station (HMS), which is located approximately 25 miles northwest of Richland, WA. No attempt should be made to infer meteorological conditions at other locations from these data.