





## Janis E. Hulla, PhD, DABT<sup>1</sup>; John E. Snawder, PhD, DABT<sup>2</sup>; Susan P. Proctor, DSc<sup>3</sup>; Gail D. Chapman, PhD, MBA<sup>4</sup>


1 Sacramento District of the US Army Corps of Engineers, Sacramento, CA, 2 Biomonitoring Research Team, National Institute for Occupational Safety and Health, Cincinnati, OH,

3 US Army Research Institute of Environmental Medicine, Natick, ME, 4 Commander, US Navy; Military Infectious Disease Research Program, US Army Medical Research and Materiel Command, Fort Detrick, MD The views expressed in this poster are those of the authors and do not reflect the official duties. Title 17 U.S.C. §105 provides that 'Copyright protection under this title is not available for any work of the United States Government.' Title 17 U.S.C. §101 defines a U.S. Government work as a work prepared by a military service member or employee of the U.S. Government as part of that person's official duties.


Abstract: The Department of Defense's (DOD's) Chemical and Materials Risk Management (CMRM) Directorate is using a scan-watch-action process to identify, rank and manage risks associated with emerging contaminants. Naphthalene is characterized as a likely human carcinogen by the National Toxicology Program (NTP) and in the Environmental Protection Agency's (EPA's) most recent draft health risk assessment. Thus, naphthalene-related environmental health regulations are evolving. The potential impacts have been assessed, using multi-criteria decision analysis, for five of the DOD's functional areas. One of the areas of concern is exposure to naphthalene among fuel handlers. To determine whether these exposures present unacceptable risk, the Army Research Office awarded a Small Business Innovative Research (SBIR) Project for the development of a miniature real-time naphthalene sensor. The National Institute of Occupational Safety and Health's (NIOSH's) Biomonitoring Research Team and investigators from the Army Research Institute for Environmental Medicine, University California-Davis, Temple University, and the Army Corps of Engineers are

collaborating on related projects. The research will validate the prototype sensor as a dosimeter by defining correlations between measured exposures and biomarkers of exposures to-be-collected from military fuel handlers. To date, naphthalene specificity with sensitivity of 0.5 mg/m<sup>3</sup> has been demonstrated and definition of the firmware chemometrics is underway. Implementation of the human subjects research protocol is pending institutional review boards' (IRBs') approval.


Introduction: The NTP conducted a series of rodent exposure studies between 1985 and 2002. In 2005, the EPA published a draft risk assessment that characterized naphthalene as a likely human carcinogen based on results from these and other rodent studies. The DOD CMRM Directorate subsequently assessed the potential impacts to five DOD functional areas and implemented risk management options that included funding projects that will develop and validate a naphthalene dosimeter. In addition to the DOD, the Army, National Institute of Environmental Health Sciences (NIEHS), National Science Foundation (NSF) and NIOSH are each funding components of the research. Collaborating scientists are from the Army Corps of Engineers, Army Research Office, Army Research Institute of Environmental Medicine, CDC Foundation, NIOSH, University of California-Davis and Temple University. Several other Defense organizations are represented on an interagency advisory committee.




Nanhthalenes are 1-3 % of iet fuels and DOD's annual consumption is approximately 5 billion gallons. In 2002, the National Research Council reported that 8 fuel is likely the largest chemical exposure war fighters experience.




Potential impacts to DOD's environmental restoration programs: If the EPA's draft risk assessment is finalized as written, a cancer potency value will be developed for the first time. New screening values for the evaluation of the vapor intrusion pathway will also be developed. The additional risk assessments may increase the cost to complete remediation and old sites may need to be reassessed. The EPA's health risk assessment is to be finalized in 2012.




An advisory group was chartered in 2008 in the Office of the Deputy Under Secretary of Defense for Installations and Environment. The Group will guide the development of the dosimeter and ensure best return on the agencies' investments.



Photon Systems' naphthalene dosimeter concept





Up to six different diodes may be incorporate to ensure the naphthalene-specificity illustrated by this principal component analysis.

Photon Systems' innovations include applying newly developed light emitting diodes (LEDs) to generate deep UV excitation of naphthalene's native fluorescence. This analytical approach offers the advantage of a favorable signal to noise ratio since there is little natural background fluorescence in this region of the spectrum as shown in the illustration below. Diodes being developed offer improved illumination of the target molecules.

Biomarker

Phase 1

Stationary Area Dosimeter


1/day X 5 locations X 2 days Dosimeter, Task specific


cinventional personal breathing

ationary Dosimeter in Area

rine- Morning Pre-shift, Post-

dosimeter subjects X (pre &






Conc. by prototype dosimeter

The performance of bench scale prototypes will be evaluated against GS/SM analyses of test atmospheres generated in an exposure chamber.

4-8/09:



8/09: Ancillary NIOSH/NORA proposal submitted. Goals are: a) extend the dosimeter's detection capability to include other polycyclic aromatic hydrocarbons, b) biomonitor transportation industry workers and c) develop clinical sampling capability. NIOSH funding awarded 11/09 for a 3-year project. Total values about \$850K.

2/10-10/10: Advisory Group reviews the protocol for NIOSH's


and Army's institutional review board submission and approval.

9/09: The Naphthalene Dosimeter Advisory Group proposes independent validation of the dosimeter's performance.


The National Science Foundation will provide about \$70K to Temple University investigators to do the evaluations.

are; a) field validate the prototype dosimeter and b) define the relationships between concentration of exposure biomarkers and concurrently collected dosimeter-measured exposures. Exposures measured by the prototype dosimeter will be compared to concurrent exposures measured by conventional pump-trap-purge technology.

10/10-10/12: The goals of the biomarker validation study



Conclusions: This poster outlines naphthalene risk management actions that integrate five multi-disciplinary research projects with funding from five federal agencies. The research will generate a real-time, data-logging naphthalene sensor and independently validate its performance. The dosimeter will provide new protection for industrial site workers, military fuel handlers and civilian transportation workers. The data generated will extend what is known about naphthalene's mode of action and fill the single most important data gap in the naphthalene health risk assessment. The governance efficiency to be captured is improved health protection at lower cost.



2/10-09/12: The Naphthalene Dosimeter Advisory Group anticipates providing the CMRM Directorate with progress reports, future work statements, budgets and justifications. Continued funding of the dosimeter biomarker \_validation study is to be determined through a competitive process.

Actions taken by the Army Research Office and Naphthalen

agement Directorate:

2005

8/06: The Army Research Office proposed the dosimeter concept to the SBIR Program.

6/07: The SBIR project is solicited

12/07: Two SBIR Phase I Awards, \$100K/6mo, each 9-11/08: SBIR projects are evaluated for Phase II award

Impact Assessment

2008

12/08: Photon Systems Awarded Phase II project

Phase II project, interim 9/09: Phase II project funded funding at \$125K/4mo at full rate, \$750K/2yr

University of California-Davis investigator's NIEHS-funded research renewal application is submitted 9/09. Goals include collection of nasal epithelium from military fuel handlers for analyses of biomarkers of exposure and disease progression. Work to be conducted under NIEHS IRB approval.

Naph. mg/m<sup>3</sup>

(Prototype)

Naph. mg/m<sup>3</sup>

Prototype

Trihydroxymercapturate,
Dihydroxydimethylthio, and
Trihydroxymethylthio Metabolites
(eliminated in urine)

Dosimeter Advisory Group: 9/06-09/07: Noblis, the Corps of Engineers and Action taken by the others take part in the Directorates's first Phase I **DOD Chemical &** Impact Assessment. It's on naphthalene. **Materials Risk Man-**

12/07: The CMRM Directorate forms the Naphthalene Dosimeter Advisory Group.

12/08-2/09: The Naphthalene Dosimeter Advisory Group proposes the biomarker validation study and DOD funds year 1 at \$110K. The fuel handlers study protocol is developed by these authors for research to be conducted under a NIOSH IRB approval.

The CMRM Directorate will continue to monitor to ensure that the naphthalene risk management actions are sufficient. The Directorate will continue to act as the contract officer's representative for the dosimeter validation study.

2007

2009 2010 201

Bases per Areas Days base Areas per Day day samples Samples

2012