

Table 1. *Illicit Drug Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs*

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	22,291	(21,636 - 22,960)	2,476	(2,382 - 2,573)	7,260	(7,065 - 7,460)	12,554	(11,980 - 13,150)	19,815	(19,171 - 20,475)
Northeast	4,406	(4,183 - 4,642)	448	(419 - 479)	1,507	(1,427 - 1,589)	2,451	(2,245 - 2,676)	3,958	(3,739 - 4,191)
Midwest	4,477	(4,282 - 4,677)	524	(496 - 554)	1,509	(1,430 - 1,590)	2,445	(2,262 - 2,639)	3,953	(3,762 - 4,149)
South	7,273	(6,944 - 7,617)	831	(785 - 880)	2,362	(2,235 - 2,493)	4,080	(3,767 - 4,417)	6,442	(6,122 - 6,779)
West	6,134	(5,803 - 6,478)	673	(624 - 724)	1,883	(1,774 - 1,998)	3,578	(3,277 - 3,902)	5,461	(5,136 - 5,802)
Alabama	282	(235 - 337)	35	(29 - 42)	91	(76 - 108)	156	(120 - 203)	247	(202 - 301)
Alaska	79	(67 - 92)	6	(5 - 8)	23	(20 - 26)	50	(40 - 62)	72	(61 - 85)
Arizona	510	(422 - 614)	59	(48 - 72)	140	(119 - 164)	312	(239 - 404)	452	(367 - 553)
Arkansas	185	(155 - 220)	23	(18 - 27)	54	(45 - 65)	108	(84 - 137)	162	(133 - 197)
California	3,149	(2,887 - 3,431)	370	(331 - 412)	999	(915 - 1,088)	1,781	(1,551 - 2,041)	2,780	(2,522 - 3,059)
Colorado	556	(479 - 642)	48	(40 - 58)	164	(144 - 186)	343	(277 - 422)	508	(433 - 593)
Connecticut	281	(237 - 330)	30	(25 - 36)	98	(84 - 112)	153	(118 - 197)	251	(209 - 299)
Delaware	62	(52 - 74)	8	(6 - 9)	22	(19 - 26)	32	(25 - 43)	55	(45 - 66)
District of Columbia	63	(54 - 74)	5	(4 - 6)	21	(18 - 24)	38	(30 - 48)	59	(49 - 69)
Florida	1,307	(1,184 - 1,446)	131	(115 - 148)	370	(337 - 406)	807	(694 - 936)	1,177	(1,056 - 1,314)
Georgia	578	(482 - 691)	73	(60 - 88)	187	(157 - 223)	317	(243 - 412)	504	(412 - 615)
Hawaii	102	(85 - 122)	9	(8 - 12)	31	(26 - 36)	62	(48 - 79)	92	(76 - 112)
Idaho	119	(99 - 141)	13	(11 - 16)	35	(30 - 41)	70	(54 - 91)	106	(87 - 128)
Illinois	886	(811 - 968)	107	(95 - 120)	322	(296 - 351)	457	(392 - 531)	779	(706 - 859)
Indiana	445	(375 - 526)	51	(42 - 62)	144	(123 - 168)	250	(194 - 320)	394	(327 - 473)
Iowa	146	(121 - 174)	23	(19 - 28)	54	(45 - 65)	69	(51 - 92)	123	(99 - 150)
Kansas	162	(135 - 194)	21	(17 - 25)	58	(48 - 69)	83	(63 - 109)	141	(115 - 172)
Kentucky	250	(208 - 299)	28	(23 - 34)	80	(67 - 95)	142	(109 - 184)	222	(181 - 270)
Louisiana	275	(230 - 327)	29	(24 - 36)	105	(89 - 123)	141	(108 - 184)	246	(203 - 296)
Maine	133	(114 - 155)	10	(9 - 12)	40	(35 - 45)	82	(65 - 103)	123	(104 - 144)
Maryland	383	(319 - 458)	46	(37 - 56)	132	(111 - 156)	205	(157 - 267)	337	(276 - 410)
Massachusetts	677	(587 - 777)	61	(51 - 73)	239	(211 - 268)	377	(303 - 466)	615	(528 - 714)
Michigan	862	(792 - 939)	87	(78 - 97)	249	(230 - 270)	526	(460 - 599)	775	(706 - 851)
Minnesota	305	(255 - 362)	34	(27 - 42)	109	(92 - 127)	163	(124 - 212)	271	(224 - 327)
Mississippi	165	(138 - 196)	23	(19 - 28)	56	(47 - 66)	86	(65 - 113)	142	(116 - 172)
Missouri	380	(317 - 455)	41	(33 - 49)	129	(110 - 152)	210	(161 - 273)	339	(279 - 413)
Montana	88	(75 - 102)	9	(7 - 10)	28	(25 - 32)	51	(41 - 64)	79	(67 - 94)

(continued)

Table 1. *Illicit Drug Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)*

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	103	(86 - 122)	12	(10 - 14)	38	(32 - 45)	53	(41 - 69)	91	(75 - 110)
Nevada	208	(175 - 247)	24	(20 - 29)	57	(49 - 66)	127	(99 - 162)	184	(152 - 223)
New Hampshire	137	(117 - 160)	14	(11 - 16)	45	(40 - 50)	79	(61 - 100)	123	(104 - 146)
New Jersey	531	(444 - 634)	61	(50 - 75)	177	(149 - 209)	292	(224 - 380)	470	(386 - 571)
New Mexico	165	(139 - 194)	20	(17 - 24)	52	(45 - 60)	93	(72 - 119)	145	(120 - 174)
New York	1,610	(1,470 - 1,762)	164	(146 - 184)	546	(504 - 590)	900	(776 - 1,043)	1,446	(1,309 - 1,596)
North Carolina	678	(571 - 805)	71	(58 - 86)	219	(189 - 253)	388	(301 - 498)	608	(504 - 731)
North Dakota	29	(23 - 35)	4	(3 - 4)	11	(9 - 14)	14	(10 - 19)	25	(20 - 31)
Ohio	753	(686 - 828)	96	(85 - 107)	253	(233 - 275)	404	(344 - 474)	658	(592 - 731)
Oklahoma	288	(241 - 344)	29	(24 - 35)	91	(77 - 108)	168	(129 - 216)	259	(213 - 314)
Oregon	405	(348 - 472)	37	(31 - 44)	112	(98 - 128)	256	(206 - 316)	368	(313 - 434)
Pennsylvania	846	(763 - 939)	91	(81 - 103)	295	(269 - 324)	459	(388 - 543)	755	(674 - 845)
Rhode Island	118	(101 - 137)	10	(9 - 12)	42	(37 - 47)	66	(52 - 83)	108	(91 - 126)
South Carolina	332	(277 - 397)	31	(25 - 37)	102	(87 - 118)	200	(154 - 259)	302	(248 - 366)
South Dakota	40	(33 - 49)	5	(4 - 7)	14	(11 - 17)	21	(15 - 28)	35	(28 - 44)
Tennessee	369	(306 - 444)	41	(34 - 49)	114	(95 - 135)	215	(165 - 279)	329	(268 - 402)
Texas	1,441	(1,309 - 1,585)	193	(170 - 219)	506	(459 - 557)	742	(630 - 873)	1,248	(1,120 - 1,389)
Utah	122	(99 - 147)	18	(14 - 22)	46	(37 - 57)	58	(43 - 78)	104	(83 - 129)
Vermont	74	(64 - 85)	6	(5 - 7)	25	(23 - 28)	43	(34 - 53)	68	(58 - 79)
Virginia	485	(411 - 571)	55	(44 - 67)	170	(145 - 199)	260	(202 - 332)	430	(359 - 514)
Washington	600	(516 - 696)	56	(46 - 67)	184	(160 - 211)	360	(290 - 445)	544	(462 - 639)
West Virginia	129	(108 - 154)	12	(10 - 15)	41	(35 - 47)	76	(59 - 99)	117	(97 - 142)
Wisconsin	367	(307 - 436)	44	(37 - 53)	127	(107 - 149)	196	(150 - 254)	322	(265 - 390)
Wyoming	32	(26 - 38)	4	(4 - 5)	11	(10 - 13)	16	(12 - 21)	27	(22 - 33)

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 2. Marijuana Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	28,996	(28,219 - 29,789)	3,388	(3,275 - 3,504)	10,279	(10,045 - 10,516)	15,329	(14,661 - 16,021)	25,608	(24,851 - 26,381)
Northeast	5,977	(5,718 - 6,249)	631	(595 - 668)	2,165	(2,083 - 2,249)	3,181	(2,948 - 3,433)	5,346	(5,095 - 5,611)
Midwest	5,963	(5,744 - 6,184)	700	(668 - 734)	2,200	(2,120 - 2,282)	3,062	(2,864 - 3,270)	5,262	(5,050 - 5,479)
South	9,195	(8,827 - 9,578)	1,132	(1,078 - 1,189)	3,249	(3,119 - 3,382)	4,815	(4,475 - 5,176)	8,063	(7,703 - 8,439)
West	7,861	(7,506 - 8,227)	925	(870 - 983)	2,664	(2,551 - 2,780)	4,272	(3,946 - 4,619)	6,936	(6,591 - 7,293)
Alabama	350	(298 - 411)	41	(34 - 49)	118	(102 - 136)	191	(150 - 244)	310	(260 - 368)
Alaska	97	(85 - 111)	9	(8 - 11)	30	(27 - 33)	58	(47 - 70)	88	(76 - 101)
Arizona	603	(510 - 709)	81	(68 - 96)	194	(169 - 221)	328	(254 - 422)	522	(434 - 624)
Arkansas	222	(188 - 261)	28	(23 - 33)	72	(62 - 82)	123	(95 - 157)	194	(162 - 232)
California	4,148	(3,856 - 4,459)	493	(447 - 542)	1,422	(1,332 - 1,515)	2,234	(1,970 - 2,528)	3,656	(3,372 - 3,959)
Colorado	698	(611 - 795)	73	(62 - 85)	225	(204 - 247)	400	(326 - 489)	625	(541 - 720)
Connecticut	407	(354 - 464)	48	(40 - 56)	146	(131 - 163)	213	(169 - 266)	359	(308 - 415)
Delaware	91	(79 - 105)	11	(10 - 13)	34	(30 - 37)	46	(36 - 59)	80	(68 - 93)
District of Columbia	83	(72 - 95)	6	(5 - 7)	30	(26 - 33)	47	(38 - 58)	77	(67 - 89)
Florida	1,671	(1,536 - 1,820)	189	(171 - 207)	530	(493 - 569)	952	(830 - 1,090)	1,482	(1,351 - 1,628)
Georgia	824	(702 - 965)	109	(91 - 129)	276	(240 - 314)	439	(341 - 562)	715	(599 - 852)
Hawaii	122	(104 - 143)	13	(10 - 15)	41	(36 - 47)	68	(53 - 86)	109	(92 - 129)
Idaho	153	(132 - 176)	18	(16 - 22)	51	(45 - 57)	83	(67 - 104)	134	(115 - 157)
Illinois	1,196	(1,108 - 1,289)	141	(128 - 156)	455	(426 - 485)	599	(524 - 685)	1,054	(970 - 1,146)
Indiana	537	(462 - 622)	65	(55 - 77)	198	(174 - 224)	274	(215 - 346)	472	(401 - 554)
Iowa	222	(187 - 260)	31	(26 - 37)	86	(74 - 98)	105	(80 - 138)	191	(158 - 227)
Kansas	222	(191 - 258)	29	(24 - 34)	89	(78 - 102)	105	(81 - 134)	194	(164 - 228)
Kentucky	289	(245 - 339)	37	(31 - 45)	102	(88 - 118)	149	(115 - 193)	252	(210 - 300)
Louisiana	319	(272 - 373)	39	(32 - 47)	127	(111 - 145)	153	(116 - 200)	280	(235 - 332)
Maine	166	(145 - 189)	15	(12 - 17)	54	(49 - 59)	97	(79 - 119)	151	(131 - 173)
Maryland	476	(405 - 558)	62	(52 - 74)	191	(166 - 217)	224	(169 - 294)	414	(347 - 493)
Massachusetts	934	(831 - 1,048)	86	(73 - 100)	339	(309 - 370)	509	(422 - 612)	848	(748 - 959)
Michigan	1,085	(1,006 - 1,170)	118	(107 - 129)	358	(336 - 380)	610	(539 - 689)	968	(891 - 1,051)
Minnesota	464	(402 - 532)	49	(41 - 58)	175	(156 - 196)	240	(190 - 302)	415	(355 - 482)
Mississippi	202	(171 - 236)	27	(22 - 32)	73	(63 - 84)	102	(79 - 132)	175	(146 - 208)
Missouri	505	(431 - 590)	58	(48 - 69)	178	(156 - 201)	269	(210 - 344)	447	(377 - 529)
Montana	119	(105 - 136)	12	(11 - 14)	41	(37 - 46)	66	(53 - 81)	107	(93 - 123)

(continued)

Table 2. Marijuana Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	140	(120 - 162)	15	(13 - 19)	56	(49 - 64)	68	(53 - 87)	124	(106 - 146)
Nevada	242	(206 - 283)	32	(27 - 38)	72	(63 - 82)	138	(108 - 175)	210	(176 - 249)
New Hampshire	172	(151 - 196)	19	(16 - 22)	59	(53 - 64)	95	(76 - 117)	153	(132 - 176)
New Jersey	770	(662 - 896)	95	(79 - 112)	278	(244 - 313)	398	(311 - 507)	676	(572 - 796)
New Mexico	205	(178 - 234)	29	(25 - 34)	73	(65 - 82)	102	(80 - 130)	175	(150 - 204)
New York	2,132	(1,974 - 2,300)	216	(195 - 239)	773	(727 - 821)	1,142	(1,003 - 1,298)	1,915	(1,763 - 2,079)
North Carolina	815	(705 - 940)	90	(75 - 106)	299	(265 - 335)	427	(337 - 538)	725	(620 - 848)
North Dakota	45	(38 - 52)	5	(4 - 6)	21	(18 - 24)	19	(15 - 25)	40	(34 - 47)
Ohio	993	(920 - 1,072)	121	(110 - 134)	372	(350 - 396)	499	(434 - 573)	871	(801 - 949)
Oklahoma	331	(281 - 389)	41	(34 - 48)	117	(101 - 134)	173	(135 - 222)	290	(242 - 346)
Oregon	507	(444 - 578)	56	(49 - 64)	156	(140 - 173)	294	(240 - 359)	451	(390 - 521)
Pennsylvania	1,162	(1,069 - 1,263)	131	(118 - 145)	428	(399 - 457)	603	(522 - 695)	1,031	(941 - 1,129)
Rhode Island	144	(126 - 164)	14	(12 - 16)	55	(50 - 61)	75	(60 - 93)	130	(113 - 150)
South Carolina	394	(335 - 461)	43	(36 - 50)	138	(121 - 157)	213	(165 - 272)	351	(295 - 417)
South Dakota	58	(50 - 68)	7	(6 - 9)	24	(21 - 28)	27	(21 - 35)	51	(43 - 61)
Tennessee	496	(421 - 581)	54	(46 - 65)	168	(146 - 192)	274	(214 - 349)	442	(370 - 524)
Texas	1,808	(1,658 - 1,971)	263	(237 - 292)	671	(619 - 725)	875	(747 - 1,022)	1,545	(1,401 - 1,702)
Utah	150	(127 - 177)	24	(19 - 29)	70	(59 - 83)	57	(41 - 78)	127	(105 - 152)
Vermont	91	(81 - 103)	8	(7 - 9)	34	(31 - 37)	50	(40 - 60)	83	(73 - 94)
Virginia	666	(575 - 771)	78	(66 - 93)	254	(224 - 286)	334	(261 - 425)	588	(500 - 689)
Washington	775	(678 - 885)	79	(67 - 93)	272	(245 - 300)	424	(342 - 523)	696	(602 - 803)
West Virginia	158	(136 - 185)	16	(13 - 19)	50	(44 - 57)	92	(73 - 115)	142	(120 - 168)
Wisconsin	495	(427 - 573)	61	(52 - 72)	188	(166 - 212)	246	(193 - 312)	434	(369 - 509)
Wyoming	41	(35 - 48)	6	(5 - 7)	16	(14 - 18)	20	(15 - 26)	36	(30 - 42)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 3. Marijuana Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	17,119	(16,556 - 17,696)	1,808	(1,726 - 1,893)	6,220	(6,036 - 6,410)	9,091	(8,609 - 9,595)	15,311	(14,761 - 15,877)
Northeast	3,506	(3,312 - 3,712)	360	(334 - 388)	1,337	(1,261 - 1,416)	1,809	(1,639 - 1,996)	3,146	(2,957 - 3,348)
Midwest	3,505	(3,335 - 3,680)	383	(359 - 408)	1,302	(1,226 - 1,381)	1,820	(1,672 - 1,979)	3,122	(2,957 - 3,292)
South	5,304	(5,033 - 5,589)	578	(539 - 620)	1,977	(1,856 - 2,103)	2,749	(2,509 - 3,010)	4,726	(4,465 - 5,002)
West	4,804	(4,515 - 5,107)	487	(448 - 529)	1,605	(1,502 - 1,713)	2,713	(2,457 - 2,992)	4,317	(4,036 - 4,614)
Alabama	192	(155 - 237)	22	(17 - 27)	72	(58 - 88)	99	(72 - 134)	170	(135 - 214)
Alaska	66	(55 - 78)	5	(4 - 6)	20	(17 - 23)	41	(32 - 52)	61	(50 - 73)
Arizona	349	(281 - 431)	41	(32 - 52)	111	(92 - 134)	197	(144 - 268)	308	(244 - 387)
Arkansas	129	(105 - 159)	15	(11 - 19)	42	(34 - 51)	73	(54 - 97)	114	(91 - 143)
California	2,487	(2,254 - 2,740)	267	(234 - 303)	863	(785 - 946)	1,358	(1,160 - 1,587)	2,221	(1,994 - 2,469)
Colorado	465	(394 - 549)	38	(30 - 47)	145	(126 - 165)	283	(222 - 358)	428	(358 - 509)
Connecticut	227	(188 - 272)	25	(20 - 31)	88	(76 - 102)	113	(83 - 153)	202	(164 - 246)
Delaware	51	(42 - 62)	6	(5 - 7)	20	(17 - 23)	26	(19 - 35)	45	(37 - 56)
District of Columbia	52	(44 - 61)	3	(3 - 4)	18	(15 - 22)	30	(23 - 39)	48	(41 - 58)
Florida	947	(847 - 1,061)	92	(80 - 106)	319	(287 - 353)	535	(448 - 639)	854	(757 - 966)
Georgia	434	(352 - 534)	53	(41 - 67)	157	(130 - 188)	224	(164 - 307)	381	(304 - 477)
Hawaii	83	(67 - 102)	7	(6 - 9)	26	(22 - 31)	49	(37 - 66)	75	(60 - 94)
Idaho	91	(74 - 110)	10	(8 - 13)	28	(24 - 33)	52	(39 - 69)	80	(65 - 100)
Illinois	690	(627 - 758)	79	(69 - 91)	274	(250 - 300)	336	(284 - 397)	610	(550 - 676)
Indiana	321	(266 - 387)	35	(28 - 44)	123	(103 - 146)	164	(123 - 217)	286	(234 - 350)
Iowa	123	(100 - 151)	16	(13 - 21)	50	(41 - 61)	57	(41 - 79)	107	(85 - 133)
Kansas	122	(99 - 150)	15	(11 - 19)	47	(38 - 57)	61	(45 - 83)	107	(86 - 134)
Kentucky	163	(132 - 201)	18	(14 - 24)	64	(52 - 77)	81	(59 - 112)	145	(116 - 181)
Louisiana	190	(155 - 232)	19	(15 - 25)	83	(70 - 99)	87	(63 - 121)	170	(137 - 211)
Maine	110	(92 - 131)	9	(7 - 11)	35	(31 - 40)	66	(51 - 86)	102	(84 - 122)
Maryland	288	(235 - 353)	32	(25 - 41)	117	(98 - 139)	139	(101 - 191)	256	(206 - 319)
Massachusetts	564	(486 - 654)	51	(42 - 62)	214	(188 - 242)	299	(237 - 376)	514	(438 - 601)
Michigan	701	(635 - 773)	67	(58 - 76)	220	(201 - 240)	414	(356 - 481)	634	(570 - 705)
Minnesota	259	(213 - 314)	25	(20 - 32)	98	(82 - 116)	136	(102 - 183)	234	(190 - 287)
Mississippi	111	(89 - 138)	13	(10 - 17)	42	(34 - 52)	56	(41 - 78)	98	(78 - 124)
Missouri	303	(247 - 370)	31	(24 - 39)	112	(94 - 133)	160	(118 - 216)	272	(219 - 337)
Montana	75	(64 - 89)	7	(6 - 9)	25	(22 - 29)	43	(34 - 56)	68	(57 - 82)

(continued)

Table 3. Marijuana Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	86	(71 - 104)	9	(7 - 11)	33	(27 - 40)	44	(33 - 59)	77	(63 - 95)
Nevada	151	(124 - 183)	16	(13 - 20)	47	(40 - 55)	88	(66 - 116)	134	(109 - 166)
New Hampshire	112	(94 - 133)	12	(10 - 15)	40	(35 - 45)	61	(46 - 80)	100	(83 - 121)
New Jersey	410	(339 - 498)	50	(40 - 63)	157	(131 - 187)	203	(151 - 273)	360	(292 - 444)
New Mexico	126	(105 - 152)	15	(12 - 19)	44	(37 - 52)	67	(50 - 90)	111	(91 - 136)
New York	1,270	(1,151 - 1,400)	130	(114 - 147)	480	(439 - 524)	660	(559 - 779)	1,140	(1,025 - 1,268)
North Carolina	540	(447 - 652)	51	(41 - 64)	195	(166 - 226)	294	(221 - 390)	489	(399 - 598)
North Dakota	22	(18 - 27)	2	(2 - 3)	9	(7 - 12)	11	(8 - 15)	20	(16 - 25)
Ohio	577	(522 - 638)	70	(62 - 80)	223	(203 - 243)	284	(237 - 339)	506	(454 - 566)
Oklahoma	206	(166 - 254)	21	(16 - 27)	73	(60 - 88)	112	(82 - 152)	185	(147 - 232)
Oregon	328	(276 - 389)	27	(22 - 33)	96	(83 - 110)	205	(160 - 261)	300	(250 - 361)
Pennsylvania	653	(586 - 729)	71	(62 - 82)	261	(236 - 288)	321	(266 - 386)	582	(517 - 655)
Rhode Island	96	(82 - 113)	8	(6 - 9)	39	(34 - 44)	50	(39 - 65)	89	(75 - 105)
South Carolina	224	(183 - 274)	20	(16 - 26)	85	(70 - 101)	119	(88 - 161)	204	(165 - 252)
South Dakota	33	(26 - 41)	4	(3 - 5)	12	(10 - 15)	17	(12 - 23)	29	(23 - 37)
Tennessee	244	(197 - 300)	26	(21 - 33)	92	(75 - 111)	126	(92 - 172)	218	(173 - 272)
Texas	1,063	(953 - 1,184)	138	(119 - 159)	415	(371 - 463)	511	(423 - 616)	926	(821 - 1,042)
Utah	68	(53 - 86)	9	(7 - 13)	29	(22 - 39)	29	(20 - 42)	58	(44 - 76)
Vermont	62	(53 - 73)	5	(4 - 6)	23	(21 - 26)	34	(27 - 43)	57	(49 - 67)
Virginia	377	(311 - 455)	40	(32 - 50)	151	(127 - 177)	186	(138 - 249)	336	(274 - 412)
Washington	492	(414 - 582)	41	(33 - 51)	161	(138 - 186)	289	(226 - 368)	450	(375 - 538)
West Virginia	92	(75 - 113)	8	(6 - 10)	35	(29 - 41)	49	(36 - 67)	84	(68 - 104)
Wisconsin	268	(218 - 327)	30	(24 - 38)	100	(83 - 120)	137	(101 - 186)	237	(191 - 295)
Wyoming	24	(20 - 29)	3	(2 - 4)	10	(8 - 11)	11	(8 - 16)	21	(17 - 26)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 4. Perceptions of Great Risk of Smoking Marijuana Once a Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	84,840	(83,479 - 86,212)	7,327	(7,185 - 7,471)	6,507	(6,323 - 6,696)	71,006	(69,685 - 72,337)	77,513	(76,166 - 78,871)
Northeast	14,568	(14,096 - 15,045)	1,118	(1,070 - 1,167)	973	(914 - 1,035)	12,477	(12,019 - 12,941)	13,450	(12,982 - 13,924)
Midwest	17,656	(17,214 - 18,104)	1,670	(1,622 - 1,719)	1,297	(1,237 - 1,358)	14,689	(14,267 - 15,116)	15,985	(15,550 - 16,428)
South	35,135	(34,367 - 35,911)	2,943	(2,861 - 3,027)	2,707	(2,599 - 2,818)	29,484	(28,743 - 30,233)	32,191	(31,433 - 32,958)
West	17,482	(16,879 - 18,099)	1,596	(1,528 - 1,665)	1,531	(1,439 - 1,626)	14,356	(13,777 - 14,949)	15,886	(15,288 - 16,500)
Alabama	1,593	(1,468 - 1,721)	151	(137 - 165)	147	(129 - 166)	1,295	(1,176 - 1,417)	1,442	(1,319 - 1,568)
Alaska	135	(119 - 153)	15	(13 - 17)	12	(10 - 14)	108	(93 - 125)	120	(104 - 138)
Arizona	1,710	(1,536 - 1,896)	146	(126 - 167)	150	(129 - 173)	1,415	(1,247 - 1,593)	1,565	(1,394 - 1,747)
Arkansas	898	(820 - 979)	79	(70 - 88)	64	(55 - 74)	755	(681 - 831)	819	(743 - 898)
California	9,430	(8,948 - 9,926)	811	(755 - 869)	875	(802 - 952)	7,745	(7,281 - 8,223)	8,619	(8,141 - 9,112)
Colorado	921	(810 - 1,041)	88	(76 - 102)	73	(60 - 87)	760	(656 - 875)	832	(725 - 951)
Connecticut	922	(829 - 1,022)	67	(57 - 77)	58	(49 - 68)	798	(708 - 893)	856	(764 - 954)
Delaware	225	(202 - 250)	19	(17 - 22)	15	(12 - 17)	192	(170 - 215)	206	(184 - 230)
District of Columbia	138	(123 - 153)	8	(7 - 10)	13	(11 - 16)	116	(102 - 131)	129	(115 - 145)
Florida	6,338	(6,066 - 6,613)	426	(400 - 453)	403	(372 - 436)	5,509	(5,244 - 5,778)	5,912	(5,641 - 6,185)
Georgia	2,793	(2,548 - 3,046)	248	(220 - 278)	224	(194 - 256)	2,321	(2,091 - 2,560)	2,545	(2,308 - 2,792)
Hawaii	333	(301 - 368)	24	(21 - 28)	25	(21 - 29)	285	(253 - 317)	309	(277 - 343)
Idaho	412	(375 - 450)	43	(38 - 48)	36	(31 - 41)	334	(299 - 370)	369	(334 - 407)
Illinois	3,482	(3,313 - 3,654)	313	(293 - 334)	273	(250 - 297)	2,896	(2,733 - 3,063)	3,169	(3,002 - 3,340)
Indiana	1,751	(1,600 - 1,909)	180	(162 - 199)	125	(108 - 145)	1,446	(1,302 - 1,596)	1,571	(1,423 - 1,726)
Iowa	828	(751 - 911)	79	(71 - 87)	62	(53 - 72)	687	(613 - 765)	749	(674 - 830)
Kansas	797	(725 - 874)	72	(64 - 80)	59	(51 - 69)	666	(598 - 738)	726	(655 - 800)
Kentucky	1,358	(1,247 - 1,474)	132	(119 - 146)	108	(94 - 123)	1,119	(1,013 - 1,228)	1,226	(1,118 - 1,340)
Louisiana	1,480	(1,364 - 1,600)	129	(116 - 143)	130	(114 - 146)	1,221	(1,111 - 1,333)	1,351	(1,237 - 1,468)
Maine	284	(251 - 318)	26	(22 - 29)	16	(14 - 20)	242	(211 - 275)	258	(226 - 292)
Maryland	1,559	(1,394 - 1,733)	120	(104 - 138)	103	(87 - 122)	1,335	(1,177 - 1,501)	1,439	(1,277 - 1,610)
Massachusetts	1,394	(1,237 - 1,564)	101	(86 - 117)	84	(69 - 103)	1,209	(1,058 - 1,372)	1,293	(1,139 - 1,460)
Michigan	2,443	(2,313 - 2,577)	233	(219 - 248)	178	(162 - 195)	2,032	(1,905 - 2,163)	2,210	(2,081 - 2,343)
Minnesota	1,211	(1,085 - 1,346)	119	(105 - 134)	92	(79 - 107)	1,000	(881 - 1,128)	1,092	(970 - 1,224)
Mississippi	1,070	(994 - 1,148)	104	(93 - 114)	94	(83 - 106)	873	(801 - 946)	967	(893 - 1,043)
Missouri	1,687	(1,538 - 1,843)	151	(135 - 168)	124	(107 - 142)	1,413	(1,270 - 1,563)	1,536	(1,390 - 1,690)
Montana	225	(202 - 250)	19	(17 - 22)	15	(12 - 17)	192	(170 - 215)	206	(184 - 230)

(continued)

Table 4. Perceptions of Great Risk of Smoking Marijuana Once a Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	504	(458 - 552)	52	(46 - 57)	38	(33 - 44)	414	(371 - 460)	452	(408 - 500)
Nevada	680	(613 - 751)	63	(55 - 72)	50	(43 - 58)	568	(504 - 635)	617	(552 - 687)
New Hampshire	271	(240 - 305)	23	(20 - 27)	15	(13 - 19)	233	(202 - 266)	248	(217 - 282)
New Jersey	2,472	(2,241 - 2,713)	187	(164 - 211)	149	(126 - 175)	2,137	(1,917 - 2,365)	2,285	(2,060 - 2,521)
New Mexico	591	(538 - 647)	43	(37 - 49)	44	(38 - 51)	505	(453 - 558)	548	(496 - 603)
New York	5,373	(5,098 - 5,654)	401	(373 - 432)	401	(365 - 439)	4,571	(4,306 - 4,843)	4,972	(4,701 - 5,249)
North Carolina	2,547	(2,313 - 2,793)	199	(175 - 225)	190	(163 - 219)	2,159	(1,934 - 2,394)	2,348	(2,117 - 2,590)
North Dakota	192	(176 - 209)	17	(15 - 19)	19	(16 - 21)	157	(142 - 172)	175	(160 - 192)
Ohio	3,111	(2,949 - 3,276)	290	(273 - 308)	209	(192 - 227)	2,613	(2,455 - 2,775)	2,821	(2,662 - 2,985)
Oklahoma	1,087	(992 - 1,186)	97	(86 - 109)	86	(74 - 100)	903	(813 - 997)	990	(896 - 1,087)
Oregon	730	(643 - 825)	70	(61 - 80)	47	(39 - 57)	613	(531 - 704)	661	(576 - 753)
Pennsylvania	3,490	(3,302 - 3,682)	283	(264 - 303)	226	(205 - 249)	2,980	(2,798 - 3,167)	3,207	(3,021 - 3,397)
Rhode Island	240	(213 - 270)	20	(17 - 23)	16	(13 - 19)	205	(179 - 233)	221	(194 - 250)
South Carolina	1,445	(1,319 - 1,576)	109	(97 - 122)	100	(87 - 116)	1,236	(1,115 - 1,360)	1,336	(1,212 - 1,464)
South Dakota	237	(216 - 260)	22	(19 - 24)	17	(15 - 20)	198	(178 - 219)	216	(195 - 238)
Tennessee	1,931	(1,752 - 2,121)	157	(141 - 174)	121	(103 - 140)	1,654	(1,480 - 1,834)	1,774	(1,597 - 1,960)
Texas	7,953	(7,610 - 8,299)	757	(714 - 802)	712	(664 - 763)	6,483	(6,150 - 6,819)	7,195	(6,858 - 7,537)
Utah	813	(750 - 878)	114	(104 - 124)	96	(84 - 109)	603	(545 - 663)	699	(638 - 762)
Vermont	122	(107 - 139)	11	(9 - 12)	8	(6 - 9)	104	(89 - 120)	111	(97 - 128)
Virginia	2,103	(1,902 - 2,315)	161	(142 - 182)	159	(136 - 185)	1,783	(1,590 - 1,985)	1,942	(1,744 - 2,150)
Washington	1,349	(1,192 - 1,518)	145	(128 - 163)	97	(82 - 115)	1,107	(959 - 1,269)	1,204	(1,052 - 1,370)
West Virginia	616	(566 - 668)	47	(42 - 52)	39	(34 - 44)	531	(482 - 581)	569	(520 - 620)
Wisconsin	1,411	(1,274 - 1,557)	143	(129 - 159)	101	(86 - 118)	1,167	(1,036 - 1,306)	1,268	(1,133 - 1,411)
Wyoming	151	(137 - 165)	15	(14 - 17)	13	(11 - 15)	123	(110 - 137)	136	(122 - 150)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 5. Average Annual Marijuana Initiates, by Age Group and State: Estimated Numbers (in Thousands), Based on 2009 and 2010 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	2,728	(2,632 - 2,827)	1,279	(1,231 - 1,328)	1,277	(1,218 - 1,338)	173	(135 - 221)	1,449	(1,371 - 1,531)
Northeast	509	(478 - 541)	233	(217 - 249)	246	(226 - 268)	30	(20 - 43)	276	(251 - 303)
Midwest	591	(561 - 622)	273	(259 - 289)	285	(266 - 306)	32	(22 - 46)	317	(292 - 344)
South	937	(889 - 988)	434	(410 - 458)	443	(412 - 477)	60	(43 - 85)	504	(463 - 548)
West	691	(649 - 737)	339	(316 - 363)	302	(274 - 332)	50	(34 - 75)	352	(317 - 392)
Alabama	35	(30 - 42)	16	(13 - 19)	17	(13 - 22)	2	(1 - 4)	20	(15 - 25)
Alaska	6	(5 - 8)	3	(3 - 4)	3	(2 - 4)	0	(0 - 1)	3	(3 - 4)
Arizona	56	(48 - 66)	28	(23 - 35)	24	(19 - 30)	4	(2 - 7)	28	(22 - 35)
Arkansas	22	(19 - 26)	10	(8 - 13)	10	(8 - 13)	1	(1 - 3)	12	(9 - 15)
California	378	(345 - 416)	185	(166 - 206)	164	(142 - 189)	29	(17 - 49)	193	(166 - 225)
Colorado	52	(45 - 61)	28	(23 - 33)	22	(17 - 27)	3	(2 - 6)	25	(20 - 31)
Connecticut	37	(31 - 43)	17	(14 - 21)	18	(14 - 22)	2	(1 - 4)	20	(16 - 24)
Delaware	8	(7 - 10)	4	(3 - 5)	4	(3 - 5)	0	(0 - 1)	4	(4 - 5)
District of Columbia	6	(5 - 7)	2	(2 - 3)	4	(3 - 5)	0	(0 - 1)	4	(3 - 5)
Florida	146	(132 - 161)	68	(61 - 77)	66	(57 - 76)	12	(7 - 19)	78	(66 - 90)
Georgia	84	(72 - 98)	42	(35 - 51)	37	(30 - 46)	5	(3 - 9)	42	(34 - 52)
Hawaii	10	(9 - 12)	4	(3 - 6)	5	(4 - 6)	1	(0 - 2)	6	(5 - 8)
Idaho	14	(12 - 17)	7	(6 - 8)	7	(5 - 8)	1	(0 - 1)	8	(6 - 9)
Illinois	116	(105 - 127)	52	(46 - 58)	58	(50 - 66)	6	(4 - 10)	64	(56 - 73)
Indiana	56	(48 - 64)	27	(22 - 32)	26	(21 - 32)	3	(2 - 5)	29	(23 - 35)
Iowa	28	(24 - 32)	13	(10 - 15)	14	(11 - 17)	1	(1 - 3)	15	(12 - 19)
Kansas	25	(22 - 30)	12	(9 - 14)	12	(10 - 15)	2	(1 - 3)	14	(11 - 17)
Kentucky	34	(29 - 39)	16	(13 - 20)	16	(13 - 19)	2	(1 - 4)	18	(15 - 22)
Louisiana	35	(30 - 41)	15	(12 - 19)	17	(14 - 21)	2	(1 - 5)	20	(16 - 24)
Maine	10	(9 - 12)	5	(4 - 6)	5	(4 - 6)	1	(0 - 1)	5	(4 - 7)
Maryland	50	(42 - 58)	22	(18 - 27)	25	(20 - 31)	2	(1 - 5)	28	(22 - 34)
Massachusetts	64	(55 - 75)	32	(26 - 38)	29	(23 - 36)	4	(2 - 7)	32	(26 - 40)
Michigan	96	(87 - 105)	43	(39 - 48)	48	(42 - 55)	4	(3 - 8)	52	(46 - 60)
Minnesota	46	(39 - 53)	20	(17 - 25)	23	(18 - 28)	3	(1 - 5)	25	(20 - 31)
Mississippi	23	(19 - 27)	10	(8 - 12)	11	(9 - 14)	1	(1 - 3)	13	(10 - 16)
Missouri	48	(41 - 56)	24	(20 - 29)	21	(17 - 26)	3	(2 - 6)	24	(19 - 30)
Montana	11	(9 - 12)	5	(4 - 5)	6	(5 - 7)	1	(0 - 1)	6	(5 - 7)

(continued)

Table 5. Average Annual Marijuana Initiates, by Age Group and State: Estimated Numbers (in Thousands), Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	15	(13 - 18)	6	(5 - 8)	8	(6 - 10)	1	(0 - 2)	9	(7 - 11)
Nevada	21	(18 - 25)	11	(9 - 14)	8	(6 - 10)	2	(1 - 3)	10	(8 - 12)
New Hampshire	13	(11 - 15)	6	(5 - 7)	6	(5 - 7)	1	(0 - 1)	7	(5 - 8)
New Jersey	81	(70 - 95)	38	(31 - 46)	38	(31 - 48)	5	(3 - 10)	43	(35 - 54)
New Mexico	20	(17 - 23)	10	(8 - 12)	8	(6 - 10)	2	(1 - 3)	9	(7 - 12)
New York	172	(157 - 190)	76	(68 - 86)	85	(74 - 97)	11	(7 - 18)	96	(84 - 110)
North Carolina	75	(64 - 87)	36	(29 - 44)	34	(27 - 43)	4	(2 - 8)	39	(31 - 48)
North Dakota	6	(5 - 7)	2	(2 - 3)	3	(3 - 4)	0	(0 - 1)	4	(3 - 5)
Ohio	99	(91 - 109)	47	(42 - 52)	47	(41 - 53)	6	(3 - 9)	53	(46 - 60)
Oklahoma	30	(25 - 35)	15	(12 - 18)	13	(10 - 17)	2	(1 - 4)	15	(12 - 19)
Oregon	35	(30 - 40)	17	(14 - 21)	15	(12 - 19)	2	(1 - 5)	17	(14 - 22)
Pennsylvania	114	(104 - 125)	50	(45 - 57)	58	(50 - 67)	6	(3 - 10)	64	(55 - 73)
Rhode Island	11	(9 - 13)	5	(4 - 6)	5	(4 - 6)	1	(0 - 2)	6	(5 - 7)
South Carolina	37	(32 - 44)	17	(14 - 20)	18	(15 - 23)	2	(1 - 4)	21	(17 - 26)
South Dakota	7	(6 - 9)	3	(2 - 4)	4	(3 - 5)	0	(0 - 1)	4	(3 - 5)
Tennessee	46	(39 - 54)	20	(16 - 24)	23	(19 - 29)	3	(2 - 6)	26	(21 - 32)
Texas	222	(203 - 244)	104	(93 - 117)	104	(91 - 119)	14	(8 - 23)	118	(103 - 136)
Utah	25	(21 - 29)	12	(9 - 14)	11	(9 - 15)	2	(1 - 3)	13	(10 - 17)
Vermont	7	(6 - 8)	3	(3 - 4)	3	(3 - 4)	0	(0 - 1)	4	(3 - 4)
Virginia	71	(61 - 82)	30	(25 - 36)	36	(30 - 45)	4	(2 - 7)	40	(33 - 49)
Washington	57	(49 - 66)	26	(22 - 32)	27	(22 - 33)	4	(2 - 7)	31	(25 - 38)
West Virginia	14	(12 - 16)	7	(5 - 8)	6	(5 - 8)	1	(1 - 2)	7	(6 - 9)
Wisconsin	49	(42 - 57)	25	(20 - 30)	22	(17 - 28)	3	(1 - 5)	24	(20 - 31)
Wyoming	5	(5 - 6)	2	(2 - 3)	3	(2 - 3)	0	(0 - 1)	3	(2 - 4)

NOTE: Average annual number of marijuana initiates = $X_1 \div 2$, where X_1 is the number of marijuana initiates in the past 24 months.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 6. *Illicit Drug Use Other Than Marijuana in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs*

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	9,106	(8,707 - 9,521)	1,107	(1,044 - 1,174)	2,760	(2,641 - 2,885)	5,238	(4,882 - 5,620)	7,999	(7,608 - 8,408)
Northeast	1,696	(1,555 - 1,850)	172	(155 - 191)	545	(503 - 590)	980	(856 - 1,121)	1,524	(1,385 - 1,678)
Midwest	1,832	(1,705 - 1,967)	241	(223 - 262)	593	(555 - 634)	998	(887 - 1,121)	1,591	(1,466 - 1,725)
South	3,247	(3,037 - 3,471)	402	(372 - 434)	928	(868 - 992)	1,916	(1,727 - 2,126)	2,845	(2,637 - 3,069)
West	2,330	(2,126 - 2,554)	292	(263 - 324)	694	(636 - 757)	1,345	(1,164 - 1,552)	2,038	(1,837 - 2,261)
Alabama	129	(100 - 165)	19	(15 - 24)	39	(30 - 51)	71	(49 - 103)	110	(82 - 147)
Alaska	22	(17 - 29)	2	(2 - 3)	6	(5 - 8)	13	(9 - 19)	20	(15 - 27)
Arizona	231	(178 - 299)	26	(20 - 34)	60	(47 - 77)	144	(100 - 208)	205	(152 - 274)
Arkansas	95	(74 - 120)	12	(9 - 15)	26	(20 - 33)	57	(40 - 80)	83	(63 - 109)
California	1,148	(994 - 1,323)	162	(139 - 189)	346	(300 - 398)	640	(510 - 803)	986	(836 - 1,161)
Colorado	172	(136 - 217)	18	(14 - 23)	54	(43 - 66)	101	(72 - 141)	154	(119 - 200)
Connecticut	111	(86 - 144)	12	(9 - 15)	34	(27 - 43)	65	(45 - 95)	99	(74 - 132)
Delaware	27	(21 - 34)	3	(2 - 4)	8	(6 - 10)	15	(11 - 22)	24	(18 - 31)
District of Columbia	24	(18 - 31)	2	(1 - 2)	7	(5 - 9)	16	(11 - 22)	22	(17 - 30)
Florida	539	(464 - 627)	60	(51 - 70)	132	(113 - 154)	347	(280 - 430)	479	(405 - 567)
Georgia	257	(200 - 331)	33	(26 - 42)	76	(59 - 98)	148	(102 - 216)	224	(168 - 300)
Hawaii	37	(28 - 49)	4	(3 - 6)	10	(8 - 14)	23	(16 - 34)	33	(24 - 45)
Idaho	49	(38 - 61)	6	(5 - 8)	15	(12 - 18)	28	(20 - 40)	42	(32 - 56)
Illinois	347	(302 - 400)	49	(41 - 57)	117	(102 - 134)	182	(143 - 230)	299	(254 - 351)
Indiana	182	(142 - 232)	25	(20 - 32)	55	(43 - 71)	101	(70 - 146)	157	(119 - 207)
Iowa	60	(45 - 78)	12	(9 - 15)	20	(15 - 27)	28	(17 - 45)	48	(34 - 68)
Kansas	78	(61 - 99)	10	(8 - 13)	27	(21 - 34)	41	(28 - 59)	68	(52 - 89)
Kentucky	122	(96 - 155)	15	(12 - 20)	36	(29 - 46)	70	(49 - 100)	106	(81 - 140)
Louisiana	151	(120 - 190)	15	(12 - 20)	50	(40 - 63)	85	(60 - 120)	136	(105 - 174)
Maine	41	(33 - 52)	4	(3 - 5)	14	(11 - 17)	23	(16 - 33)	37	(29 - 48)
Maryland	151	(113 - 202)	19	(15 - 25)	40	(29 - 53)	92	(61 - 138)	132	(94 - 183)
Massachusetts	237	(189 - 296)	21	(16 - 27)	86	(69 - 106)	130	(92 - 183)	216	(168 - 276)
Michigan	313	(274 - 359)	36	(31 - 42)	94	(82 - 106)	183	(149 - 226)	277	(238 - 322)
Minnesota	124	(95 - 161)	15	(11 - 19)	41	(32 - 53)	68	(45 - 101)	109	(81 - 147)
Mississippi	80	(63 - 102)	14	(11 - 17)	26	(20 - 33)	41	(28 - 60)	67	(50 - 88)
Missouri	147	(114 - 191)	20	(15 - 25)	48	(37 - 62)	80	(53 - 119)	128	(95 - 172)
Montana	26	(20 - 33)	3	(3 - 4)	8	(7 - 11)	14	(9 - 20)	22	(17 - 30)

(continued)

Table 6. *Illicit Drug Use Other Than Marijuana in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)*

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	44	(33 - 57)	6	(4 - 8)	14	(11 - 18)	24	(16 - 35)	38	(28 - 51)
Nevada	101	(79 - 129)	11	(8 - 14)	27	(21 - 33)	64	(45 - 90)	90	(69 - 118)
New Hampshire	52	(42 - 66)	5	(3 - 6)	18	(14 - 21)	30	(21 - 43)	48	(37 - 61)
New Jersey	241	(185 - 314)	25	(19 - 33)	72	(56 - 92)	145	(98 - 211)	217	(161 - 291)
New Mexico	65	(51 - 83)	9	(7 - 11)	20	(16 - 25)	37	(25 - 53)	57	(43 - 75)
New York	606	(522 - 704)	61	(51 - 72)	189	(164 - 217)	357	(284 - 448)	546	(462 - 644)
North Carolina	262	(205 - 334)	30	(23 - 39)	76	(59 - 97)	155	(109 - 222)	232	(175 - 305)
North Dakota	13	(10 - 17)	2	(2 - 3)	5	(3 - 6)	6	(4 - 9)	11	(8 - 15)
Ohio	333	(291 - 381)	43	(37 - 50)	106	(94 - 119)	184	(148 - 230)	290	(248 - 339)
Oklahoma	136	(107 - 173)	13	(10 - 17)	44	(34 - 56)	79	(56 - 111)	123	(94 - 160)
Oregon	140	(111 - 178)	16	(12 - 20)	40	(31 - 50)	85	(60 - 120)	125	(95 - 163)
Pennsylvania	340	(292 - 397)	39	(33 - 46)	109	(94 - 126)	192	(151 - 245)	301	(253 - 358)
Rhode Island	42	(33 - 54)	4	(3 - 5)	15	(11 - 18)	24	(16 - 34)	38	(29 - 50)
South Carolina	146	(109 - 194)	14	(11 - 18)	33	(25 - 45)	98	(67 - 144)	132	(96 - 181)
South Dakota	16	(12 - 21)	3	(2 - 4)	5	(4 - 7)	8	(5 - 13)	13	(9 - 19)
Tennessee	193	(150 - 249)	22	(17 - 28)	53	(41 - 67)	119	(83 - 169)	171	(128 - 228)
Texas	642	(557 - 740)	99	(84 - 117)	193	(166 - 224)	350	(278 - 439)	543	(460 - 641)
Utah	79	(63 - 99)	10	(8 - 13)	30	(24 - 38)	39	(27 - 56)	69	(53 - 90)
Vermont	25	(20 - 31)	2	(1 - 2)	9	(8 - 11)	13	(10 - 19)	23	(18 - 29)
Virginia	228	(177 - 293)	26	(20 - 33)	70	(55 - 88)	132	(91 - 191)	202	(152 - 268)
Washington	245	(193 - 311)	23	(17 - 29)	73	(58 - 91)	150	(107 - 210)	223	(171 - 289)
West Virginia	64	(51 - 82)	6	(5 - 8)	19	(16 - 24)	39	(27 - 55)	58	(45 - 76)
Wisconsin	175	(138 - 222)	21	(17 - 27)	61	(48 - 78)	92	(64 - 134)	154	(117 - 201)
Wyoming	15	(11 - 19)	2	(2 - 3)	5	(4 - 6)	8	(5 - 12)	13	(9 - 17)

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 7. Cocaine Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	4,670	(4,390 - 4,967)	241	(212 - 274)	1,692	(1,600 - 1,789)	2,738	(2,492 - 3,005)	4,430	(4,153 - 4,722)
Northeast	1,048	(937 - 1,172)	41	(34 - 49)	386	(348 - 427)	622	(526 - 734)	1,007	(897 - 1,131)
Midwest	853	(768 - 945)	50	(42 - 58)	317	(287 - 350)	486	(411 - 574)	803	(719 - 896)
South	1,539	(1,396 - 1,697)	80	(69 - 93)	519	(470 - 573)	941	(813 - 1,087)	1,459	(1,317 - 1,616)
West	1,230	(1,094 - 1,381)	70	(57 - 85)	471	(421 - 525)	689	(575 - 826)	1,160	(1,026 - 1,310)
Alabama	40	(26 - 63)	3	(2 - 4)	13	(9 - 21)	24	(13 - 44)	38	(23 - 61)
Alaska	13	(9 - 18)	1	(0 - 1)	5	(4 - 7)	8	(5 - 12)	12	(9 - 18)
Arizona	105	(73 - 150)	6	(4 - 9)	39	(29 - 54)	59	(36 - 97)	99	(67 - 144)
Arkansas	27	(18 - 40)	2	(1 - 3)	9	(6 - 13)	16	(10 - 28)	25	(16 - 39)
California	636	(529 - 763)	40	(30 - 53)	244	(204 - 291)	352	(267 - 466)	597	(491 - 724)
Colorado	126	(95 - 168)	4	(3 - 6)	46	(36 - 60)	75	(50 - 112)	122	(90 - 164)
Connecticut	50	(35 - 71)	3	(2 - 4)	19	(14 - 27)	28	(16 - 46)	47	(32 - 68)
Delaware	13	(9 - 19)	1	(0 - 1)	5	(3 - 6)	8	(5 - 13)	13	(9 - 18)
District of Columbia	16	(11 - 23)	0	(0 - 0)	4	(3 - 5)	12	(8 - 19)	16	(11 - 23)
Florida	287	(235 - 352)	12	(9 - 15)	93	(78 - 112)	182	(137 - 242)	275	(223 - 341)
Georgia	111	(75 - 164)	6	(4 - 8)	36	(25 - 52)	69	(41 - 115)	105	(69 - 159)
Hawaii	17	(11 - 24)	1	(1 - 1)	6	(4 - 8)	10	(6 - 17)	16	(11 - 24)
Idaho	13	(9 - 19)	1	(1 - 2)	6	(4 - 9)	6	(3 - 11)	12	(8 - 18)
Illinois	183	(151 - 222)	9	(7 - 12)	65	(55 - 78)	109	(82 - 144)	174	(142 - 213)
Indiana	61	(42 - 90)	5	(3 - 7)	24	(16 - 34)	33	(19 - 57)	57	(37 - 86)
Iowa	26	(17 - 38)	3	(2 - 4)	11	(7 - 16)	12	(7 - 23)	23	(15 - 36)
Kansas	36	(26 - 51)	2	(2 - 3)	15	(11 - 21)	19	(11 - 31)	34	(23 - 49)
Kentucky	45	(30 - 66)	4	(2 - 5)	17	(11 - 24)	25	(14 - 43)	41	(27 - 63)
Louisiana	60	(42 - 87)	3	(2 - 4)	19	(14 - 27)	38	(24 - 61)	58	(39 - 84)
Maine	22	(17 - 30)	1	(1 - 1)	9	(7 - 12)	13	(8 - 20)	21	(16 - 30)
Maryland	84	(57 - 125)	4	(2 - 5)	26	(18 - 37)	54	(33 - 89)	81	(53 - 121)
Massachusetts	166	(124 - 221)	5	(3 - 7)	62	(48 - 80)	99	(66 - 146)	161	(119 - 216)
Michigan	138	(113 - 168)	7	(5 - 9)	50	(42 - 60)	81	(61 - 109)	131	(107 - 161)
Minnesota	55	(37 - 80)	4	(3 - 6)	21	(15 - 30)	30	(17 - 51)	51	(34 - 76)
Mississippi	38	(26 - 56)	2	(1 - 2)	11	(8 - 16)	25	(16 - 41)	37	(25 - 55)
Missouri	84	(59 - 118)	4	(3 - 6)	29	(21 - 41)	50	(31 - 79)	79	(55 - 114)
Montana	15	(11 - 21)	1	(1 - 1)	6	(5 - 9)	8	(5 - 13)	14	(10 - 20)

(continued)

Table 7. Cocaine Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	20	(14 - 29)	1	(1 - 2)	8	(6 - 12)	11	(6 - 18)	19	(13 - 28)
Nevada	52	(37 - 72)	2	(1 - 3)	17	(13 - 22)	33	(21 - 50)	49	(35 - 69)
New Hampshire	31	(23 - 41)	1	(1 - 2)	12	(9 - 15)	18	(12 - 26)	30	(22 - 40)
New Jersey	160	(115 - 224)	6	(4 - 10)	52	(38 - 71)	102	(66 - 157)	154	(109 - 218)
New Mexico	40	(29 - 53)	2	(1 - 3)	15	(12 - 20)	22	(14 - 34)	37	(27 - 51)
New York	369	(307 - 444)	15	(11 - 19)	144	(122 - 169)	211	(160 - 279)	355	(293 - 430)
North Carolina	139	(97 - 198)	6	(4 - 9)	43	(31 - 59)	90	(57 - 142)	133	(91 - 193)
North Dakota	5	(4 - 8)	0	(0 - 1)	2	(2 - 4)	2	(1 - 4)	5	(3 - 7)
Ohio	142	(115 - 175)	9	(7 - 12)	50	(42 - 60)	83	(60 - 114)	133	(106 - 167)
Oklahoma	41	(27 - 60)	3	(2 - 5)	17	(11 - 25)	21	(12 - 37)	38	(25 - 58)
Oregon	62	(46 - 85)	4	(2 - 5)	25	(19 - 34)	34	(22 - 53)	59	(42 - 82)
Pennsylvania	207	(170 - 253)	8	(6 - 11)	70	(58 - 85)	128	(97 - 170)	199	(161 - 245)
Rhode Island	26	(19 - 36)	1	(1 - 1)	10	(7 - 13)	15	(10 - 24)	25	(18 - 35)
South Carolina	79	(55 - 114)	2	(2 - 4)	20	(14 - 28)	57	(36 - 89)	77	(52 - 112)
South Dakota	8	(5 - 11)	1	(0 - 1)	3	(2 - 5)	4	(2 - 7)	7	(4 - 11)
Tennessee	64	(42 - 97)	4	(3 - 7)	19	(13 - 29)	40	(23 - 69)	59	(38 - 93)
Texas	347	(286 - 421)	24	(18 - 31)	132	(110 - 159)	191	(142 - 256)	324	(263 - 397)
Utah	28	(20 - 41)	3	(2 - 4)	13	(9 - 19)	13	(7 - 22)	26	(17 - 38)
Vermont	16	(13 - 21)	1	(0 - 1)	7	(6 - 9)	9	(6 - 13)	16	(12 - 21)
Virginia	119	(86 - 163)	5	(4 - 7)	43	(32 - 58)	70	(46 - 109)	114	(81 - 158)
Washington	117	(84 - 161)	5	(4 - 8)	45	(33 - 60)	67	(42 - 105)	111	(79 - 156)
West Virginia	29	(21 - 40)	1	(1 - 2)	11	(8 - 14)	17	(11 - 26)	27	(19 - 39)
Wisconsin	95	(68 - 132)	4	(3 - 6)	38	(28 - 51)	53	(33 - 84)	91	(64 - 128)
Wyoming	6	(4 - 9)	0	(0 - 1)	3	(2 - 4)	3	(2 - 5)	6	(4 - 9)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 8. Nonmedical Use of Pain Relievers in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	12,346	(11,887 - 12,822)	1,575	(1,500 - 1,653)	3,903	(3,763 - 4,048)	6,868	(6,460 - 7,301)	10,771	(10,321 - 11,240)
Northeast	2,091	(1,950 - 2,243)	231	(210 - 254)	730	(686 - 776)	1,131	(1,010 - 1,265)	1,860	(1,722 - 2,010)
Midwest	2,656	(2,513 - 2,804)	353	(331 - 377)	889	(844 - 937)	1,413	(1,289 - 1,547)	2,303	(2,163 - 2,449)
South	4,411	(4,167 - 4,668)	582	(545 - 622)	1,378	(1,309 - 1,451)	2,450	(2,238 - 2,682)	3,829	(3,591 - 4,082)
West	3,188	(2,951 - 3,443)	408	(373 - 447)	906	(844 - 971)	1,874	(1,663 - 2,110)	2,780	(2,549 - 3,031)
Alabama	180	(146 - 220)	27	(22 - 33)	57	(46 - 70)	95	(70 - 129)	152	(120 - 192)
Alaska	30	(24 - 37)	4	(3 - 5)	9	(8 - 11)	17	(12 - 23)	26	(21 - 33)
Arizona	338	(270 - 421)	41	(33 - 51)	89	(73 - 108)	208	(152 - 285)	297	(231 - 379)
Arkansas	131	(106 - 161)	17	(14 - 21)	37	(31 - 45)	76	(56 - 102)	113	(90 - 143)
California	1,493	(1,324 - 1,684)	205	(178 - 236)	419	(369 - 474)	870	(724 - 1,043)	1,288	(1,124 - 1,476)
Colorado	257	(209 - 316)	28	(22 - 34)	74	(62 - 89)	155	(115 - 208)	229	(182 - 288)
Connecticut	122	(97 - 152)	14	(11 - 18)	42	(34 - 51)	66	(47 - 92)	107	(84 - 137)
Delaware	41	(33 - 50)	4	(3 - 5)	13	(11 - 15)	24	(17 - 32)	37	(29 - 46)
District of Columbia	22	(17 - 28)	2	(1 - 2)	7	(6 - 9)	13	(10 - 18)	20	(16 - 26)
Florida	679	(593 - 776)	80	(69 - 93)	181	(159 - 205)	418	(344 - 507)	598	(515 - 696)
Georgia	337	(274 - 414)	49	(40 - 62)	110	(90 - 134)	178	(130 - 242)	288	(228 - 363)
Hawaii	44	(35 - 57)	5	(4 - 6)	12	(10 - 15)	27	(19 - 38)	39	(30 - 52)
Idaho	76	(62 - 92)	10	(8 - 12)	23	(19 - 27)	43	(32 - 58)	66	(53 - 82)
Illinois	418	(369 - 473)	58	(49 - 67)	147	(130 - 165)	214	(174 - 262)	360	(313 - 414)
Indiana	302	(250 - 363)	40	(32 - 49)	103	(86 - 123)	159	(119 - 211)	262	(213 - 322)
Iowa	93	(74 - 114)	15	(12 - 19)	32	(25 - 40)	46	(32 - 64)	77	(60 - 99)
Kansas	108	(89 - 131)	15	(12 - 19)	37	(30 - 45)	55	(41 - 74)	92	(74 - 115)
Kentucky	191	(157 - 231)	25	(20 - 31)	61	(51 - 73)	105	(79 - 140)	166	(133 - 205)
Louisiana	207	(171 - 251)	23	(19 - 29)	73	(61 - 88)	110	(82 - 147)	184	(149 - 226)
Maine	51	(41 - 63)	6	(5 - 7)	18	(15 - 21)	27	(20 - 38)	45	(36 - 57)
Maryland	200	(157 - 253)	26	(21 - 33)	63	(50 - 78)	111	(78 - 156)	173	(133 - 226)
Massachusetts	283	(233 - 343)	28	(22 - 35)	101	(83 - 121)	155	(116 - 206)	255	(207 - 315)
Michigan	460	(410 - 515)	53	(46 - 60)	147	(133 - 162)	260	(216 - 312)	407	(358 - 462)
Minnesota	179	(144 - 222)	24	(19 - 30)	63	(51 - 77)	92	(66 - 129)	155	(122 - 198)
Mississippi	121	(98 - 148)	21	(17 - 26)	37	(30 - 45)	63	(46 - 86)	100	(78 - 126)
Missouri	254	(209 - 307)	32	(26 - 40)	85	(70 - 103)	136	(102 - 182)	221	(178 - 274)
Montana	41	(34 - 50)	5	(4 - 6)	14	(11 - 16)	23	(17 - 31)	36	(29 - 45)

(continued)

Table 8. Nonmedical Use of Pain Relievers in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	57	(46 - 71)	8	(6 - 10)	20	(16 - 25)	29	(21 - 40)	49	(39 - 62)
Nevada	128	(104 - 157)	16	(13 - 21)	34	(28 - 41)	78	(58 - 104)	112	(89 - 140)
New Hampshire	61	(49 - 74)	6	(5 - 8)	21	(18 - 25)	33	(25 - 45)	54	(43 - 67)
New Jersey	301	(243 - 372)	34	(27 - 44)	105	(86 - 127)	162	(117 - 223)	266	(211 - 336)
New Mexico	95	(76 - 117)	13	(11 - 17)	26	(21 - 31)	56	(41 - 76)	81	(64 - 103)
New York	729	(646 - 823)	79	(68 - 93)	258	(231 - 289)	391	(322 - 474)	649	(568 - 742)
North Carolina	347	(281 - 428)	50	(40 - 61)	104	(85 - 127)	193	(142 - 261)	297	(234 - 376)
North Dakota	22	(18 - 27)	3	(3 - 4)	8	(7 - 10)	11	(8 - 15)	19	(15 - 24)
Ohio	525	(472 - 583)	70	(62 - 80)	165	(150 - 181)	289	(243 - 344)	454	(403 - 512)
Oklahoma	209	(170 - 256)	23	(19 - 29)	66	(54 - 79)	120	(89 - 161)	186	(149 - 232)
Oregon	215	(176 - 262)	23	(18 - 28)	59	(49 - 70)	133	(100 - 176)	192	(154 - 239)
Pennsylvania	466	(410 - 530)	55	(48 - 64)	157	(140 - 176)	254	(207 - 311)	411	(357 - 474)
Rhode Island	53	(43 - 65)	5	(4 - 6)	19	(16 - 23)	29	(21 - 39)	48	(38 - 60)
South Carolina	189	(153 - 234)	21	(17 - 26)	59	(48 - 72)	109	(80 - 148)	168	(133 - 213)
South Dakota	24	(19 - 30)	4	(3 - 5)	8	(6 - 10)	12	(9 - 18)	20	(16 - 26)
Tennessee	232	(188 - 285)	30	(25 - 38)	77	(63 - 94)	125	(91 - 171)	201	(159 - 254)
Texas	909	(805 - 1,025)	130	(112 - 150)	298	(265 - 336)	481	(395 - 584)	779	(679 - 893)
Utah	106	(88 - 129)	17	(13 - 21)	39	(32 - 48)	51	(37 - 68)	90	(72 - 112)
Vermont	26	(21 - 32)	3	(2 - 3)	10	(8 - 12)	14	(10 - 19)	23	(19 - 29)
Virginia	330	(271 - 401)	42	(34 - 52)	108	(90 - 130)	180	(134 - 241)	288	(231 - 358)
Washington	344	(283 - 417)	39	(31 - 47)	102	(86 - 121)	203	(153 - 269)	305	(247 - 377)
West Virginia	86	(71 - 105)	9	(8 - 12)	26	(22 - 31)	50	(38 - 68)	77	(62 - 96)
Wisconsin	215	(173 - 266)	31	(25 - 39)	74	(60 - 91)	110	(78 - 153)	184	(143 - 234)
Wyoming	20	(16 - 25)	3	(2 - 4)	7	(5 - 8)	11	(8 - 15)	17	(14 - 22)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 9. Alcohol Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	131,017	(129,609 - 132,407)	3,484	(3,374 - 3,597)	20,834	(20,580 - 21,087)	106,699	(105,352 - 108,024)	127,533	(126,145 - 128,899)
Northeast	26,537	(25,919 - 27,151)	690	(654 - 729)	4,114	(4,003 - 4,222)	21,733	(21,178 - 22,283)	25,847	(25,228 - 26,461)
Midwest	30,202	(29,551 - 30,855)	763	(730 - 797)	4,818	(4,694 - 4,939)	24,622	(24,050 - 25,191)	29,439	(28,786 - 30,089)
South	44,208	(43,120 - 45,282)	1,204	(1,150 - 1,261)	7,056	(6,852 - 7,258)	35,949	(35,001 - 36,880)	43,004	(41,916 - 44,075)
West	30,069	(29,260 - 30,872)	827	(776 - 879)	4,847	(4,692 - 4,999)	24,396	(23,670 - 25,111)	29,243	(28,433 - 30,040)
Alabama	1,674	(1,550 - 1,800)	46	(38 - 54)	270	(248 - 292)	1,359	(1,244 - 1,475)	1,628	(1,505 - 1,753)
Alaska	293	(274 - 312)	8	(7 - 10)	48	(45 - 52)	237	(219 - 254)	285	(266 - 304)
Arizona	2,573	(2,378 - 2,772)	73	(61 - 88)	389	(360 - 417)	2,112	(1,928 - 2,294)	2,500	(2,306 - 2,693)
Arkansas	976	(898 - 1,056)	33	(28 - 39)	165	(153 - 177)	778	(705 - 853)	943	(866 - 1,022)
California	15,204	(14,627 - 15,780)	434	(394 - 478)	2,639	(2,529 - 2,745)	12,132	(11,594 - 12,666)	14,770	(14,193 - 15,342)
Colorado	2,585	(2,446 - 2,719)	67	(57 - 79)	383	(361 - 402)	2,135	(2,004 - 2,258)	2,518	(2,379 - 2,649)
Connecticut	1,747	(1,645 - 1,848)	50	(43 - 59)	258	(243 - 272)	1,439	(1,342 - 1,532)	1,697	(1,594 - 1,795)
Delaware	416	(390 - 441)	11	(9 - 13)	63	(59 - 66)	342	(317 - 366)	405	(379 - 430)
District of Columbia	317	(300 - 333)	5	(4 - 6)	62	(59 - 65)	249	(233 - 265)	312	(295 - 328)
Florida	7,884	(7,580 - 8,187)	192	(174 - 211)	1,051	(1,003 - 1,098)	6,642	(6,355 - 6,926)	7,692	(7,390 - 7,995)
Georgia	3,793	(3,522 - 4,065)	116	(99 - 137)	609	(566 - 651)	3,068	(2,816 - 3,319)	3,676	(3,408 - 3,943)
Hawaii	503	(466 - 540)	12	(10 - 14)	78	(72 - 84)	412	(379 - 446)	491	(455 - 527)
Idaho	590	(549 - 631)	17	(15 - 20)	94	(87 - 100)	479	(441 - 517)	573	(532 - 613)
Illinois	5,753	(5,559 - 5,944)	145	(131 - 160)	974	(938 - 1,008)	4,634	(4,455 - 4,811)	5,608	(5,416 - 5,797)
Indiana	2,606	(2,433 - 2,780)	67	(56 - 79)	422	(394 - 449)	2,117	(1,956 - 2,277)	2,539	(2,367 - 2,710)
Iowa	1,403	(1,319 - 1,486)	41	(35 - 47)	239	(225 - 252)	1,123	(1,044 - 1,199)	1,362	(1,278 - 1,442)
Kansas	1,271	(1,196 - 1,345)	36	(30 - 41)	209	(196 - 222)	1,026	(956 - 1,094)	1,235	(1,160 - 1,308)
Kentucky	1,391	(1,278 - 1,508)	44	(36 - 52)	221	(203 - 240)	1,126	(1,021 - 1,234)	1,347	(1,235 - 1,462)
Louisiana	1,710	(1,588 - 1,833)	55	(47 - 65)	313	(292 - 334)	1,341	(1,229 - 1,454)	1,654	(1,533 - 1,775)
Maine	634	(597 - 670)	13	(11 - 16)	83	(78 - 88)	537	(502 - 572)	621	(584 - 657)
Maryland	2,450	(2,277 - 2,622)	58	(49 - 69)	364	(336 - 391)	2,028	(1,867 - 2,186)	2,392	(2,220 - 2,561)
Massachusetts	3,554	(3,383 - 3,717)	77	(65 - 90)	569	(541 - 595)	2,907	(2,747 - 3,058)	3,477	(3,307 - 3,637)
Michigan	4,553	(4,397 - 4,707)	110	(100 - 121)	687	(661 - 713)	3,756	(3,610 - 3,899)	4,443	(4,287 - 4,597)
Minnesota	2,571	(2,426 - 2,715)	54	(46 - 64)	389	(366 - 411)	2,128	(1,990 - 2,259)	2,517	(2,370 - 2,656)
Mississippi	973	(897 - 1,050)	31	(26 - 37)	178	(165 - 191)	764	(694 - 835)	942	(867 - 1,017)
Missouri	2,505	(2,340 - 2,669)	72	(62 - 84)	389	(363 - 414)	2,043	(1,888 - 2,197)	2,433	(2,269 - 2,595)
Montana	474	(447 - 502)	14	(12 - 16)	73	(68 - 77)	387	(361 - 413)	460	(433 - 487)

(continued)

Table 9. Alcohol Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	759	(709 - 808)	19	(16 - 23)	136	(128 - 144)	603	(556 - 650)	739	(690 - 788)
Nevada	1,173	(1,095 - 1,250)	31	(26 - 37)	150	(140 - 161)	992	(918 - 1,064)	1,143	(1,065 - 1,218)
New Hampshire	719	(682 - 754)	18	(16 - 21)	104	(99 - 108)	597	(562 - 629)	700	(664 - 735)
New Jersey	4,041	(3,799 - 4,279)	106	(90 - 124)	565	(529 - 598)	3,370	(3,141 - 3,593)	3,935	(3,694 - 4,169)
New Mexico	748	(691 - 806)	23	(20 - 28)	129	(119 - 138)	596	(543 - 649)	725	(668 - 781)
New York	9,113	(8,791 - 9,431)	253	(230 - 277)	1,490	(1,435 - 1,542)	7,371	(7,070 - 7,667)	8,861	(8,539 - 9,177)
North Carolina	3,745	(3,489 - 4,001)	89	(74 - 106)	584	(545 - 622)	3,072	(2,829 - 3,314)	3,656	(3,402 - 3,908)
North Dakota	322	(304 - 339)	8	(7 - 9)	65	(62 - 68)	249	(232 - 265)	314	(296 - 331)
Ohio	5,052	(4,868 - 5,235)	127	(116 - 140)	794	(766 - 821)	4,131	(3,957 - 4,303)	4,925	(4,742 - 5,106)
Oklahoma	1,407	(1,300 - 1,515)	43	(36 - 51)	255	(237 - 272)	1,109	(1,009 - 1,210)	1,364	(1,258 - 1,470)
Oregon	1,961	(1,857 - 2,063)	50	(43 - 57)	269	(254 - 284)	1,642	(1,543 - 1,736)	1,912	(1,807 - 2,011)
Pennsylvania	5,852	(5,628 - 6,073)	152	(138 - 167)	902	(867 - 936)	4,798	(4,589 - 5,004)	5,700	(5,477 - 5,920)
Rhode Island	545	(512 - 577)	13	(11 - 15)	91	(85 - 96)	442	(411 - 471)	532	(500 - 563)
South Carolina	1,723	(1,596 - 1,851)	43	(36 - 51)	266	(246 - 285)	1,415	(1,296 - 1,534)	1,680	(1,554 - 1,806)
South Dakota	383	(361 - 405)	10	(8 - 12)	65	(62 - 69)	308	(287 - 328)	373	(351 - 395)
Tennessee	2,107	(1,927 - 2,292)	55	(47 - 65)	332	(305 - 359)	1,719	(1,550 - 1,893)	2,051	(1,872 - 2,234)
Texas	9,674	(9,299 - 10,048)	284	(256 - 314)	1,666	(1,599 - 1,733)	7,724	(7,378 - 8,067)	9,390	(9,018 - 9,759)
Utah	572	(510 - 637)	19	(15 - 25)	123	(109 - 139)	429	(373 - 490)	552	(492 - 616)
Vermont	331	(314 - 348)	8	(7 - 9)	53	(50 - 55)	271	(254 - 287)	323	(306 - 340)
Virginia	3,349	(3,132 - 3,564)	81	(68 - 95)	558	(524 - 591)	2,710	(2,506 - 2,912)	3,268	(3,053 - 3,481)
Washington	3,156	(2,971 - 3,337)	70	(59 - 83)	435	(407 - 463)	2,650	(2,477 - 2,817)	3,085	(2,901 - 3,264)
West Virginia	621	(568 - 675)	19	(16 - 22)	99	(92 - 106)	503	(453 - 554)	602	(550 - 655)
Wisconsin	3,025	(2,878 - 3,165)	74	(64 - 85)	447	(422 - 471)	2,503	(2,365 - 2,633)	2,951	(2,804 - 3,089)
Wyoming	236	(221 - 252)	7	(6 - 8)	37	(35 - 40)	192	(178 - 206)	230	(214 - 245)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 10. Binge Alcohol Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	59,113	(58,039 - 60,199)	2,051	(1,965 - 2,139)	13,920	(13,664 - 14,178)	43,142	(42,137 - 44,162)	57,063	(55,990 - 58,147)
Northeast	11,537	(11,147 - 11,936)	402	(373 - 433)	2,751	(2,673 - 2,830)	8,384	(8,014 - 8,766)	11,135	(10,749 - 11,531)
Midwest	14,112	(13,751 - 14,480)	468	(441 - 496)	3,329	(3,251 - 3,407)	10,316	(9,972 - 10,668)	13,644	(13,284 - 14,009)
South	19,965	(19,394 - 20,546)	677	(638 - 719)	4,602	(4,480 - 4,725)	14,686	(14,135 - 15,250)	19,288	(18,719 - 19,867)
West	13,499	(13,030 - 13,979)	504	(463 - 548)	3,238	(3,132 - 3,346)	9,757	(9,313 - 10,215)	12,995	(12,531 - 13,471)
Alabama	809	(716 - 911)	26	(21 - 32)	169	(150 - 190)	614	(528 - 710)	783	(691 - 884)
Alaska	137	(123 - 152)	5	(4 - 7)	33	(30 - 37)	98	(86 - 112)	132	(118 - 146)
Arizona	1,264	(1,122 - 1,417)	47	(38 - 58)	271	(244 - 299)	946	(816 - 1,090)	1,217	(1,076 - 1,367)
Arkansas	448	(397 - 505)	18	(15 - 22)	104	(92 - 116)	327	(280 - 379)	430	(379 - 486)
California	6,896	(6,509 - 7,299)	266	(232 - 304)	1,726	(1,636 - 1,817)	4,905	(4,541 - 5,289)	6,630	(6,247 - 7,030)
Colorado	1,081	(973 - 1,197)	37	(30 - 45)	255	(234 - 276)	789	(690 - 898)	1,044	(937 - 1,159)
Connecticut	776	(699 - 858)	32	(26 - 39)	183	(167 - 199)	561	(490 - 639)	744	(667 - 825)
Delaware	178	(160 - 199)	6	(5 - 8)	43	(39 - 46)	129	(112 - 149)	172	(154 - 192)
District of Columbia	152	(138 - 166)	2	(2 - 3)	40	(37 - 44)	109	(96 - 123)	149	(136 - 164)
Florida	3,215	(3,015 - 3,425)	102	(89 - 117)	659	(621 - 699)	2,453	(2,261 - 2,658)	3,113	(2,915 - 3,322)
Georgia	1,590	(1,416 - 1,780)	59	(48 - 73)	390	(352 - 430)	1,141	(980 - 1,321)	1,530	(1,358 - 1,718)
Hawaii	249	(223 - 277)	8	(6 - 10)	54	(48 - 59)	188	(164 - 214)	241	(216 - 269)
Idaho	272	(244 - 303)	11	(9 - 13)	64	(58 - 70)	198	(171 - 226)	261	(233 - 292)
Illinois	2,760	(2,619 - 2,905)	92	(81 - 104)	681	(650 - 712)	1,987	(1,852 - 2,127)	2,668	(2,528 - 2,812)
Indiana	1,189	(1,068 - 1,320)	41	(34 - 50)	286	(259 - 313)	862	(752 - 984)	1,148	(1,028 - 1,277)
Iowa	686	(622 - 752)	25	(21 - 30)	171	(158 - 185)	489	(432 - 552)	661	(598 - 725)
Kansas	590	(532 - 652)	21	(17 - 26)	142	(128 - 156)	427	(374 - 485)	569	(511 - 631)
Kentucky	770	(690 - 856)	27	(22 - 34)	174	(158 - 192)	568	(495 - 650)	743	(663 - 828)
Louisiana	840	(753 - 934)	27	(22 - 33)	201	(183 - 221)	612	(532 - 700)	813	(726 - 906)
Maine	259	(233 - 288)	8	(7 - 10)	59	(54 - 64)	192	(167 - 220)	251	(225 - 280)
Maryland	952	(838 - 1,077)	35	(28 - 43)	223	(198 - 249)	695	(590 - 812)	917	(804 - 1,042)
Massachusetts	1,428	(1,294 - 1,570)	46	(38 - 56)	372	(343 - 403)	1,009	(886 - 1,144)	1,382	(1,249 - 1,523)
Michigan	2,197	(2,079 - 2,319)	66	(58 - 74)	474	(452 - 496)	1,657	(1,545 - 1,775)	2,131	(2,014 - 2,252)
Minnesota	1,130	(1,021 - 1,246)	33	(27 - 40)	266	(244 - 288)	831	(731 - 941)	1,097	(988 - 1,212)
Mississippi	477	(425 - 534)	17	(13 - 21)	111	(99 - 123)	349	(302 - 402)	460	(408 - 517)
Missouri	1,183	(1,065 - 1,310)	45	(37 - 55)	255	(232 - 278)	883	(772 - 1,004)	1,138	(1,021 - 1,263)
Montana	229	(208 - 251)	9	(8 - 11)	55	(51 - 59)	164	(145 - 185)	220	(199 - 241)

(continued)

Table 10. Binge Alcohol Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	362	(326 - 401)	13	(10 - 15)	99	(91 - 107)	251	(218 - 287)	350	(314 - 388)
Nevada	573	(512 - 638)	18	(14 - 22)	108	(98 - 118)	447	(390 - 509)	555	(495 - 619)
New Hampshire	271	(243 - 301)	11	(9 - 13)	68	(62 - 73)	192	(166 - 221)	260	(232 - 290)
New Jersey	1,841	(1,656 - 2,040)	61	(49 - 75)	401	(366 - 436)	1,379	(1,206 - 1,569)	1,780	(1,596 - 1,977)
New Mexico	365	(326 - 407)	15	(12 - 18)	86	(77 - 94)	265	(229 - 304)	350	(312 - 392)
New York	3,936	(3,696 - 4,187)	145	(128 - 164)	944	(898 - 991)	2,847	(2,620 - 3,088)	3,791	(3,553 - 4,040)
North Carolina	1,579	(1,405 - 1,770)	50	(40 - 61)	368	(331 - 405)	1,162	(1,002 - 1,340)	1,530	(1,357 - 1,718)
North Dakota	160	(147 - 174)	5	(4 - 6)	48	(44 - 51)	108	(96 - 121)	155	(142 - 169)
Ohio	2,261	(2,137 - 2,391)	77	(68 - 86)	544	(521 - 567)	1,641	(1,521 - 1,767)	2,184	(2,061 - 2,314)
Oklahoma	646	(575 - 723)	26	(21 - 32)	164	(147 - 182)	456	(393 - 527)	620	(550 - 697)
Oregon	719	(649 - 797)	25	(20 - 30)	171	(155 - 187)	524	(457 - 597)	695	(625 - 771)
Pennsylvania	2,648	(2,497 - 2,805)	87	(77 - 99)	628	(597 - 658)	1,933	(1,791 - 2,082)	2,561	(2,411 - 2,717)
Rhode Island	236	(211 - 262)	7	(6 - 9)	59	(54 - 65)	169	(147 - 194)	229	(204 - 254)
South Carolina	831	(737 - 935)	22	(17 - 27)	173	(155 - 193)	636	(549 - 734)	810	(716 - 912)
South Dakota	184	(167 - 201)	6	(5 - 7)	46	(43 - 50)	131	(116 - 148)	178	(161 - 195)
Tennessee	917	(798 - 1,049)	33	(26 - 40)	206	(183 - 230)	678	(567 - 806)	884	(766 - 1,015)
Texas	4,720	(4,463 - 4,986)	166	(146 - 188)	1,113	(1,057 - 1,169)	3,441	(3,196 - 3,700)	4,554	(4,299 - 4,819)
Utah	286	(245 - 333)	14	(11 - 19)	89	(77 - 103)	183	(147 - 226)	272	(232 - 318)
Vermont	142	(128 - 156)	5	(4 - 6)	37	(34 - 40)	100	(88 - 114)	137	(124 - 152)
Virginia	1,525	(1,371 - 1,692)	50	(41 - 61)	393	(359 - 428)	1,082	(941 - 1,238)	1,475	(1,323 - 1,640)
Washington	1,325	(1,192 - 1,468)	45	(37 - 55)	302	(275 - 329)	978	(854 - 1,115)	1,280	(1,148 - 1,422)
West Virginia	314	(279 - 354)	11	(9 - 14)	71	(64 - 78)	233	(199 - 270)	303	(268 - 342)
Wisconsin	1,410	(1,283 - 1,544)	44	(37 - 53)	318	(292 - 343)	1,048	(930 - 1,175)	1,366	(1,240 - 1,498)
Wyoming	102	(92 - 113)	4	(3 - 6)	26	(23 - 28)	72	(63 - 82)	98	(88 - 109)

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 11. Perceptions of Great Risk of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	107,623	(106,311 - 108,933)	9,796	(9,647 - 9,946)	11,312	(11,089 - 11,537)	86,515	(85,246 - 87,781)	97,828	(96,524 - 99,128)
Northeast	18,825	(18,356 - 19,298)	1,651	(1,600 - 1,702)	1,875	(1,806 - 1,944)	15,300	(14,849 - 15,754)	17,174	(16,715 - 17,638)
Midwest	21,534	(21,092 - 21,976)	2,045	(1,997 - 2,094)	2,189	(2,127 - 2,252)	17,299	(16,874 - 17,724)	19,488	(19,055 - 19,922)
South	41,596	(40,857 - 42,337)	3,755	(3,674 - 3,837)	4,444	(4,339 - 4,551)	33,397	(32,680 - 34,114)	37,841	(37,114 - 38,569)
West	25,669	(25,049 - 26,289)	2,344	(2,271 - 2,419)	2,804	(2,706 - 2,904)	20,520	(19,920 - 21,121)	23,324	(22,712 - 23,937)
Alabama	1,772	(1,652 - 1,894)	167	(154 - 180)	203	(185 - 221)	1,403	(1,288 - 1,519)	1,605	(1,488 - 1,724)
Alaska	236	(218 - 254)	25	(23 - 27)	25	(22 - 29)	185	(169 - 202)	211	(193 - 228)
Arizona	2,350	(2,177 - 2,527)	227	(207 - 247)	257	(231 - 283)	1,866	(1,702 - 2,033)	2,123	(1,955 - 2,295)
Arkansas	1,076	(1,003 - 1,151)	101	(93 - 110)	108	(98 - 119)	867	(797 - 938)	975	(904 - 1,047)
California	13,684	(13,180 - 14,189)	1,254	(1,193 - 1,316)	1,581	(1,495 - 1,669)	10,849	(10,362 - 11,337)	12,430	(11,935 - 12,927)
Colorado	1,610	(1,475 - 1,749)	137	(124 - 151)	158	(141 - 177)	1,315	(1,187 - 1,447)	1,473	(1,341 - 1,608)
Connecticut	1,247	(1,152 - 1,346)	114	(104 - 125)	116	(103 - 129)	1,017	(926 - 1,111)	1,133	(1,039 - 1,230)
Delaware	299	(276 - 323)	26	(24 - 29)	30	(27 - 34)	242	(221 - 265)	273	(250 - 296)
District of Columbia	213	(197 - 229)	16	(15 - 17)	28	(25 - 32)	169	(154 - 184)	197	(182 - 213)
Florida	7,347	(7,076 - 7,618)	568	(542 - 595)	701	(665 - 737)	6,078	(5,815 - 6,342)	6,779	(6,512 - 7,046)
Georgia	3,622	(3,375 - 3,872)	358	(329 - 386)	379	(345 - 415)	2,885	(2,651 - 3,121)	3,264	(3,024 - 3,508)
Hawaii	432	(397 - 467)	40	(36 - 43)	46	(41 - 52)	345	(312 - 380)	392	(358 - 427)
Idaho	480	(443 - 519)	49	(45 - 53)	53	(47 - 58)	379	(344 - 415)	432	(395 - 469)
Illinois	4,400	(4,219 - 4,584)	424	(403 - 446)	458	(432 - 485)	3,518	(3,343 - 3,694)	3,976	(3,798 - 4,156)
Indiana	2,074	(1,918 - 2,235)	206	(188 - 224)	211	(189 - 235)	1,657	(1,510 - 1,808)	1,868	(1,716 - 2,025)
Iowa	921	(846 - 1,000)	87	(79 - 95)	95	(85 - 106)	739	(668 - 813)	834	(761 - 911)
Kansas	884	(816 - 953)	82	(74 - 89)	94	(84 - 105)	708	(644 - 773)	802	(736 - 870)
Kentucky	1,483	(1,377 - 1,592)	137	(125 - 149)	153	(138 - 169)	1,193	(1,093 - 1,296)	1,347	(1,243 - 1,452)
Louisiana	1,660	(1,546 - 1,775)	162	(150 - 175)	209	(191 - 227)	1,289	(1,182 - 1,397)	1,498	(1,387 - 1,610)
Maine	425	(390 - 461)	32	(29 - 35)	35	(31 - 39)	358	(324 - 392)	393	(359 - 428)
Maryland	2,066	(1,911 - 2,223)	191	(175 - 208)	222	(199 - 245)	1,653	(1,506 - 1,802)	1,874	(1,723 - 2,028)
Massachusetts	2,100	(1,934 - 2,270)	176	(160 - 193)	212	(189 - 238)	1,711	(1,556 - 1,872)	1,924	(1,763 - 2,090)
Michigan	3,283	(3,148 - 3,420)	320	(304 - 336)	362	(342 - 382)	2,602	(2,471 - 2,735)	2,963	(2,831 - 3,097)
Minnesota	1,617	(1,486 - 1,753)	150	(136 - 164)	154	(137 - 172)	1,313	(1,187 - 1,443)	1,467	(1,338 - 1,601)
Mississippi	1,143	(1,071 - 1,216)	118	(109 - 127)	139	(128 - 151)	886	(817 - 954)	1,025	(955 - 1,095)
Missouri	1,955	(1,809 - 2,104)	182	(167 - 199)	203	(182 - 224)	1,570	(1,432 - 1,711)	1,772	(1,631 - 1,918)
Montana	299	(275 - 324)	24	(22 - 27)	29	(26 - 33)	245	(222 - 269)	275	(251 - 299)

(continued)

Table 11. Perceptions of Great Risk of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	546	(502 - 592)	53	(48 - 58)	55	(49 - 62)	439	(397 - 482)	494	(451 - 538)
Nevada	917	(847 - 989)	84	(77 - 92)	86	(77 - 96)	747	(680 - 815)	833	(765 - 903)
New Hampshire	431	(397 - 466)	36	(32 - 39)	35	(31 - 39)	360	(328 - 393)	395	(362 - 429)
New Jersey	3,011	(2,781 - 3,246)	257	(233 - 282)	287	(257 - 320)	2,467	(2,249 - 2,689)	2,754	(2,531 - 2,982)
New Mexico	782	(730 - 834)	64	(59 - 70)	84	(76 - 92)	634	(585 - 683)	717	(667 - 768)
New York	6,918	(6,628 - 7,212)	625	(595 - 656)	728	(686 - 772)	5,565	(5,284 - 5,849)	6,293	(6,006 - 6,583)
North Carolina	3,361	(3,117 - 3,608)	275	(251 - 301)	339	(307 - 371)	2,747	(2,514 - 2,982)	3,085	(2,847 - 3,326)
North Dakota	192	(177 - 208)	16	(15 - 18)	23	(20 - 26)	153	(139 - 167)	176	(161 - 191)
Ohio	3,785	(3,628 - 3,945)	353	(337 - 369)	338	(319 - 359)	3,094	(2,942 - 3,248)	3,432	(3,277 - 3,590)
Oklahoma	1,253	(1,158 - 1,349)	114	(103 - 125)	142	(127 - 157)	997	(908 - 1,088)	1,139	(1,047 - 1,233)
Oregon	1,284	(1,192 - 1,378)	107	(98 - 117)	113	(100 - 126)	1,064	(977 - 1,153)	1,177	(1,088 - 1,268)
Pennsylvania	4,160	(3,969 - 4,354)	367	(348 - 386)	407	(381 - 433)	3,386	(3,200 - 3,576)	3,793	(3,605 - 3,984)
Rhode Island	337	(309 - 366)	29	(26 - 32)	38	(34 - 43)	270	(244 - 297)	308	(281 - 336)
South Carolina	1,752	(1,631 - 1,875)	149	(137 - 162)	190	(174 - 207)	1,413	(1,296 - 1,530)	1,603	(1,484 - 1,723)
South Dakota	223	(202 - 245)	20	(18 - 22)	23	(20 - 27)	179	(160 - 200)	203	(183 - 224)
Tennessee	2,347	(2,180 - 2,517)	204	(189 - 219)	223	(202 - 245)	1,920	(1,759 - 2,083)	2,143	(1,979 - 2,311)
Texas	8,777	(8,455 - 9,101)	876	(833 - 920)	1,035	(983 - 1,087)	6,866	(6,557 - 7,177)	7,901	(7,585 - 8,218)
Utah	1,099	(1,032 - 1,166)	124	(114 - 134)	145	(131 - 159)	830	(768 - 892)	975	(910 - 1,040)
Vermont	197	(180 - 214)	15	(13 - 16)	17	(15 - 19)	165	(149 - 182)	182	(166 - 199)
Virginia	2,745	(2,547 - 2,946)	241	(220 - 262)	281	(253 - 310)	2,223	(2,034 - 2,415)	2,504	(2,311 - 2,701)
Washington	2,321	(2,153 - 2,493)	192	(175 - 210)	209	(186 - 233)	1,920	(1,760 - 2,083)	2,129	(1,964 - 2,297)
West Virginia	680	(630 - 730)	51	(46 - 56)	63	(56 - 69)	566	(519 - 615)	629	(581 - 678)
Wisconsin	1,653	(1,514 - 1,796)	153	(138 - 168)	172	(153 - 193)	1,328	(1,198 - 1,463)	1,500	(1,366 - 1,640)
Wyoming	175	(161 - 189)	16	(15 - 18)	18	(16 - 21)	140	(127 - 153)	158	(145 - 172)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 12. Alcohol Use and Binge Alcohol Use in the Past Month among Persons Aged 12 to 20, by State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	Alcohol Use in Past Month (Estimate)	Alcohol Use in Past Month (95% Confidence Interval)	Binge Alcohol Use in Past Month (Estimate)	Binge Alcohol Use in Past Month (95% Confidence Interval)
Total U.S.	10,163	(9,932 - 10,397)	6,684	(6,481 - 6,893)
Northeast	2,050	(1,988 - 2,113)	1,335	(1,279 - 1,394)
Midwest	2,286	(2,230 - 2,343)	1,580	(1,525 - 1,637)
South	3,519	(3,429 - 3,611)	2,224	(2,140 - 2,310)
West	2,308	(2,224 - 2,394)	1,545	(1,467 - 1,627)
Alabama	132	(118 - 148)	81	(70 - 93)
Alaska	23	(21 - 26)	16	(15 - 19)
Arizona	195	(175 - 217)	136	(118 - 155)
Arkansas	75	(67 - 85)	45	(39 - 52)
California	1,209	(1,139 - 1,282)	806	(743 - 873)
Colorado	170	(154 - 187)	104	(92 - 118)
Connecticut	137	(124 - 149)	95	(84 - 108)
Delaware	34	(31 - 37)	23	(21 - 26)
District of Columbia	17	(15 - 19)	10	(8 - 11)
Florida	549	(519 - 581)	329	(303 - 357)
Georgia	281	(252 - 311)	176	(154 - 200)
Hawaii	37	(33 - 42)	26	(22 - 30)
Idaho	52	(47 - 57)	34	(30 - 38)
Illinois	477	(454 - 502)	325	(304 - 348)
Indiana	192	(172 - 213)	134	(118 - 152)
Iowa	108	(98 - 118)	77	(68 - 86)
Kansas	103	(94 - 113)	70	(62 - 79)
Kentucky	126	(114 - 140)	90	(79 - 101)
Louisiana	153	(139 - 168)	91	(80 - 104)
Maine	37	(33 - 41)	26	(23 - 29)
Maryland	170	(153 - 188)	105	(92 - 121)
Massachusetts	233	(214 - 253)	157	(140 - 175)
Michigan	318	(301 - 336)	214	(199 - 231)
Minnesota	180	(165 - 197)	132	(117 - 147)
Mississippi	99	(89 - 110)	61	(53 - 69)
Missouri	181	(163 - 201)	118	(103 - 135)
Montana	38	(34 - 41)	27	(24 - 30)

(continued)

Table 12. Alcohol Use and Binge Alcohol Use in the Past Month among Persons Aged 12 to 20, by State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	Alcohol Use in Past Month (Estimate)	Alcohol Use in Past Month (95% Confidence Interval)	Binge Alcohol Use in Past Month (Estimate)	Alcohol Use in Past Month (95% Confidence Interval)
Nebraska	62	(56 - 69)	45	(40 - 51)
Nevada	81	(73 - 90)	57	(50 - 64)
New Hampshire	54	(49 - 58)	35	(32 - 40)
New Jersey	297	(270 - 327)	203	(180 - 228)
New Mexico	64	(58 - 72)	42	(37 - 48)
New York	768	(731 - 806)	466	(433 - 501)
North Carolina	273	(245 - 302)	164	(142 - 188)
North Dakota	28	(26 - 31)	20	(18 - 22)
Ohio	401	(381 - 421)	281	(263 - 299)
Oklahoma	116	(104 - 129)	72	(62 - 83)
Oregon	133	(120 - 146)	78	(69 - 89)
Pennsylvania	455	(431 - 480)	307	(286 - 330)
Rhode Island	43	(39 - 47)	28	(25 - 31)
South Carolina	125	(112 - 139)	78	(67 - 89)
South Dakota	31	(28 - 34)	21	(19 - 23)
Tennessee	171	(151 - 192)	116	(101 - 133)
Texas	855	(810 - 902)	555	(515 - 597)
Utah	64	(54 - 75)	49	(41 - 59)
Vermont	26	(24 - 28)	18	(16 - 20)
Virginia	294	(269 - 321)	196	(174 - 219)
Washington	221	(200 - 243)	155	(138 - 174)
West Virginia	49	(44 - 55)	33	(29 - 38)
Wisconsin	204	(186 - 222)	143	(127 - 159)
Wyoming	21	(19 - 23)	15	(13 - 17)

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on at least 1 day in the past 30 days.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 13. Tobacco Product Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	69,738	(68,537 - 70,965)	2,753	(2,653 - 2,856)	13,960	(13,720 - 14,201)	53,025	(51,901 - 54,177)	66,985	(65,801 - 68,194)
Northeast	12,206	(11,765 - 12,659)	465	(436 - 497)	2,488	(2,413 - 2,564)	9,253	(8,820 - 9,699)	11,740	(11,306 - 12,186)
Midwest	16,632	(16,176 - 17,094)	675	(643 - 709)	3,396	(3,321 - 3,471)	12,561	(12,111 - 13,020)	15,957	(15,507 - 16,411)
South	26,860	(26,105 - 27,642)	1,014	(967 - 1,064)	5,089	(4,967 - 5,212)	20,756	(20,009 - 21,532)	25,845	(25,104 - 26,613)
West	14,040	(13,472 - 14,623)	598	(556 - 643)	2,987	(2,883 - 3,093)	10,455	(9,901 - 11,031)	13,442	(12,884 - 14,015)
Alabama	1,269	(1,161 - 1,383)	45	(38 - 53)	228	(209 - 248)	996	(895 - 1,103)	1,225	(1,118 - 1,336)
Alaska	177	(160 - 195)	8	(6 - 9)	37	(34 - 41)	132	(117 - 149)	169	(153 - 187)
Arizona	1,393	(1,249 - 1,547)	62	(51 - 75)	264	(238 - 290)	1,067	(936 - 1,210)	1,331	(1,190 - 1,481)
Arkansas	794	(728 - 862)	32	(27 - 38)	137	(126 - 148)	625	(563 - 689)	762	(698 - 828)
California	6,561	(6,140 - 6,999)	281	(248 - 317)	1,468	(1,383 - 1,555)	4,813	(4,411 - 5,240)	6,280	(5,867 - 6,712)
Colorado	1,110	(1,003 - 1,224)	47	(39 - 56)	234	(213 - 256)	828	(730 - 935)	1,062	(957 - 1,174)
Connecticut	738	(659 - 824)	31	(25 - 37)	155	(141 - 169)	552	(479 - 633)	708	(629 - 791)
Delaware	213	(193 - 235)	8	(7 - 10)	40	(36 - 43)	165	(146 - 186)	205	(184 - 226)
District of Columbia	141	(127 - 156)	3	(3 - 4)	31	(28 - 34)	108	(95 - 121)	138	(124 - 153)
Florida	4,078	(3,852 - 4,317)	139	(125 - 155)	695	(656 - 734)	3,244	(3,023 - 3,475)	3,939	(3,716 - 4,174)
Georgia	2,115	(1,907 - 2,338)	95	(79 - 113)	424	(386 - 464)	1,596	(1,406 - 1,803)	2,021	(1,818 - 2,238)
Hawaii	237	(212 - 266)	9	(7 - 11)	48	(43 - 54)	180	(156 - 206)	228	(203 - 256)
Idaho	345	(312 - 379)	16	(13 - 18)	71	(65 - 78)	258	(228 - 290)	329	(297 - 363)
Illinois	3,050	(2,892 - 3,214)	115	(103 - 129)	643	(614 - 673)	2,292	(2,140 - 2,450)	2,935	(2,779 - 3,096)
Indiana	1,615	(1,479 - 1,759)	64	(54 - 76)	332	(306 - 359)	1,219	(1,093 - 1,353)	1,551	(1,416 - 1,692)
Iowa	770	(701 - 843)	31	(27 - 37)	158	(145 - 171)	581	(517 - 649)	739	(670 - 809)
Kansas	702	(640 - 766)	31	(26 - 36)	146	(133 - 159)	525	(469 - 585)	671	(610 - 734)
Kentucky	1,265	(1,161 - 1,372)	50	(43 - 59)	216	(200 - 233)	998	(900 - 1,100)	1,214	(1,112 - 1,320)
Louisiana	1,099	(1,002 - 1,201)	40	(33 - 47)	221	(202 - 240)	838	(748 - 934)	1,059	(963 - 1,159)
Maine	326	(295 - 359)	12	(10 - 14)	60	(56 - 65)	254	(225 - 285)	314	(284 - 346)
Maryland	1,141	(1,007 - 1,286)	42	(34 - 52)	221	(198 - 247)	877	(755 - 1,012)	1,099	(968 - 1,241)
Massachusetts	1,355	(1,220 - 1,500)	51	(43 - 61)	325	(295 - 355)	979	(857 - 1,113)	1,304	(1,171 - 1,446)
Michigan	2,500	(2,373 - 2,632)	98	(88 - 108)	498	(476 - 520)	1,905	(1,782 - 2,032)	2,403	(2,277 - 2,532)
Minnesota	1,194	(1,089 - 1,305)	46	(39 - 55)	274	(254 - 294)	874	(777 - 979)	1,148	(1,044 - 1,256)
Mississippi	819	(751 - 891)	34	(28 - 40)	158	(145 - 170)	628	(564 - 695)	786	(718 - 855)
Missouri	1,701	(1,558 - 1,852)	75	(64 - 88)	306	(282 - 331)	1,320	(1,186 - 1,460)	1,626	(1,485 - 1,773)
Montana	262	(240 - 286)	12	(10 - 14)	53	(49 - 57)	197	(176 - 220)	250	(228 - 274)

(continued)

Table 13. Tobacco Product Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	424	(386 - 465)	17	(15 - 20)	90	(83 - 98)	317	(281 - 354)	407	(369 - 447)
Nevada	589	(528 - 655)	21	(17 - 26)	102	(92 - 112)	466	(408 - 528)	568	(508 - 633)
New Hampshire	327	(296 - 359)	14	(12 - 17)	66	(61 - 71)	247	(219 - 277)	313	(283 - 344)
New Jersey	1,853	(1,668 - 2,053)	68	(55 - 83)	369	(336 - 403)	1,417	(1,243 - 1,606)	1,786	(1,603 - 1,982)
New Mexico	444	(399 - 490)	21	(18 - 25)	98	(90 - 107)	324	(283 - 368)	422	(379 - 468)
New York	3,970	(3,731 - 4,219)	151	(134 - 169)	815	(769 - 861)	3,004	(2,775 - 3,247)	3,819	(3,583 - 4,065)
North Carolina	2,314	(2,107 - 2,533)	80	(67 - 95)	428	(392 - 465)	1,806	(1,612 - 2,012)	2,234	(2,030 - 2,449)
North Dakota	155	(141 - 169)	6	(5 - 7)	39	(36 - 43)	109	(97 - 123)	148	(135 - 163)
Ohio	2,989	(2,840 - 3,142)	130	(118 - 142)	578	(555 - 601)	2,281	(2,138 - 2,430)	2,859	(2,713 - 3,011)
Oklahoma	1,089	(999 - 1,183)	42	(35 - 50)	216	(199 - 233)	832	(748 - 919)	1,048	(958 - 1,139)
Oregon	914	(831 - 1,003)	37	(32 - 44)	172	(157 - 187)	705	(627 - 789)	877	(795 - 964)
Pennsylvania	3,251	(3,078 - 3,431)	125	(112 - 138)	613	(583 - 642)	2,514	(2,347 - 2,688)	3,127	(2,956 - 3,304)
Rhode Island	239	(215 - 266)	9	(7 - 10)	54	(49 - 59)	177	(155 - 201)	231	(207 - 256)
South Carolina	1,141	(1,038 - 1,249)	41	(34 - 48)	214	(196 - 233)	886	(791 - 988)	1,100	(999 - 1,206)
South Dakota	208	(189 - 228)	9	(7 - 10)	45	(41 - 48)	155	(138 - 174)	200	(181 - 219)
Tennessee	1,805	(1,650 - 1,968)	65	(55 - 76)	316	(292 - 340)	1,424	(1,278 - 1,577)	1,740	(1,586 - 1,899)
Texas	5,186	(4,898 - 5,482)	210	(188 - 235)	1,079	(1,024 - 1,135)	3,897	(3,624 - 4,183)	4,976	(4,692 - 5,269)
Utah	364	(316 - 416)	19	(14 - 24)	95	(83 - 109)	251	(209 - 298)	346	(299 - 397)
Vermont	145	(131 - 161)	6	(5 - 7)	32	(29 - 34)	108	(95 - 123)	140	(126 - 155)
Virginia	1,769	(1,603 - 1,946)	65	(55 - 78)	369	(338 - 401)	1,335	(1,181 - 1,501)	1,704	(1,541 - 1,877)
Washington	1,493	(1,355 - 1,639)	58	(48 - 69)	313	(287 - 340)	1,122	(994 - 1,259)	1,435	(1,300 - 1,578)
West Virginia	620	(571 - 671)	23	(19 - 26)	95	(88 - 102)	502	(455 - 551)	597	(548 - 647)
Wisconsin	1,322	(1,200 - 1,452)	52	(44 - 62)	286	(262 - 311)	984	(872 - 1,104)	1,270	(1,150 - 1,397)
Wyoming	150	(138 - 164)	7	(6 - 8)	31	(29 - 33)	112	(100 - 125)	143	(131 - 156)

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 14. Cigarette Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	58,524	(57,392 - 59,682)	2,128	(2,037 - 2,223)	11,870	(11,644 - 12,098)	44,526	(43,460 - 45,621)	56,396	(55,275 - 57,542)
Northeast	10,322	(9,910 - 10,749)	365	(339 - 393)	2,150	(2,072 - 2,229)	7,808	(7,397 - 8,235)	9,957	(9,549 - 10,380)
Midwest	13,879	(13,450 - 14,314)	530	(502 - 560)	2,880	(2,806 - 2,956)	10,468	(10,036 - 10,910)	13,348	(12,925 - 13,777)
South	22,440	(21,727 - 23,182)	781	(737 - 828)	4,308	(4,182 - 4,435)	17,352	(16,628 - 18,106)	21,659	(20,954 - 22,392)
West	11,883	(11,334 - 12,450)	452	(415 - 492)	2,532	(2,428 - 2,640)	8,899	(8,362 - 9,462)	11,431	(10,889 - 11,992)
Alabama	1,052	(957 - 1,152)	34	(28 - 41)	191	(172 - 212)	826	(737 - 922)	1,018	(924 - 1,117)
Alaska	145	(130 - 162)	5	(4 - 6)	32	(29 - 35)	109	(95 - 124)	141	(126 - 157)
Arizona	1,197	(1,067 - 1,339)	53	(43 - 66)	220	(196 - 247)	924	(804 - 1,055)	1,144	(1,015 - 1,283)
Arkansas	666	(605 - 731)	24	(20 - 29)	119	(107 - 130)	524	(467 - 585)	643	(582 - 707)
California	5,600	(5,198 - 6,022)	212	(183 - 244)	1,248	(1,163 - 1,337)	4,141	(3,761 - 4,549)	5,389	(4,992 - 5,806)
Colorado	872	(779 - 974)	33	(27 - 41)	187	(167 - 208)	652	(566 - 747)	839	(746 - 939)
Connecticut	627	(557 - 705)	25	(20 - 31)	133	(119 - 148)	469	(405 - 542)	603	(532 - 678)
Delaware	174	(156 - 193)	6	(5 - 8)	34	(30 - 37)	134	(117 - 152)	167	(149 - 187)
District of Columbia	120	(107 - 135)	2	(2 - 3)	25	(22 - 29)	93	(80 - 106)	118	(105 - 132)
Florida	3,472	(3,256 - 3,702)	106	(93 - 121)	587	(551 - 624)	2,779	(2,566 - 3,004)	3,366	(3,151 - 3,595)
Georgia	1,771	(1,577 - 1,980)	76	(62 - 93)	358	(320 - 398)	1,336	(1,158 - 1,534)	1,695	(1,503 - 1,902)
Hawaii	207	(183 - 234)	7	(5 - 9)	43	(38 - 48)	158	(136 - 183)	201	(177 - 227)
Idaho	294	(266 - 325)	12	(10 - 14)	62	(56 - 68)	220	(194 - 249)	283	(254 - 313)
Illinois	2,587	(2,443 - 2,736)	91	(80 - 104)	556	(528 - 585)	1,939	(1,801 - 2,085)	2,496	(2,353 - 2,643)
Indiana	1,363	(1,238 - 1,497)	49	(40 - 60)	282	(256 - 309)	1,032	(916 - 1,157)	1,314	(1,190 - 1,446)
Iowa	635	(573 - 701)	26	(21 - 31)	132	(119 - 145)	478	(421 - 540)	609	(548 - 674)
Kansas	548	(496 - 603)	24	(20 - 29)	119	(107 - 132)	405	(358 - 457)	524	(472 - 578)
Kentucky	1,066	(971 - 1,167)	37	(30 - 45)	181	(164 - 198)	848	(759 - 944)	1,029	(935 - 1,128)
Louisiana	946	(852 - 1,046)	32	(26 - 39)	194	(175 - 213)	720	(634 - 814)	914	(821 - 1,013)
Maine	279	(251 - 309)	9	(7 - 11)	52	(47 - 57)	218	(192 - 246)	270	(242 - 299)
Maryland	931	(818 - 1,055)	34	(27 - 42)	186	(163 - 211)	711	(608 - 828)	897	(786 - 1,021)
Massachusetts	1,152	(1,034 - 1,280)	39	(31 - 48)	278	(250 - 306)	836	(728 - 955)	1,113	(996 - 1,240)
Michigan	2,155	(2,039 - 2,276)	78	(70 - 87)	430	(408 - 452)	1,647	(1,534 - 1,766)	2,077	(1,962 - 2,198)
Minnesota	1,014	(916 - 1,119)	37	(30 - 45)	236	(215 - 257)	742	(651 - 842)	977	(879 - 1,081)
Mississippi	651	(592 - 714)	25	(21 - 31)	131	(119 - 144)	495	(440 - 553)	626	(567 - 688)
Missouri	1,355	(1,231 - 1,486)	55	(46 - 67)	255	(231 - 279)	1,044	(929 - 1,168)	1,299	(1,177 - 1,430)
Montana	204	(185 - 224)	9	(7 - 11)	41	(37 - 45)	154	(136 - 174)	195	(176 - 215)

(continued)

Table 14. Cigarette Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	352	(318 - 389)	13	(11 - 16)	80	(73 - 88)	259	(227 - 294)	339	(305 - 376)
Nevada	509	(453 - 570)	16	(12 - 20)	85	(76 - 95)	408	(355 - 466)	494	(438 - 554)
New Hampshire	273	(246 - 302)	11	(9 - 13)	58	(53 - 63)	204	(179 - 232)	262	(235 - 291)
New Jersey	1,537	(1,367 - 1,723)	53	(43 - 66)	326	(293 - 361)	1,158	(999 - 1,334)	1,484	(1,316 - 1,668)
New Mexico	369	(331 - 410)	16	(13 - 20)	83	(75 - 91)	270	(235 - 309)	353	(315 - 393)
New York	3,472	(3,251 - 3,704)	118	(102 - 135)	709	(664 - 756)	2,645	(2,432 - 2,871)	3,354	(3,135 - 3,584)
North Carolina	1,990	(1,795 - 2,198)	59	(48 - 74)	378	(342 - 415)	1,552	(1,370 - 1,749)	1,930	(1,737 - 2,137)
North Dakota	122	(110 - 135)	5	(4 - 6)	31	(28 - 34)	86	(75 - 99)	117	(105 - 130)
Ohio	2,471	(2,333 - 2,615)	104	(93 - 116)	482	(459 - 506)	1,884	(1,749 - 2,026)	2,367	(2,230 - 2,510)
Oklahoma	865	(783 - 951)	32	(26 - 38)	183	(165 - 201)	650	(575 - 732)	833	(752 - 919)
Oregon	769	(694 - 850)	27	(22 - 34)	152	(137 - 167)	590	(519 - 666)	741	(667 - 822)
Pennsylvania	2,654	(2,496 - 2,820)	99	(88 - 112)	520	(490 - 550)	2,035	(1,882 - 2,197)	2,555	(2,398 - 2,720)
Rhode Island	207	(186 - 230)	7	(5 - 8)	46	(41 - 51)	154	(135 - 176)	201	(179 - 223)
South Carolina	992	(894 - 1,097)	32	(26 - 39)	182	(165 - 200)	778	(686 - 878)	960	(863 - 1,064)
South Dakota	167	(150 - 185)	7	(5 - 8)	36	(33 - 40)	124	(108 - 141)	160	(143 - 178)
Tennessee	1,451	(1,311 - 1,599)	45	(37 - 55)	257	(232 - 282)	1,149	(1,017 - 1,290)	1,405	(1,266 - 1,552)
Texas	4,335	(4,063 - 4,617)	170	(148 - 195)	918	(863 - 974)	3,248	(2,990 - 3,521)	4,165	(3,896 - 4,445)
Utah	343	(297 - 393)	15	(11 - 20)	89	(76 - 103)	239	(199 - 285)	328	(283 - 377)
Vermont	120	(108 - 134)	4	(3 - 5)	27	(25 - 30)	88	(77 - 101)	116	(103 - 129)
Virginia	1,466	(1,322 - 1,621)	51	(42 - 63)	304	(274 - 335)	1,111	(978 - 1,256)	1,415	(1,271 - 1,569)
Washington	1,257	(1,134 - 1,389)	41	(33 - 51)	266	(240 - 293)	950	(837 - 1,074)	1,216	(1,094 - 1,347)
West Virginia	493	(449 - 540)	15	(13 - 19)	81	(74 - 88)	397	(354 - 442)	478	(433 - 524)
Wisconsin	1,109	(1,001 - 1,225)	42	(34 - 51)	241	(217 - 266)	827	(728 - 934)	1,068	(961 - 1,182)
Wyoming	115	(104 - 127)	6	(5 - 7)	25	(23 - 27)	84	(74 - 96)	109	(98 - 121)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 15. Perceptions of Great Risk of Smoking One or More Packs of Cigarettes Per Day, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	179,963	(178,777 - 181,117)	16,012	(15,856 - 16,166)	22,037	(21,827 - 22,245)	141,914	(140,761 - 143,032)	163,951	(162,783 - 165,086)
Northeast	34,149	(33,700 - 34,585)	2,907	(2,857 - 2,957)	4,076	(3,998 - 4,153)	27,165	(26,750 - 27,566)	31,241	(30,805 - 31,665)
Midwest	37,252	(36,768 - 37,729)	3,457	(3,405 - 3,508)	4,529	(4,449 - 4,609)	29,266	(28,821 - 29,702)	33,795	(33,322 - 34,261)
South	65,648	(64,899 - 66,374)	5,759	(5,679 - 5,839)	7,910	(7,786 - 8,032)	51,979	(51,278 - 52,652)	59,888	(59,154 - 60,599)
West	42,914	(42,319 - 43,491)	3,888	(3,814 - 3,961)	5,522	(5,411 - 5,629)	33,504	(32,951 - 34,036)	39,026	(38,444 - 39,588)
Alabama	2,616	(2,501 - 2,725)	227	(213 - 239)	312	(292 - 331)	2,077	(1,969 - 2,179)	2,389	(2,276 - 2,496)
Alaska	373	(356 - 389)	39	(37 - 41)	51	(48 - 54)	283	(267 - 298)	334	(317 - 349)
Arizona	3,838	(3,676 - 3,991)	350	(330 - 370)	462	(436 - 486)	3,025	(2,873 - 3,165)	3,487	(3,328 - 3,636)
Arkansas	1,661	(1,595 - 1,724)	147	(138 - 154)	184	(173 - 195)	1,331	(1,268 - 1,389)	1,515	(1,450 - 1,576)
California	22,768	(22,311 - 23,206)	2,066	(2,006 - 2,124)	3,077	(2,989 - 3,162)	17,625	(17,193 - 18,033)	20,702	(20,254 - 21,129)
Colorado	2,984	(2,865 - 3,097)	255	(241 - 267)	365	(346 - 383)	2,365	(2,252 - 2,469)	2,730	(2,613 - 2,839)
Connecticut	2,152	(2,067 - 2,232)	195	(185 - 204)	248	(235 - 262)	1,709	(1,627 - 1,783)	1,957	(1,873 - 2,035)
Delaware	538	(517 - 558)	45	(43 - 48)	62	(59 - 65)	431	(410 - 449)	493	(472 - 512)
District of Columbia	387	(374 - 400)	23	(21 - 24)	60	(57 - 63)	304	(291 - 316)	365	(351 - 377)
Florida	11,638	(11,405 - 11,861)	890	(864 - 915)	1,276	(1,239 - 1,312)	9,472	(9,249 - 9,684)	10,748	(10,519 - 10,967)
Georgia	5,634	(5,419 - 5,837)	527	(499 - 553)	701	(665 - 735)	4,407	(4,203 - 4,595)	5,108	(4,898 - 5,304)
Hawaii	756	(725 - 786)	61	(58 - 64)	87	(82 - 92)	608	(578 - 636)	695	(665 - 724)
Idaho	817	(782 - 852)	85	(80 - 89)	103	(97 - 109)	629	(595 - 661)	732	(697 - 766)
Illinois	7,337	(7,165 - 7,503)	686	(666 - 706)	909	(880 - 938)	5,741	(5,578 - 5,898)	6,650	(6,482 - 6,813)
Indiana	3,489	(3,336 - 3,636)	340	(322 - 356)	414	(389 - 438)	2,735	(2,590 - 2,873)	3,149	(3,000 - 3,292)
Iowa	1,629	(1,553 - 1,702)	146	(137 - 153)	205	(192 - 217)	1,279	(1,207 - 1,347)	1,483	(1,409 - 1,555)
Kansas	1,485	(1,417 - 1,550)	133	(125 - 141)	204	(192 - 216)	1,148	(1,083 - 1,208)	1,351	(1,285 - 1,415)
Kentucky	2,219	(2,108 - 2,326)	212	(200 - 223)	249	(233 - 265)	1,757	(1,652 - 1,858)	2,007	(1,898 - 2,112)
Louisiana	2,627	(2,527 - 2,722)	239	(226 - 252)	343	(325 - 360)	2,045	(1,950 - 2,133)	2,388	(2,290 - 2,480)
Maine	775	(742 - 806)	65	(62 - 68)	78	(74 - 83)	632	(600 - 662)	710	(678 - 741)
Maryland	3,537	(3,399 - 3,664)	298	(282 - 313)	429	(407 - 450)	2,809	(2,680 - 2,926)	3,238	(3,104 - 3,362)
Massachusetts	4,240	(4,099 - 4,371)	348	(331 - 363)	524	(497 - 550)	3,368	(3,234 - 3,490)	3,892	(3,753 - 4,021)
Michigan	5,676	(5,540 - 5,809)	527	(512 - 542)	687	(666 - 707)	4,463	(4,333 - 4,588)	5,150	(5,015 - 5,280)
Minnesota	2,975	(2,842 - 3,102)	279	(264 - 292)	354	(334 - 374)	2,342	(2,216 - 2,460)	2,696	(2,566 - 2,821)
Mississippi	1,640	(1,572 - 1,705)	152	(143 - 160)	213	(202 - 224)	1,275	(1,211 - 1,334)	1,488	(1,422 - 1,551)
Missouri	3,277	(3,130 - 3,418)	295	(278 - 312)	378	(355 - 401)	2,604	(2,462 - 2,737)	2,982	(2,837 - 3,120)
Montana	556	(532 - 578)	47	(45 - 50)	67	(63 - 70)	442	(419 - 463)	508	(485 - 531)

(continued)

Table 15. Perceptions of Great Risk of Smoking One or More Packs of Cigarettes Per Day, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	982	(940 - 1,022)	89	(84 - 94)	129	(122 - 137)	764	(724 - 801)	893	(851 - 932)
Nevada	1,518	(1,453 - 1,578)	140	(132 - 147)	174	(165 - 183)	1,204	(1,143 - 1,260)	1,378	(1,315 - 1,437)
New Hampshire	785	(751 - 818)	67	(63 - 70)	90	(85 - 95)	628	(596 - 659)	719	(685 - 750)
New Jersey	5,361	(5,164 - 5,545)	467	(444 - 489)	602	(571 - 632)	4,292	(4,104 - 4,463)	4,894	(4,701 - 5,073)
New Mexico	1,187	(1,144 - 1,229)	103	(97 - 108)	148	(139 - 155)	937	(896 - 975)	1,085	(1,042 - 1,125)
New York	12,527	(12,271 - 12,772)	1,073	(1,043 - 1,103)	1,558	(1,513 - 1,601)	9,896	(9,654 - 10,124)	11,454	(11,203 - 11,692)
North Carolina	5,273	(5,050 - 5,486)	483	(458 - 507)	627	(593 - 660)	4,163	(3,950 - 4,363)	4,790	(4,570 - 4,998)
North Dakota	366	(351 - 381)	31	(29 - 32)	59	(56 - 63)	276	(261 - 290)	335	(320 - 350)
Ohio	6,359	(6,204 - 6,511)	593	(576 - 610)	735	(711 - 758)	5,031	(4,881 - 5,176)	5,766	(5,613 - 5,915)
Oklahoma	1,975	(1,886 - 2,061)	172	(161 - 182)	251	(234 - 267)	1,552	(1,468 - 1,632)	1,804	(1,716 - 1,887)
Oregon	2,262	(2,167 - 2,351)	188	(178 - 197)	243	(228 - 257)	1,831	(1,740 - 1,915)	2,074	(1,980 - 2,162)
Pennsylvania	7,288	(7,098 - 7,471)	611	(590 - 631)	842	(812 - 871)	5,836	(5,654 - 6,010)	6,677	(6,491 - 6,858)
Rhode Island	639	(612 - 664)	52	(49 - 55)	87	(82 - 91)	500	(475 - 523)	587	(561 - 611)
South Carolina	2,634	(2,525 - 2,737)	218	(206 - 230)	303	(286 - 320)	2,112	(2,008 - 2,208)	2,416	(2,309 - 2,516)
South Dakota	440	(420 - 460)	41	(39 - 43)	58	(55 - 62)	341	(322 - 359)	399	(379 - 418)
Tennessee	3,444	(3,277 - 3,605)	300	(283 - 316)	375	(350 - 398)	2,770	(2,609 - 2,921)	3,144	(2,979 - 3,302)
Texas	14,195	(13,891 - 14,489)	1,350	(1,309 - 1,391)	1,850	(1,794 - 1,905)	10,995	(10,704 - 11,272)	12,845	(12,545 - 13,133)
Utah	1,596	(1,542 - 1,646)	185	(176 - 193)	260	(246 - 273)	1,150	(1,101 - 1,195)	1,411	(1,358 - 1,460)
Vermont	381	(366 - 396)	30	(28 - 31)	47	(44 - 49)	304	(290 - 318)	351	(336 - 366)
Virginia	4,633	(4,450 - 4,805)	397	(377 - 416)	569	(539 - 598)	3,667	(3,495 - 3,825)	4,235	(4,056 - 4,403)
Washington	3,951	(3,806 - 4,089)	342	(325 - 358)	447	(422 - 470)	3,163	(3,024 - 3,292)	3,609	(3,466 - 3,745)
West Virginia	996	(949 - 1,042)	80	(76 - 85)	105	(98 - 111)	811	(765 - 855)	916	(869 - 961)
Wisconsin	3,237	(3,103 - 3,365)	297	(283 - 311)	396	(373 - 419)	2,543	(2,415 - 2,664)	2,940	(2,807 - 3,066)
Wyoming	308	(295 - 320)	28	(26 - 29)	38	(36 - 40)	242	(230 - 253)	280	(268 - 292)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 16. Alcohol Dependence or Abuse in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	18,365	(17,797 - 18,949)	1,123	(1,057 - 1,192)	5,382	(5,216 - 5,553)	11,860	(11,329 - 12,414)	17,242	(16,682 - 17,819)
Northeast	3,366	(3,172 - 3,572)	197	(178 - 217)	1,031	(970 - 1,096)	2,138	(1,959 - 2,332)	3,169	(2,979 - 3,372)
Midwest	4,121	(3,934 - 4,315)	246	(228 - 266)	1,247	(1,184 - 1,312)	2,628	(2,455 - 2,812)	3,874	(3,691 - 4,065)
South	6,313	(6,011 - 6,629)	375	(344 - 409)	1,785	(1,686 - 1,888)	4,152	(3,869 - 4,454)	5,937	(5,640 - 6,249)
West	4,566	(4,287 - 4,860)	305	(275 - 338)	1,319	(1,234 - 1,409)	2,942	(2,687 - 3,219)	4,261	(3,988 - 4,550)
Alabama	265	(220 - 318)	16	(12 - 21)	71	(59 - 84)	178	(140 - 226)	249	(206 - 300)
Alaska	43	(36 - 52)	3	(2 - 4)	13	(11 - 15)	28	(22 - 35)	40	(34 - 48)
Arizona	414	(345 - 495)	26	(20 - 34)	119	(101 - 139)	269	(210 - 343)	388	(321 - 467)
Arkansas	156	(130 - 186)	11	(8 - 14)	43	(36 - 51)	102	(80 - 130)	145	(121 - 175)
California	2,422	(2,200 - 2,664)	167	(142 - 196)	689	(623 - 761)	1,566	(1,366 - 1,793)	2,255	(2,038 - 2,492)
Colorado	359	(304 - 423)	22	(17 - 28)	110	(96 - 126)	227	(180 - 286)	337	(284 - 399)
Connecticut	231	(194 - 274)	14	(11 - 18)	70	(59 - 82)	148	(116 - 187)	218	(182 - 259)
Delaware	53	(44 - 64)	3	(2 - 4)	16	(13 - 18)	34	(27 - 44)	50	(41 - 61)
District of Columbia	49	(41 - 58)	1	(1 - 2)	15	(13 - 17)	33	(26 - 41)	48	(40 - 56)
Florida	1,054	(947 - 1,173)	59	(50 - 70)	255	(229 - 284)	739	(640 - 853)	994	(889 - 1,113)
Georgia	498	(415 - 598)	30	(23 - 39)	148	(126 - 175)	320	(249 - 410)	468	(387 - 566)
Hawaii	81	(66 - 98)	5	(4 - 7)	22	(19 - 27)	53	(41 - 69)	76	(62 - 93)
Idaho	94	(79 - 110)	7	(5 - 9)	28	(24 - 33)	58	(46 - 73)	87	(73 - 103)
Illinois	765	(692 - 845)	43	(36 - 51)	247	(225 - 270)	475	(409 - 551)	722	(650 - 800)
Indiana	381	(321 - 450)	23	(18 - 29)	112	(95 - 131)	245	(195 - 308)	357	(300 - 425)
Iowa	206	(175 - 241)	13	(10 - 16)	69	(60 - 80)	124	(99 - 155)	194	(163 - 227)
Kansas	182	(155 - 213)	12	(10 - 15)	59	(51 - 69)	110	(88 - 138)	170	(143 - 200)
Kentucky	219	(182 - 262)	14	(10 - 18)	63	(53 - 74)	143	(112 - 181)	205	(170 - 247)
Louisiana	241	(201 - 288)	12	(9 - 17)	75	(63 - 88)	154	(120 - 197)	229	(190 - 274)
Maine	73	(61 - 87)	4	(3 - 5)	21	(18 - 25)	47	(37 - 60)	69	(57 - 83)
Maryland	303	(246 - 371)	15	(11 - 20)	85	(70 - 103)	202	(155 - 263)	288	(232 - 355)
Massachusetts	464	(395 - 542)	25	(20 - 32)	152	(132 - 175)	286	(228 - 356)	438	(372 - 515)
Michigan	640	(578 - 707)	37	(31 - 43)	173	(157 - 190)	430	(374 - 493)	603	(543 - 669)
Minnesota	341	(289 - 402)	20	(16 - 26)	102	(88 - 119)	219	(174 - 274)	321	(270 - 380)
Mississippi	154	(129 - 183)	9	(7 - 12)	45	(38 - 54)	99	(78 - 126)	145	(120 - 173)
Missouri	328	(273 - 392)	22	(17 - 28)	95	(80 - 113)	210	(164 - 268)	306	(253 - 368)
Montana	76	(64 - 88)	5	(4 - 6)	25	(22 - 29)	45	(36 - 57)	71	(60 - 83)

(continued)

Table 16. Alcohol Dependence or Abuse in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	112	(95 - 131)	7	(5 - 8)	39	(33 - 45)	67	(53 - 84)	105	(89 - 124)
Nevada	181	(152 - 216)	11	(9 - 15)	47	(40 - 54)	123	(97 - 155)	170	(142 - 204)
New Hampshire	84	(71 - 100)	5	(4 - 7)	26	(23 - 30)	53	(41 - 67)	79	(66 - 94)
New Jersey	464	(385 - 559)	27	(21 - 36)	127	(107 - 149)	310	(240 - 398)	437	(360 - 529)
New Mexico	125	(105 - 149)	9	(7 - 11)	37	(31 - 44)	79	(62 - 100)	116	(97 - 139)
New York	1,204	(1,084 - 1,337)	75	(64 - 88)	371	(338 - 407)	758	(650 - 882)	1,129	(1,011 - 1,260)
North Carolina	479	(401 - 573)	26	(20 - 34)	136	(115 - 160)	317	(249 - 403)	453	(377 - 545)
North Dakota	45	(39 - 52)	2	(2 - 3)	18	(16 - 21)	25	(20 - 31)	43	(37 - 50)
Ohio	706	(640 - 778)	43	(37 - 50)	205	(188 - 224)	457	(396 - 527)	663	(598 - 735)
Oklahoma	229	(193 - 271)	14	(11 - 18)	68	(57 - 81)	147	(117 - 185)	216	(181 - 256)
Oregon	256	(216 - 304)	15	(11 - 19)	66	(56 - 77)	175	(139 - 219)	241	(202 - 288)
Pennsylvania	725	(647 - 812)	40	(34 - 48)	226	(204 - 249)	459	(390 - 540)	685	(609 - 771)
Rhode Island	78	(65 - 94)	4	(3 - 5)	23	(20 - 27)	51	(40 - 65)	75	(62 - 90)
South Carolina	301	(249 - 364)	14	(10 - 18)	70	(58 - 83)	218	(171 - 277)	288	(237 - 349)
South Dakota	61	(51 - 71)	4	(3 - 5)	19	(16 - 22)	38	(30 - 47)	57	(48 - 67)
Tennessee	333	(276 - 401)	18	(14 - 24)	90	(76 - 107)	225	(175 - 288)	315	(259 - 382)
Texas	1,361	(1,231 - 1,504)	97	(82 - 115)	415	(375 - 457)	849	(734 - 981)	1,264	(1,137 - 1,404)
Utah	124	(102 - 150)	10	(8 - 13)	44	(36 - 53)	70	(53 - 92)	114	(92 - 139)
Vermont	43	(36 - 50)	2	(2 - 3)	14	(12 - 17)	26	(21 - 33)	41	(34 - 48)
Virginia	523	(444 - 614)	30	(23 - 38)	163	(141 - 188)	330	(262 - 414)	493	(416 - 582)
Washington	355	(296 - 424)	22	(17 - 29)	107	(90 - 127)	226	(176 - 288)	333	(276 - 400)
West Virginia	93	(77 - 112)	6	(4 - 7)	26	(22 - 31)	62	(48 - 79)	88	(72 - 106)
Wisconsin	355	(300 - 420)	20	(16 - 26)	107	(91 - 125)	228	(181 - 287)	335	(281 - 398)
Wyoming	35	(30 - 41)	2	(2 - 3)	11	(10 - 13)	21	(17 - 27)	33	(28 - 39)

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 17. Alcohol Dependence in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	8,643	(8,243 - 9,060)	443	(404 - 485)	2,226	(2,120 - 2,338)	5,974	(5,592 - 6,380)	8,200	(7,805 - 8,614)
Northeast	1,568	(1,429 - 1,719)	78	(68 - 91)	425	(390 - 464)	1,064	(934 - 1,212)	1,489	(1,353 - 1,639)
Midwest	1,867	(1,739 - 2,003)	94	(84 - 106)	520	(486 - 556)	1,253	(1,130 - 1,389)	1,772	(1,646 - 1,907)
South	2,989	(2,770 - 3,224)	149	(131 - 168)	730	(679 - 785)	2,110	(1,899 - 2,343)	2,840	(2,624 - 3,074)
West	2,220	(2,015 - 2,443)	121	(102 - 143)	551	(500 - 606)	1,548	(1,354 - 1,767)	2,098	(1,897 - 2,319)
Alabama	125	(98 - 158)	6	(4 - 8)	30	(23 - 39)	89	(64 - 122)	119	(93 - 152)
Alaska	21	(16 - 26)	1	(1 - 1)	5	(4 - 7)	15	(11 - 20)	20	(15 - 25)
Arizona	205	(159 - 262)	10	(7 - 15)	52	(40 - 66)	142	(101 - 200)	194	(150 - 251)
Arkansas	75	(60 - 94)	4	(3 - 6)	19	(15 - 24)	52	(38 - 70)	71	(55 - 90)
California	1,206	(1,041 - 1,395)	67	(52 - 86)	285	(244 - 332)	855	(700 - 1,041)	1,139	(977 - 1,327)
Colorado	153	(121 - 192)	8	(6 - 11)	38	(30 - 48)	107	(78 - 146)	145	(114 - 184)
Connecticut	96	(75 - 122)	5	(4 - 8)	25	(20 - 33)	65	(46 - 90)	90	(70 - 116)
Delaware	24	(19 - 30)	1	(1 - 2)	6	(5 - 8)	16	(11 - 22)	22	(17 - 28)
District of Columbia	24	(19 - 30)	0	(0 - 1)	6	(5 - 8)	17	(13 - 23)	23	(19 - 29)
Florida	531	(451 - 625)	24	(18 - 30)	104	(88 - 123)	403	(326 - 498)	507	(428 - 601)
Georgia	228	(174 - 299)	11	(8 - 17)	56	(43 - 73)	161	(111 - 233)	217	(164 - 287)
Hawaii	37	(28 - 47)	2	(1 - 3)	10	(7 - 13)	25	(18 - 35)	35	(27 - 45)
Idaho	43	(35 - 53)	3	(2 - 3)	12	(10 - 15)	28	(21 - 38)	41	(33 - 51)
Illinois	354	(305 - 411)	16	(13 - 21)	110	(96 - 127)	227	(182 - 283)	338	(289 - 394)
Indiana	170	(136 - 212)	10	(7 - 14)	51	(41 - 64)	109	(78 - 151)	160	(127 - 202)
Iowa	85	(68 - 107)	5	(4 - 7)	27	(21 - 34)	54	(38 - 75)	80	(63 - 102)
Kansas	74	(60 - 92)	4	(3 - 6)	25	(19 - 31)	45	(33 - 63)	70	(56 - 87)
Kentucky	103	(79 - 134)	5	(4 - 8)	26	(20 - 33)	72	(50 - 102)	97	(74 - 128)
Louisiana	114	(91 - 144)	5	(4 - 8)	31	(24 - 39)	78	(56 - 108)	109	(86 - 139)
Maine	34	(26 - 43)	2	(1 - 2)	8	(6 - 10)	24	(17 - 34)	32	(25 - 42)
Maryland	158	(122 - 203)	7	(5 - 10)	36	(27 - 46)	115	(83 - 161)	151	(116 - 196)
Massachusetts	203	(164 - 251)	10	(7 - 13)	60	(48 - 74)	134	(98 - 182)	194	(155 - 241)
Michigan	304	(263 - 352)	14	(11 - 18)	66	(57 - 77)	223	(184 - 271)	290	(249 - 337)
Minnesota	167	(133 - 208)	8	(6 - 11)	44	(35 - 54)	115	(85 - 156)	159	(126 - 200)
Mississippi	75	(59 - 94)	4	(2 - 5)	20	(16 - 25)	51	(37 - 70)	71	(56 - 90)
Missouri	155	(122 - 196)	8	(6 - 11)	39	(30 - 50)	108	(78 - 149)	147	(115 - 188)
Montana	30	(24 - 38)	2	(1 - 3)	11	(9 - 13)	18	(12 - 25)	28	(23 - 36)

(continued)

Table 17. Alcohol Dependence in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	46	(37 - 58)	2	(2 - 3)	14	(11 - 18)	30	(21 - 42)	44	(34 - 56)
Nevada	80	(63 - 101)	4	(3 - 6)	20	(16 - 25)	56	(40 - 77)	76	(59 - 96)
New Hampshire	37	(30 - 47)	2	(2 - 3)	10	(8 - 13)	25	(18 - 34)	35	(28 - 44)
New Jersey	221	(172 - 284)	11	(8 - 16)	52	(40 - 66)	158	(113 - 221)	210	(161 - 272)
New Mexico	64	(51 - 81)	4	(3 - 5)	17	(14 - 22)	43	(31 - 60)	61	(47 - 78)
New York	593	(509 - 690)	31	(24 - 39)	167	(145 - 193)	395	(316 - 492)	562	(479 - 658)
North Carolina	241	(192 - 302)	11	(8 - 16)	57	(44 - 73)	173	(127 - 234)	230	(181 - 290)
North Dakota	20	(16 - 24)	1	(1 - 1)	7	(6 - 9)	11	(8 - 16)	19	(15 - 23)
Ohio	305	(263 - 354)	15	(12 - 19)	85	(74 - 96)	205	(166 - 253)	290	(248 - 338)
Oklahoma	93	(74 - 116)	5	(3 - 7)	28	(22 - 36)	60	(43 - 82)	88	(69 - 111)
Oregon	131	(102 - 168)	6	(4 - 8)	27	(21 - 34)	99	(71 - 135)	125	(97 - 162)
Pennsylvania	329	(278 - 388)	15	(12 - 20)	88	(75 - 103)	226	(178 - 285)	314	(264 - 373)
Rhode Island	35	(28 - 44)	1	(1 - 2)	9	(7 - 12)	24	(18 - 33)	34	(26 - 43)
South Carolina	148	(114 - 193)	5	(3 - 7)	26	(20 - 34)	117	(84 - 163)	144	(109 - 188)
South Dakota	25	(20 - 31)	1	(1 - 2)	7	(6 - 9)	16	(11 - 22)	23	(18 - 29)
Tennessee	167	(131 - 214)	8	(5 - 11)	37	(28 - 48)	123	(89 - 169)	160	(124 - 206)
Texas	619	(533 - 718)	39	(31 - 49)	170	(146 - 196)	411	(332 - 507)	580	(496 - 678)
Utah	64	(50 - 80)	5	(3 - 7)	21	(16 - 27)	38	(27 - 54)	59	(46 - 75)
Vermont	20	(16 - 25)	1	(1 - 1)	5	(4 - 7)	14	(10 - 19)	19	(15 - 24)
Virginia	215	(173 - 267)	11	(8 - 15)	68	(55 - 85)	136	(99 - 187)	204	(163 - 256)
Washington	170	(134 - 214)	9	(7 - 13)	48	(38 - 61)	112	(80 - 156)	160	(126 - 204)
West Virginia	49	(38 - 63)	2	(2 - 3)	11	(9 - 14)	35	(25 - 50)	47	(36 - 61)
Wisconsin	161	(128 - 203)	8	(6 - 11)	44	(35 - 56)	109	(80 - 150)	154	(121 - 194)
Wyoming	16	(13 - 20)	1	(1 - 1)	5	(4 - 6)	10	(7 - 14)	15	(12 - 19)

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 18. *Illicit Drug Dependence or Abuse in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs*

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	7,129	(6,808 - 7,464)	1,106	(1,045 - 1,170)	2,637	(2,522 - 2,757)	3,386	(3,106 - 3,692)	6,023	(5,713 - 6,349)
Northeast	1,364	(1,260 - 1,477)	194	(176 - 214)	534	(494 - 576)	636	(549 - 737)	1,170	(1,069 - 1,282)
Midwest	1,480	(1,382 - 1,584)	237	(218 - 257)	558	(523 - 595)	685	(602 - 778)	1,243	(1,148 - 1,345)
South	2,494	(2,333 - 2,666)	379	(349 - 411)	874	(819 - 931)	1,242	(1,102 - 1,399)	2,115	(1,959 - 2,284)
West	1,791	(1,638 - 1,957)	297	(268 - 329)	671	(616 - 730)	823	(698 - 970)	1,494	(1,348 - 1,656)
Alabama	104	(82 - 132)	15	(11 - 19)	35	(27 - 45)	54	(37 - 79)	89	(68 - 117)
Alaska	15	(12 - 19)	3	(2 - 3)	6	(5 - 8)	7	(5 - 10)	13	(10 - 17)
Arizona	164	(129 - 207)	26	(20 - 35)	57	(45 - 72)	80	(54 - 119)	137	(105 - 179)
Arkansas	62	(50 - 78)	10	(8 - 13)	20	(15 - 25)	33	(23 - 47)	52	(40 - 68)
California	950	(830 - 1,087)	158	(135 - 186)	355	(310 - 405)	437	(340 - 561)	792	(676 - 927)
Colorado	132	(105 - 165)	24	(18 - 31)	52	(41 - 64)	56	(37 - 85)	108	(83 - 140)
Connecticut	86	(68 - 108)	13	(10 - 16)	34	(27 - 43)	39	(26 - 58)	73	(56 - 95)
Delaware	20	(16 - 25)	3	(3 - 4)	8	(6 - 10)	9	(6 - 14)	17	(13 - 22)
District of Columbia	20	(15 - 26)	1	(1 - 2)	6	(5 - 8)	12	(9 - 18)	19	(14 - 24)
Florida	421	(364 - 488)	58	(49 - 69)	137	(119 - 157)	227	(177 - 290)	363	(307 - 430)
Georgia	175	(135 - 227)	32	(23 - 43)	61	(47 - 79)	82	(52 - 130)	143	(105 - 195)
Hawaii	27	(21 - 35)	4	(3 - 5)	10	(7 - 13)	14	(9 - 21)	23	(17 - 31)
Idaho	39	(31 - 49)	6	(5 - 8)	14	(11 - 18)	19	(13 - 27)	33	(26 - 42)
Illinois	284	(247 - 326)	45	(38 - 54)	117	(102 - 134)	122	(93 - 159)	238	(203 - 279)
Indiana	157	(125 - 196)	22	(17 - 29)	56	(45 - 71)	78	(54 - 113)	135	(104 - 174)
Iowa	52	(40 - 68)	11	(9 - 15)	19	(14 - 26)	22	(13 - 36)	41	(29 - 57)
Kansas	58	(45 - 73)	9	(7 - 12)	25	(19 - 31)	24	(16 - 37)	49	(37 - 64)
Kentucky	96	(77 - 119)	16	(12 - 21)	35	(28 - 43)	45	(31 - 66)	80	(62 - 103)
Louisiana	99	(79 - 123)	14	(11 - 18)	38	(30 - 48)	46	(31 - 68)	85	(66 - 109)
Maine	29	(23 - 37)	4	(3 - 5)	13	(10 - 16)	13	(9 - 19)	25	(20 - 33)
Maryland	139	(109 - 177)	18	(13 - 24)	52	(41 - 66)	69	(47 - 102)	121	(93 - 159)
Massachusetts	185	(148 - 231)	26	(20 - 34)	71	(57 - 88)	88	(61 - 128)	159	(124 - 204)
Michigan	228	(199 - 262)	36	(31 - 43)	84	(73 - 95)	109	(85 - 139)	192	(164 - 225)
Minnesota	109	(87 - 137)	17	(13 - 23)	43	(34 - 54)	49	(33 - 73)	92	(71 - 119)
Mississippi	71	(57 - 88)	10	(7 - 13)	25	(20 - 31)	36	(25 - 51)	61	(48 - 78)
Missouri	132	(106 - 165)	20	(15 - 26)	51	(40 - 64)	62	(42 - 90)	112	(87 - 145)
Montana	22	(18 - 28)	4	(3 - 5)	9	(7 - 11)	10	(6 - 14)	18	(14 - 24)

(continued)

Table 18. Illicit Drug Dependence or Abuse in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	33	(25 - 42)	6	(4 - 7)	12	(9 - 16)	15	(10 - 23)	27	(20 - 37)
Nevada	62	(48 - 80)	9	(7 - 13)	18	(14 - 23)	35	(24 - 51)	53	(40 - 71)
New Hampshire	38	(31 - 48)	7	(5 - 9)	15	(12 - 18)	17	(11 - 25)	32	(25 - 41)
New Jersey	188	(148 - 238)	27	(21 - 36)	68	(53 - 86)	93	(63 - 136)	160	(123 - 210)
New Mexico	52	(42 - 65)	11	(9 - 14)	19	(15 - 24)	22	(15 - 33)	41	(31 - 53)
New York	490	(426 - 564)	68	(57 - 81)	201	(176 - 228)	222	(171 - 287)	422	(360 - 495)
North Carolina	211	(167 - 267)	30	(23 - 39)	76	(61 - 96)	105	(72 - 154)	182	(139 - 237)
North Dakota	11	(8 - 14)	2	(1 - 2)	5	(4 - 6)	4	(3 - 7)	9	(7 - 12)
Ohio	273	(239 - 313)	42	(36 - 50)	97	(86 - 110)	133	(104 - 170)	231	(197 - 270)
Oklahoma	92	(73 - 117)	14	(11 - 19)	34	(26 - 43)	44	(30 - 65)	78	(60 - 102)
Oregon	102	(82 - 128)	16	(12 - 20)	37	(29 - 46)	49	(34 - 72)	86	(67 - 111)
Pennsylvania	300	(259 - 348)	43	(36 - 52)	112	(98 - 129)	145	(111 - 188)	257	(217 - 304)
Rhode Island	31	(24 - 38)	4	(3 - 5)	13	(11 - 17)	13	(9 - 19)	26	(20 - 34)
South Carolina	106	(84 - 135)	14	(11 - 19)	34	(27 - 43)	58	(40 - 84)	92	(70 - 121)
South Dakota	16	(12 - 20)	3	(2 - 4)	6	(4 - 7)	7	(5 - 11)	13	(9 - 17)
Tennessee	136	(107 - 171)	20	(15 - 26)	42	(33 - 54)	74	(51 - 106)	116	(88 - 151)
Texas	516	(449 - 592)	92	(77 - 110)	185	(160 - 213)	238	(185 - 307)	423	(360 - 498)
Utah	64	(52 - 79)	12	(9 - 15)	27	(22 - 35)	25	(17 - 36)	52	(41 - 67)
Vermont	17	(14 - 21)	2	(2 - 3)	8	(6 - 10)	7	(5 - 10)	15	(12 - 19)
Virginia	184	(147 - 230)	28	(21 - 36)	70	(56 - 87)	87	(59 - 127)	156	(121 - 202)
Washington	151	(120 - 189)	22	(17 - 29)	64	(51 - 80)	65	(44 - 97)	129	(100 - 167)
West Virginia	42	(32 - 53)	5	(3 - 6)	16	(13 - 20)	21	(14 - 31)	37	(28 - 49)
Wisconsin	127	(101 - 161)	23	(17 - 30)	44	(34 - 57)	60	(41 - 89)	104	(79 - 137)
Wyoming	10	(8 - 13)	2	(2 - 3)	4	(3 - 5)	4	(3 - 7)	8	(6 - 11)

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 19. *Illicit Drug Dependence in the Past Year*, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	4,868	(4,608 - 5,143)	596	(552 - 643)	1,836	(1,738 - 1,939)	2,436	(2,204 - 2,693)	4,272	(4,017 - 4,543)
Northeast	989	(896 - 1,091)	107	(94 - 122)	390	(354 - 429)	492	(416 - 582)	882	(791 - 984)
Midwest	1,021	(939 - 1,108)	127	(114 - 142)	381	(352 - 413)	512	(443 - 591)	893	(814 - 979)
South	1,679	(1,541 - 1,830)	203	(183 - 226)	603	(554 - 655)	872	(753 - 1,011)	1,475	(1,342 - 1,623)
West	1,180	(1,063 - 1,308)	158	(138 - 181)	462	(416 - 512)	560	(465 - 673)	1,022	(909 - 1,147)
Alabama	69	(53 - 90)	8	(6 - 11)	25	(18 - 34)	36	(24 - 53)	61	(45 - 82)
Alaska	11	(8 - 14)	2	(1 - 2)	4	(3 - 5)	5	(4 - 8)	9	(7 - 12)
Arizona	104	(80 - 134)	14	(10 - 20)	39	(29 - 53)	51	(34 - 74)	90	(67 - 120)
Arkansas	44	(34 - 57)	5	(4 - 8)	13	(10 - 18)	25	(17 - 36)	39	(29 - 52)
California	620	(531 - 724)	81	(66 - 101)	249	(211 - 293)	290	(222 - 378)	539	(453 - 640)
Colorado	76	(58 - 98)	13	(9 - 19)	28	(20 - 38)	35	(23 - 54)	63	(46 - 85)
Connecticut	60	(46 - 77)	7	(5 - 10)	23	(17 - 30)	30	(20 - 45)	53	(39 - 70)
Delaware	15	(12 - 20)	2	(1 - 3)	6	(5 - 8)	7	(5 - 10)	13	(10 - 18)
District of Columbia	13	(10 - 17)	1	(0 - 1)	4	(3 - 6)	8	(6 - 12)	12	(9 - 17)
Florida	258	(216 - 309)	30	(24 - 37)	91	(76 - 108)	137	(104 - 182)	228	(188 - 278)
Georgia	114	(83 - 158)	17	(12 - 25)	37	(25 - 53)	60	(37 - 98)	97	(67 - 141)
Hawaii	16	(12 - 22)	2	(1 - 3)	6	(4 - 9)	8	(5 - 13)	14	(10 - 20)
Idaho	27	(21 - 34)	4	(3 - 5)	10	(8 - 13)	14	(9 - 20)	23	(18 - 31)
Illinois	199	(169 - 233)	23	(18 - 30)	83	(70 - 97)	93	(70 - 123)	175	(147 - 209)
Indiana	110	(84 - 142)	13	(9 - 18)	40	(29 - 53)	57	(39 - 84)	97	(73 - 129)
Iowa	35	(26 - 48)	7	(5 - 10)	12	(8 - 17)	17	(10 - 27)	29	(20 - 41)
Kansas	37	(28 - 49)	5	(3 - 7)	15	(11 - 21)	18	(12 - 27)	33	(24 - 44)
Kentucky	71	(56 - 91)	8	(6 - 11)	27	(21 - 36)	36	(25 - 52)	63	(48 - 83)
Louisiana	73	(56 - 94)	8	(6 - 12)	28	(21 - 37)	37	(25 - 53)	64	(48 - 85)
Maine	23	(18 - 29)	2	(2 - 3)	10	(8 - 13)	11	(7 - 16)	21	(16 - 27)
Maryland	100	(76 - 131)	10	(7 - 14)	37	(27 - 50)	53	(36 - 78)	90	(67 - 121)
Massachusetts	133	(104 - 169)	15	(11 - 20)	53	(40 - 69)	66	(45 - 95)	118	(90 - 154)
Michigan	156	(134 - 183)	19	(15 - 24)	60	(51 - 70)	77	(59 - 101)	137	(115 - 164)
Minnesota	66	(50 - 86)	9	(6 - 13)	24	(17 - 33)	33	(22 - 49)	57	(42 - 77)
Mississippi	48	(38 - 61)	5	(4 - 7)	17	(13 - 23)	26	(18 - 37)	43	(33 - 56)
Missouri	93	(73 - 119)	11	(8 - 16)	36	(27 - 47)	46	(32 - 67)	82	(62 - 107)
Montana	16	(12 - 20)	2	(1 - 3)	6	(5 - 8)	8	(5 - 11)	14	(11 - 18)

(continued)

Table 19. *Illicit Drug Dependence in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)*

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	24	(18 - 31)	3	(2 - 5)	8	(6 - 11)	12	(8 - 18)	20	(15 - 28)
Nevada	40	(31 - 53)	6	(4 - 8)	12	(9 - 17)	22	(15 - 33)	35	(25 - 47)
New Hampshire	26	(21 - 34)	3	(2 - 5)	10	(8 - 13)	13	(9 - 19)	23	(17 - 30)
New Jersey	130	(100 - 169)	14	(10 - 20)	48	(36 - 65)	67	(46 - 99)	115	(87 - 154)
New Mexico	32	(25 - 41)	6	(4 - 8)	13	(10 - 18)	13	(9 - 20)	27	(20 - 36)
New York	364	(309 - 427)	37	(29 - 46)	147	(126 - 173)	179	(136 - 235)	327	(274 - 390)
North Carolina	149	(116 - 192)	17	(12 - 23)	56	(42 - 73)	77	(52 - 112)	132	(100 - 174)
North Dakota	8	(6 - 11)	1	(1 - 2)	3	(2 - 5)	3	(2 - 6)	7	(5 - 10)
Ohio	196	(167 - 229)	22	(18 - 27)	68	(59 - 80)	105	(81 - 135)	173	(145 - 207)
Oklahoma	59	(46 - 77)	7	(5 - 10)	21	(16 - 29)	31	(21 - 45)	52	(39 - 69)
Oregon	75	(58 - 96)	9	(6 - 12)	27	(21 - 36)	39	(26 - 56)	66	(50 - 87)
Pennsylvania	220	(186 - 260)	25	(20 - 31)	84	(71 - 100)	111	(84 - 146)	195	(162 - 235)
Rhode Island	22	(17 - 28)	3	(2 - 4)	9	(7 - 12)	10	(7 - 15)	19	(14 - 25)
South Carolina	66	(50 - 86)	7	(5 - 10)	22	(16 - 30)	37	(25 - 55)	59	(44 - 80)
South Dakota	11	(8 - 14)	2	(1 - 3)	4	(3 - 5)	5	(3 - 8)	9	(6 - 13)
Tennessee	91	(70 - 118)	10	(7 - 14)	29	(21 - 39)	52	(36 - 76)	81	(61 - 108)
Texas	349	(296 - 411)	51	(41 - 63)	129	(108 - 155)	169	(128 - 223)	298	(248 - 359)
Utah	43	(34 - 54)	6	(4 - 9)	18	(14 - 24)	19	(13 - 27)	37	(28 - 48)
Vermont	12	(10 - 16)	1	(1 - 2)	6	(4 - 7)	5	(4 - 8)	11	(9 - 14)
Virginia	125	(98 - 160)	15	(11 - 21)	46	(34 - 61)	64	(44 - 93)	110	(84 - 145)
Washington	113	(89 - 143)	13	(10 - 18)	46	(35 - 60)	53	(37 - 78)	99	(77 - 129)
West Virginia	34	(26 - 43)	3	(2 - 4)	14	(11 - 18)	17	(11 - 25)	31	(23 - 40)
Wisconsin	87	(67 - 113)	12	(9 - 17)	29	(21 - 39)	46	(31 - 67)	75	(55 - 101)
Wyoming	7	(5 - 9)	1	(1 - 2)	2	(2 - 3)	3	(2 - 5)	6	(4 - 8)

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 20. Dependence or Abuse of Illicit Drugs or Alcohol in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	22,428	(21,791 - 23,081)	1,762	(1,683 - 1,845)	6,778	(6,593 - 6,967)	13,887	(13,299 - 14,499)	20,666	(20,040 - 21,307)
Northeast	4,215	(4,007 - 4,434)	315	(291 - 341)	1,308	(1,243 - 1,375)	2,591	(2,396 - 2,801)	3,899	(3,697 - 4,113)
Midwest	4,964	(4,753 - 5,182)	384	(361 - 409)	1,526	(1,460 - 1,594)	3,054	(2,855 - 3,265)	4,579	(4,372 - 4,794)
South	7,686	(7,354 - 8,033)	589	(551 - 629)	2,280	(2,177 - 2,388)	4,818	(4,504 - 5,151)	7,098	(6,773 - 7,437)
West	5,563	(5,267 - 5,872)	474	(436 - 515)	1,664	(1,574 - 1,758)	3,425	(3,153 - 3,718)	5,089	(4,799 - 5,392)
Alabama	322	(275 - 377)	25	(20 - 31)	91	(78 - 106)	206	(166 - 256)	297	(252 - 350)
Alaska	51	(43 - 59)	4	(3 - 5)	16	(13 - 18)	31	(25 - 38)	47	(40 - 55)
Arizona	502	(428 - 586)	41	(32 - 51)	151	(131 - 174)	310	(248 - 387)	462	(391 - 543)
Arkansas	186	(158 - 219)	15	(12 - 19)	53	(45 - 61)	118	(94 - 147)	171	(143 - 203)
California	2,974	(2,727 - 3,239)	258	(226 - 293)	878	(805 - 956)	1,838	(1,616 - 2,086)	2,716	(2,475 - 2,976)
Colorado	421	(361 - 490)	36	(29 - 45)	130	(115 - 148)	255	(204 - 316)	385	(327 - 452)
Connecticut	276	(235 - 322)	20	(16 - 26)	85	(75 - 97)	171	(136 - 213)	256	(217 - 300)
Delaware	65	(56 - 77)	5	(4 - 7)	21	(18 - 24)	39	(31 - 50)	60	(51 - 71)
District of Columbia	62	(53 - 72)	2	(2 - 3)	18	(15 - 20)	42	(34 - 52)	60	(51 - 70)
Florida	1,293	(1,178 - 1,422)	93	(81 - 106)	326	(298 - 356)	874	(766 - 997)	1,201	(1,087 - 1,327)
Georgia	591	(502 - 695)	48	(38 - 61)	192	(167 - 221)	351	(275 - 446)	543	(458 - 644)
Hawaii	92	(78 - 109)	7	(6 - 10)	27	(23 - 31)	58	(46 - 73)	85	(71 - 101)
Idaho	120	(104 - 138)	11	(8 - 13)	37	(32 - 42)	73	(59 - 89)	110	(94 - 127)
Illinois	928	(848 - 1,015)	71	(61 - 82)	306	(282 - 332)	551	(480 - 632)	858	(780 - 943)
Indiana	467	(403 - 540)	34	(27 - 43)	140	(121 - 161)	292	(237 - 358)	432	(370 - 503)
Iowa	229	(197 - 264)	19	(15 - 23)	76	(67 - 87)	134	(108 - 166)	210	(179 - 244)
Kansas	207	(179 - 239)	17	(14 - 21)	69	(60 - 79)	121	(98 - 150)	190	(163 - 221)
Kentucky	273	(232 - 320)	22	(17 - 28)	81	(70 - 94)	170	(135 - 212)	251	(212 - 296)
Louisiana	301	(257 - 352)	21	(16 - 26)	100	(87 - 115)	180	(143 - 226)	280	(237 - 330)
Maine	90	(77 - 105)	6	(5 - 8)	27	(23 - 31)	57	(45 - 71)	84	(71 - 98)
Maryland	386	(325 - 458)	26	(20 - 33)	114	(98 - 133)	246	(194 - 310)	360	(301 - 430)
Massachusetts	600	(520 - 692)	42	(34 - 52)	192	(169 - 217)	367	(297 - 451)	559	(480 - 648)
Michigan	774	(709 - 844)	59	(52 - 67)	216	(200 - 235)	498	(438 - 565)	714	(651 - 784)
Minnesota	396	(339 - 459)	29	(23 - 36)	121	(106 - 138)	246	(198 - 304)	367	(312 - 428)
Mississippi	186	(158 - 218)	15	(12 - 19)	58	(50 - 67)	113	(89 - 142)	170	(144 - 201)
Missouri	394	(336 - 461)	33	(26 - 41)	119	(103 - 137)	243	(193 - 304)	361	(305 - 427)
Montana	87	(75 - 100)	7	(6 - 9)	29	(25 - 32)	51	(41 - 63)	80	(68 - 92)

(continued)

Table 20. Dependence or Abuse of Illicit Drugs or Alcohol in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	133	(114 - 154)	9	(7 - 12)	45	(39 - 51)	79	(63 - 98)	124	(106 - 144)
Nevada	212	(181 - 247)	17	(13 - 21)	55	(48 - 63)	140	(113 - 173)	195	(165 - 230)
New Hampshire	110	(95 - 128)	10	(8 - 13)	34	(30 - 38)	67	(53 - 83)	100	(85 - 118)
New Jersey	565	(482 - 661)	42	(33 - 54)	163	(140 - 188)	360	(287 - 450)	523	(443 - 617)
New Mexico	155	(134 - 180)	16	(13 - 19)	46	(40 - 52)	94	(75 - 116)	139	(119 - 163)
New York	1,498	(1,371 - 1,636)	117	(102 - 135)	480	(442 - 519)	901	(785 - 1,034)	1,381	(1,258 - 1,516)
North Carolina	598	(506 - 704)	43	(34 - 54)	177	(153 - 204)	378	(299 - 476)	555	(467 - 659)
North Dakota	51	(44 - 58)	3	(3 - 4)	20	(18 - 23)	27	(22 - 34)	47	(41 - 54)
Ohio	869	(795 - 950)	68	(60 - 77)	260	(241 - 280)	541	(472 - 619)	801	(729 - 881)
Oklahoma	285	(242 - 334)	22	(17 - 27)	86	(74 - 100)	177	(142 - 221)	263	(222 - 310)
Oregon	311	(266 - 363)	25	(21 - 31)	82	(72 - 94)	203	(163 - 251)	285	(242 - 336)
Pennsylvania	926	(838 - 1,023)	68	(59 - 78)	280	(256 - 306)	578	(500 - 668)	858	(772 - 953)
Rhode Island	96	(82 - 113)	6	(5 - 8)	30	(26 - 34)	60	(48 - 75)	90	(76 - 106)
South Carolina	354	(299 - 418)	22	(18 - 28)	90	(77 - 103)	242	(193 - 302)	332	(278 - 394)
South Dakota	68	(59 - 79)	6	(4 - 7)	21	(18 - 24)	42	(34 - 51)	63	(54 - 73)
Tennessee	421	(356 - 496)	30	(24 - 37)	118	(102 - 137)	273	(217 - 342)	391	(328 - 464)
Texas	1,625	(1,482 - 1,781)	147	(128 - 168)	520	(478 - 565)	958	(831 - 1,103)	1,478	(1,339 - 1,630)
Utah	161	(136 - 189)	16	(12 - 20)	62	(52 - 73)	83	(65 - 107)	145	(122 - 172)
Vermont	53	(46 - 61)	4	(3 - 5)	19	(16 - 21)	31	(25 - 39)	49	(42 - 58)
Virginia	620	(537 - 714)	46	(37 - 57)	200	(176 - 226)	374	(303 - 461)	574	(494 - 666)
Washington	438	(373 - 514)	33	(26 - 42)	139	(121 - 159)	266	(211 - 336)	405	(342 - 479)
West Virginia	118	(100 - 140)	8	(6 - 10)	35	(30 - 40)	76	(60 - 95)	110	(93 - 131)
Wisconsin	448	(386 - 520)	36	(29 - 45)	133	(115 - 151)	279	(225 - 345)	412	(352 - 482)
Wyoming	39	(34 - 45)	4	(3 - 4)	12	(11 - 14)	23	(18 - 29)	36	(30 - 42)

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 21. Needing But Not Receiving Treatment for Illicit Drug Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	6,399	(6,093 - 6,719)	1,031	(973 - 1,093)	2,415	(2,304 - 2,531)	2,952	(2,687 - 3,243)	5,367	(5,072 - 5,679)
Northeast	1,194	(1,104 - 1,292)	181	(163 - 200)	482	(446 - 522)	531	(457 - 617)	1,013	(927 - 1,108)
Midwest	1,322	(1,236 - 1,413)	220	(202 - 239)	512	(479 - 548)	590	(518 - 670)	1,102	(1,021 - 1,189)
South	2,233	(2,083 - 2,393)	354	(325 - 384)	803	(747 - 864)	1,076	(948 - 1,220)	1,879	(1,736 - 2,035)
West	1,650	(1,510 - 1,803)	277	(248 - 310)	617	(563 - 675)	756	(641 - 891)	1,373	(1,240 - 1,519)
Alabama	92	(73 - 115)	14	(10 - 18)	33	(26 - 43)	45	(32 - 64)	78	(61 - 101)
Alaska	15	(12 - 19)	2	(2 - 3)	5	(4 - 7)	7	(5 - 11)	13	(10 - 17)
Arizona	146	(116 - 182)	24	(18 - 32)	54	(42 - 68)	68	(48 - 97)	122	(94 - 156)
Arkansas	54	(43 - 67)	9	(7 - 12)	18	(14 - 23)	26	(19 - 37)	45	(35 - 58)
California	890	(779 - 1,017)	151	(128 - 179)	325	(282 - 374)	414	(325 - 526)	739	(633 - 861)
Colorado	119	(97 - 146)	22	(16 - 29)	49	(39 - 62)	48	(33 - 68)	97	(77 - 122)
Connecticut	73	(58 - 90)	12	(9 - 15)	30	(23 - 38)	31	(22 - 45)	61	(48 - 78)
Delaware	18	(14 - 22)	3	(2 - 4)	7	(5 - 9)	8	(5 - 12)	15	(12 - 19)
District of Columbia	14	(11 - 18)	1	(1 - 2)	6	(4 - 7)	8	(5 - 11)	13	(10 - 17)
Florida	384	(332 - 444)	54	(45 - 65)	125	(107 - 145)	204	(161 - 259)	329	(280 - 389)
Georgia	161	(125 - 208)	29	(21 - 40)	55	(41 - 73)	77	(51 - 117)	132	(98 - 178)
Hawaii	25	(19 - 32)	3	(2 - 5)	9	(7 - 12)	12	(8 - 19)	22	(16 - 29)
Idaho	35	(28 - 44)	6	(4 - 8)	13	(10 - 17)	16	(12 - 23)	30	(23 - 38)
Illinois	259	(227 - 295)	43	(35 - 51)	110	(96 - 127)	106	(83 - 136)	216	(186 - 251)
Indiana	140	(113 - 174)	21	(16 - 28)	52	(41 - 66)	67	(47 - 95)	119	(94 - 152)
Iowa	50	(39 - 63)	11	(8 - 15)	18	(14 - 24)	21	(14 - 32)	39	(29 - 52)
Kansas	49	(39 - 61)	8	(6 - 11)	21	(16 - 28)	19	(13 - 28)	41	(32 - 52)
Kentucky	88	(70 - 109)	14	(11 - 19)	32	(25 - 41)	41	(29 - 58)	73	(57 - 93)
Louisiana	90	(71 - 112)	13	(10 - 17)	35	(27 - 45)	42	(29 - 60)	77	(60 - 99)
Maine	27	(21 - 34)	4	(3 - 5)	11	(9 - 14)	12	(8 - 18)	23	(18 - 30)
Maryland	120	(96 - 151)	17	(13 - 23)	48	(37 - 62)	55	(38 - 79)	103	(81 - 132)
Massachusetts	163	(132 - 201)	26	(20 - 34)	66	(53 - 83)	71	(50 - 102)	137	(109 - 173)
Michigan	199	(173 - 229)	33	(28 - 39)	72	(63 - 83)	93	(72 - 121)	165	(141 - 195)
Minnesota	100	(81 - 124)	16	(12 - 21)	40	(32 - 51)	44	(31 - 62)	84	(66 - 106)
Mississippi	65	(52 - 80)	9	(7 - 12)	24	(19 - 30)	32	(23 - 45)	56	(44 - 71)
Missouri	122	(98 - 152)	18	(14 - 24)	46	(36 - 58)	57	(40 - 82)	103	(81 - 132)
Montana	20	(16 - 24)	3	(3 - 4)	8	(6 - 10)	8	(6 - 12)	16	(13 - 21)

(continued)

Table 21. Needing But Not Receiving Treatment for Illicit Drug Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	29	(22 - 37)	5	(4 - 7)	11	(8 - 14)	13	(8 - 19)	23	(18 - 31)
Nevada	54	(42 - 68)	9	(6 - 12)	17	(13 - 21)	28	(20 - 41)	45	(35 - 59)
New Hampshire	33	(26 - 40)	6	(5 - 8)	13	(11 - 17)	13	(9 - 19)	26	(20 - 34)
New Jersey	155	(125 - 193)	26	(19 - 34)	60	(46 - 77)	70	(49 - 99)	129	(101 - 166)
New Mexico	48	(39 - 59)	10	(8 - 13)	18	(14 - 23)	20	(13 - 29)	38	(29 - 49)
New York	447	(391 - 510)	61	(51 - 74)	186	(163 - 211)	200	(156 - 257)	386	(333 - 447)
North Carolina	191	(152 - 240)	28	(21 - 37)	69	(55 - 87)	94	(66 - 135)	164	(127 - 211)
North Dakota	10	(8 - 13)	2	(1 - 2)	5	(3 - 6)	4	(2 - 6)	8	(6 - 11)
Ohio	238	(210 - 269)	39	(33 - 46)	90	(79 - 102)	109	(87 - 137)	199	(173 - 229)
Oklahoma	79	(63 - 99)	13	(10 - 17)	29	(22 - 37)	37	(26 - 52)	66	(51 - 85)
Oregon	91	(74 - 112)	14	(11 - 19)	32	(25 - 41)	44	(31 - 61)	76	(60 - 97)
Pennsylvania	255	(222 - 294)	41	(34 - 49)	97	(84 - 113)	117	(91 - 152)	215	(183 - 252)
Rhode Island	27	(22 - 33)	4	(3 - 5)	12	(10 - 15)	11	(7 - 15)	23	(18 - 29)
South Carolina	94	(75 - 119)	14	(10 - 18)	31	(24 - 40)	50	(35 - 71)	81	(62 - 104)
South Dakota	14	(11 - 17)	3	(2 - 4)	5	(4 - 7)	6	(4 - 9)	11	(8 - 14)
Tennessee	123	(98 - 155)	18	(14 - 23)	39	(30 - 51)	67	(47 - 94)	106	(81 - 137)
Texas	465	(405 - 533)	87	(72 - 104)	174	(149 - 203)	204	(159 - 262)	378	(322 - 444)
Utah	56	(45 - 68)	11	(8 - 15)	23	(18 - 30)	22	(15 - 31)	45	(35 - 57)
Vermont	15	(12 - 18)	2	(1 - 3)	7	(6 - 8)	6	(4 - 8)	13	(10 - 16)
Virginia	159	(129 - 196)	27	(20 - 35)	64	(50 - 81)	68	(48 - 97)	132	(104 - 168)
Washington	143	(116 - 176)	19	(14 - 26)	59	(47 - 74)	64	(45 - 90)	123	(98 - 155)
West Virginia	36	(28 - 45)	4	(3 - 6)	14	(11 - 18)	17	(12 - 25)	31	(24 - 40)
Wisconsin	113	(91 - 142)	22	(17 - 28)	41	(32 - 53)	51	(35 - 74)	92	(71 - 119)
Wyoming	9	(7 - 12)	2	(1 - 2)	4	(3 - 5)	4	(3 - 6)	8	(6 - 10)

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but not receiving treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 22. Needing But Not Receiving Treatment for Alcohol Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	17,376	(16,819 - 17,951)	1,080	(1,016 - 1,148)	5,163	(4,999 - 5,333)	11,133	(10,616 - 11,674)	16,297	(15,747 - 16,863)
Northeast	3,205	(3,018 - 3,404)	190	(172 - 209)	987	(934 - 1,043)	2,028	(1,859 - 2,213)	3,015	(2,833 - 3,210)
Midwest	3,856	(3,672 - 4,049)	235	(217 - 253)	1,180	(1,128 - 1,234)	2,442	(2,272 - 2,624)	3,622	(3,441 - 3,811)
South	5,975	(5,667 - 6,298)	367	(339 - 396)	1,716	(1,632 - 1,803)	3,892	(3,609 - 4,196)	5,608	(5,306 - 5,926)
West	4,340	(4,066 - 4,630)	289	(259 - 322)	1,280	(1,203 - 1,361)	2,771	(2,522 - 3,042)	4,051	(3,784 - 4,335)
Alabama	252	(209 - 303)	16	(13 - 21)	66	(55 - 79)	170	(134 - 216)	236	(195 - 286)
Alaska	40	(33 - 47)	3	(2 - 3)	12	(10 - 14)	25	(19 - 32)	37	(31 - 45)
Arizona	383	(318 - 459)	25	(19 - 32)	116	(99 - 136)	241	(188 - 309)	358	(295 - 431)
Arkansas	145	(122 - 174)	10	(8 - 13)	41	(34 - 49)	94	(74 - 120)	135	(112 - 163)
California	2,343	(2,116 - 2,591)	160	(135 - 189)	675	(611 - 745)	1,508	(1,306 - 1,738)	2,183	(1,962 - 2,426)
Colorado	337	(283 - 401)	20	(16 - 26)	106	(91 - 123)	211	(165 - 269)	317	(264 - 379)
Connecticut	221	(185 - 264)	13	(10 - 17)	67	(57 - 78)	141	(110 - 180)	208	(172 - 250)
Delaware	49	(41 - 59)	3	(2 - 4)	15	(12 - 17)	31	(24 - 40)	46	(38 - 56)
District of Columbia	47	(40 - 56)	1	(1 - 2)	14	(12 - 17)	32	(25 - 40)	46	(39 - 55)
Florida	991	(884 - 1,110)	59	(50 - 69)	243	(217 - 271)	689	(592 - 801)	932	(828 - 1,049)
Georgia	464	(384 - 560)	29	(22 - 38)	142	(119 - 168)	294	(227 - 379)	435	(358 - 528)
Hawaii	78	(64 - 96)	5	(4 - 6)	22	(19 - 27)	51	(39 - 67)	74	(59 - 91)
Idaho	87	(74 - 103)	6	(5 - 8)	27	(23 - 32)	53	(42 - 67)	81	(68 - 96)
Illinois	731	(661 - 809)	42	(35 - 50)	236	(215 - 260)	453	(390 - 526)	690	(621 - 766)
Indiana	352	(297 - 416)	22	(17 - 27)	105	(88 - 124)	225	(179 - 282)	331	(277 - 393)
Iowa	196	(166 - 231)	12	(10 - 16)	66	(56 - 76)	119	(93 - 150)	184	(154 - 218)
Kansas	169	(143 - 199)	11	(9 - 14)	55	(47 - 64)	103	(81 - 129)	158	(133 - 187)
Kentucky	208	(173 - 250)	14	(10 - 18)	59	(49 - 70)	136	(106 - 173)	195	(160 - 235)
Louisiana	233	(195 - 278)	12	(9 - 16)	73	(62 - 85)	148	(116 - 188)	221	(184 - 264)
Maine	70	(59 - 83)	4	(3 - 5)	20	(17 - 24)	46	(36 - 58)	66	(55 - 79)
Maryland	282	(231 - 344)	15	(11 - 20)	82	(68 - 99)	185	(142 - 240)	267	(217 - 327)
Massachusetts	443	(376 - 521)	25	(19 - 32)	146	(126 - 169)	272	(216 - 342)	419	(354 - 494)
Michigan	582	(523 - 646)	35	(30 - 41)	161	(147 - 177)	385	(332 - 447)	547	(490 - 610)
Minnesota	319	(270 - 377)	19	(15 - 25)	98	(85 - 114)	202	(159 - 254)	300	(251 - 356)
Mississippi	147	(123 - 175)	9	(7 - 12)	43	(37 - 51)	94	(74 - 120)	138	(114 - 165)
Missouri	309	(254 - 375)	21	(17 - 27)	89	(74 - 105)	199	(153 - 258)	288	(235 - 352)
Montana	71	(61 - 83)	5	(4 - 6)	24	(21 - 27)	43	(34 - 54)	67	(57 - 78)

(continued)

Table 22. Needing But Not Receiving Treatment for Alcohol Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	108	(92 - 127)	6	(5 - 8)	38	(33 - 44)	64	(51 - 80)	102	(86 - 120)
Nevada	172	(143 - 206)	11	(8 - 14)	45	(38 - 52)	116	(92 - 147)	161	(133 - 194)
New Hampshire	81	(68 - 95)	5	(4 - 7)	25	(21 - 29)	50	(39 - 64)	75	(63 - 90)
New Jersey	440	(364 - 533)	27	(21 - 35)	121	(101 - 144)	293	(228 - 375)	414	(339 - 504)
New Mexico	117	(98 - 138)	8	(7 - 11)	35	(30 - 41)	73	(58 - 92)	108	(91 - 129)
New York	1,146	(1,032 - 1,271)	71	(60 - 84)	357	(325 - 392)	718	(616 - 834)	1,075	(964 - 1,198)
North Carolina	459	(379 - 556)	26	(20 - 34)	132	(110 - 156)	301	(234 - 387)	433	(355 - 527)
North Dakota	43	(37 - 49)	2	(2 - 3)	18	(15 - 20)	23	(18 - 29)	40	(35 - 47)
Ohio	657	(592 - 728)	40	(35 - 47)	195	(179 - 212)	422	(363 - 489)	616	(553 - 687)
Oklahoma	213	(177 - 256)	13	(10 - 17)	65	(55 - 77)	135	(105 - 173)	200	(165 - 241)
Oregon	229	(192 - 272)	14	(11 - 17)	60	(51 - 71)	155	(123 - 194)	215	(180 - 258)
Pennsylvania	689	(614 - 774)	39	(33 - 47)	215	(194 - 237)	435	(369 - 513)	650	(576 - 733)
Rhode Island	74	(61 - 89)	4	(3 - 5)	22	(18 - 26)	48	(37 - 62)	70	(58 - 85)
South Carolina	287	(235 - 350)	14	(10 - 18)	67	(56 - 80)	206	(160 - 264)	273	(223 - 335)
South Dakota	56	(48 - 66)	4	(3 - 5)	18	(15 - 21)	34	(27 - 43)	52	(44 - 62)
Tennessee	322	(266 - 387)	18	(14 - 24)	88	(73 - 104)	216	(168 - 276)	304	(250 - 368)
Texas	1,304	(1,172 - 1,450)	93	(78 - 109)	405	(368 - 445)	807	(690 - 941)	1,212	(1,083 - 1,353)
Utah	117	(95 - 143)	10	(7 - 13)	44	(35 - 53)	64	(47 - 87)	107	(86 - 133)
Vermont	41	(35 - 48)	2	(2 - 3)	14	(12 - 16)	25	(20 - 32)	39	(33 - 46)
Virginia	484	(410 - 571)	29	(22 - 37)	157	(135 - 181)	299	(237 - 375)	456	(384 - 539)
Washington	335	(277 - 404)	21	(16 - 27)	103	(87 - 122)	211	(163 - 272)	314	(258 - 381)
West Virginia	85	(70 - 104)	5	(4 - 7)	24	(20 - 29)	56	(42 - 72)	80	(65 - 98)
Wisconsin	334	(281 - 396)	20	(15 - 25)	101	(86 - 119)	213	(168 - 269)	314	(263 - 375)
Wyoming	32	(27 - 38)	2	(2 - 3)	11	(9 - 12)	19	(15 - 24)	30	(25 - 35)

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but not receiving treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 23. Serious Mental Illness in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Total U.S.	11,251	(10,767 - 11,759)	2,554	(2,446 - 2,668)	8,697	(8,226 - 9,196)
Northeast	2,007	(1,854 - 2,173)	468	(436 - 503)	1,539	(1,395 - 1,697)
Midwest	2,592	(2,433 - 2,760)	587	(554 - 622)	2,005	(1,852 - 2,169)
South	4,116	(3,858 - 4,393)	855	(805 - 907)	3,262	(3,011 - 3,533)
West	2,537	(2,340 - 2,749)	645	(595 - 698)	1,892	(1,705 - 2,099)
Alabama	205	(166 - 252)	36	(29 - 44)	169	(132 - 216)
Alaska	25	(21 - 31)	6	(5 - 8)	19	(15 - 24)
Arizona	251	(204 - 308)	54	(45 - 66)	197	(153 - 253)
Arkansas	134	(110 - 163)	26	(21 - 31)	108	(85 - 137)
California	1,183	(1,028 - 1,359)	340	(298 - 387)	843	(698 - 1,016)
Colorado	192	(157 - 235)	44	(36 - 53)	148	(116 - 190)
Connecticut	130	(105 - 159)	32	(26 - 39)	98	(75 - 126)
Delaware	33	(27 - 40)	7	(6 - 9)	25	(20 - 32)
District of Columbia	22	(18 - 27)	6	(5 - 7)	17	(13 - 22)
Florida	703	(609 - 811)	125	(108 - 143)	579	(488 - 685)
Georgia	324	(258 - 404)	63	(50 - 79)	260	(199 - 340)
Hawaii	42	(33 - 53)	9	(7 - 12)	33	(24 - 43)
Idaho	67	(56 - 81)	16	(13 - 20)	51	(40 - 65)
Illinois	442	(388 - 503)	115	(101 - 129)	327	(277 - 387)
Indiana	265	(219 - 321)	59	(49 - 71)	206	(163 - 261)
Iowa	126	(102 - 154)	28	(23 - 34)	98	(76 - 126)
Kansas	95	(77 - 116)	24	(19 - 30)	71	(55 - 91)
Kentucky	182	(149 - 222)	32	(26 - 40)	150	(118 - 190)
Louisiana	164	(134 - 201)	36	(30 - 45)	128	(100 - 163)
Maine	54	(44 - 66)	11	(9 - 14)	43	(34 - 55)
Maryland	184	(148 - 228)	44	(35 - 55)	140	(107 - 183)
Massachusetts	260	(213 - 316)	62	(51 - 75)	198	(155 - 253)
Michigan	391	(343 - 445)	91	(81 - 102)	300	(254 - 354)
Minnesota	204	(168 - 248)	47	(38 - 57)	157	(124 - 200)
Mississippi	110	(90 - 134)	22	(18 - 27)	88	(69 - 112)
Missouri	255	(209 - 311)	50	(41 - 61)	205	(162 - 260)
Montana	42	(34 - 52)	9	(7 - 11)	33	(26 - 43)

(continued)

Table 23. Serious Mental Illness in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Nebraska	64	(52 - 78)	16	(13 - 20)	48	(37 - 61)
Nevada	102	(83 - 126)	20	(16 - 24)	83	(64 - 106)
New Hampshire	55	(45 - 67)	13	(11 - 16)	42	(33 - 54)
New Jersey	283	(230 - 348)	64	(52 - 79)	219	(170 - 282)
New Mexico	70	(57 - 85)	17	(14 - 21)	52	(41 - 67)
New York	713	(622 - 817)	170	(150 - 192)	543	(458 - 644)
North Carolina	318	(256 - 393)	63	(51 - 78)	254	(196 - 329)
North Dakota	22	(18 - 27)	7	(6 - 8)	15	(12 - 20)
Ohio	483	(425 - 548)	95	(84 - 107)	388	(332 - 453)
Oklahoma	141	(115 - 172)	34	(27 - 41)	107	(84 - 137)
Oregon	164	(136 - 199)	33	(27 - 41)	131	(104 - 165)
Pennsylvania	438	(380 - 505)	100	(87 - 115)	338	(283 - 404)
Rhode Island	49	(40 - 60)	11	(9 - 13)	38	(30 - 49)
South Carolina	165	(134 - 204)	32	(26 - 39)	134	(104 - 171)
South Dakota	28	(22 - 35)	6	(5 - 8)	21	(16 - 28)
Tennessee	272	(221 - 333)	47	(38 - 58)	225	(176 - 285)
Texas	794	(694 - 908)	207	(183 - 233)	588	(492 - 701)
Utah	117	(98 - 140)	36	(30 - 43)	81	(64 - 103)
Vermont	25	(21 - 30)	6	(5 - 8)	19	(15 - 24)
Virginia	283	(230 - 348)	62	(50 - 75)	221	(171 - 286)
Washington	259	(214 - 313)	54	(45 - 66)	205	(162 - 258)
West Virginia	83	(67 - 102)	14	(11 - 17)	69	(54 - 88)
Wisconsin	217	(178 - 263)	50	(41 - 61)	166	(130 - 212)
Wyoming	21	(17 - 25)	5	(4 - 6)	16	(12 - 20)

NOTE: Serious Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment in carrying out major life activities. For details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 24. Any Mental Illness in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Total U.S.	45,705	(44,761 - 46,669)	10,212	(10,012 - 10,413)	35,493	(34,578 - 36,433)
Northeast	8,472	(8,132 - 8,823)	1,894	(1,828 - 1,961)	6,578	(6,247 - 6,922)
Midwest	10,221	(9,891 - 10,557)	2,304	(2,242 - 2,366)	7,917	(7,597 - 8,247)
South	16,372	(15,843 - 16,920)	3,530	(3,429 - 3,632)	12,842	(12,328 - 13,379)
West	10,639	(10,193 - 11,102)	2,484	(2,388 - 2,582)	8,155	(7,723 - 8,607)
Alabama	816	(722 - 918)	157	(140 - 174)	660	(569 - 760)
Alaska	105	(93 - 118)	25	(22 - 27)	80	(69 - 93)
Arizona	964	(847 - 1,092)	217	(195 - 240)	747	(635 - 873)
Arkansas	454	(404 - 508)	92	(84 - 102)	361	(314 - 414)
California	5,161	(4,800 - 5,544)	1,289	(1,211 - 1,371)	3,872	(3,526 - 4,245)
Colorado	746	(660 - 841)	168	(151 - 186)	578	(496 - 670)
Connecticut	513	(452 - 580)	118	(107 - 131)	395	(337 - 460)
Delaware	131	(116 - 147)	29	(26 - 32)	102	(88 - 118)
District of Columbia	103	(92 - 115)	26	(23 - 28)	78	(67 - 89)
Florida	2,593	(2,408 - 2,791)	505	(472 - 539)	2,089	(1,908 - 2,283)
Georgia	1,269	(1,121 - 1,433)	302	(271 - 336)	967	(825 - 1,127)
Hawaii	188	(166 - 213)	40	(35 - 45)	148	(127 - 173)
Idaho	278	(248 - 309)	58	(53 - 64)	220	(192 - 251)
Illinois	1,827	(1,705 - 1,956)	453	(427 - 480)	1,374	(1,256 - 1,501)
Indiana	1,011	(903 - 1,127)	224	(203 - 246)	787	(683 - 901)
Iowa	469	(417 - 525)	110	(99 - 121)	359	(311 - 414)
Kansas	384	(337 - 434)	93	(82 - 104)	291	(248 - 340)
Kentucky	684	(607 - 766)	131	(117 - 145)	553	(480 - 634)
Louisiana	682	(609 - 762)	158	(142 - 174)	524	(455 - 601)
Maine	206	(183 - 232)	43	(39 - 47)	164	(141 - 189)
Maryland	824	(722 - 938)	184	(164 - 206)	640	(543 - 750)
Massachusetts	1,049	(938 - 1,169)	255	(231 - 280)	794	(689 - 911)
Michigan	1,620	(1,513 - 1,734)	342	(323 - 362)	1,278	(1,174 - 1,388)
Minnesota	774	(686 - 870)	183	(165 - 201)	592	(508 - 686)
Mississippi	433	(385 - 485)	99	(89 - 109)	334	(288 - 385)
Missouri	931	(827 - 1,044)	204	(184 - 225)	727	(628 - 837)
Montana	166	(147 - 185)	35	(31 - 38)	131	(113 - 150)

(continued)

Table 24. Any Mental Illness in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Nebraska	262	(232 - 296)	64	(58 - 70)	199	(170 - 231)
Nevada	425	(375 - 480)	74	(66 - 82)	352	(303 - 406)
New Hampshire	214	(190 - 240)	46	(42 - 51)	168	(145 - 193)
New Jersey	1,274	(1,126 - 1,436)	269	(242 - 298)	1,004	(864 - 1,163)
New Mexico	272	(238 - 309)	65	(58 - 72)	207	(175 - 243)
New York	3,041	(2,831 - 3,264)	679	(637 - 722)	2,363	(2,159 - 2,581)
North Carolina	1,325	(1,172 - 1,494)	271	(243 - 301)	1,054	(908 - 1,218)
North Dakota	93	(82 - 105)	26	(23 - 29)	67	(57 - 78)
Ohio	1,878	(1,753 - 2,009)	381	(361 - 402)	1,497	(1,377 - 1,625)
Oklahoma	587	(522 - 658)	138	(124 - 152)	449	(388 - 517)
Oregon	625	(555 - 702)	126	(114 - 140)	499	(432 - 573)
Pennsylvania	1,882	(1,746 - 2,027)	419	(393 - 447)	1,463	(1,331 - 1,605)
Rhode Island	188	(167 - 212)	41	(37 - 45)	147	(127 - 170)
South Carolina	649	(573 - 733)	134	(120 - 149)	515	(443 - 597)
South Dakota	113	(99 - 127)	27	(24 - 31)	85	(73 - 100)
Tennessee	1,084	(959 - 1,218)	204	(184 - 225)	880	(761 - 1,012)
Texas	3,243	(3,016 - 3,483)	784	(736 - 833)	2,459	(2,243 - 2,692)
Utah	484	(438 - 533)	139	(126 - 153)	345	(303 - 391)
Vermont	104	(93 - 116)	24	(22 - 26)	80	(69 - 92)
Virginia	1,175	(1,045 - 1,318)	262	(236 - 289)	914	(790 - 1,053)
Washington	1,138	(1,017 - 1,269)	228	(205 - 251)	910	(795 - 1,037)
West Virginia	318	(283 - 357)	55	(50 - 61)	263	(229 - 301)
Wisconsin	858	(764 - 962)	196	(177 - 217)	662	(572 - 763)
Wyoming	87	(78 - 97)	21	(19 - 22)	67	(58 - 76)

NOTE: Any Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), regardless of the level of impairment in carrying out major life activities. For details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 25. Had Serious Thoughts of Suicide in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Total U.S.	8,620	(8,195 - 9,068)	2,150	(2,056 - 2,248)	6,470	(6,061 - 6,908)
Northeast	1,610	(1,470 - 1,762)	403	(369 - 439)	1,207	(1,080 - 1,349)
Midwest	2,019	(1,872 - 2,177)	499	(466 - 535)	1,520	(1,384 - 1,669)
South	2,948	(2,712 - 3,206)	728	(676 - 785)	2,220	(2,003 - 2,461)
West	2,043	(1,853 - 2,252)	520	(474 - 570)	1,523	(1,347 - 1,721)
Alabama	144	(112 - 185)	34	(27 - 43)	110	(81 - 149)
Alaska	20	(15 - 25)	5	(4 - 6)	15	(11 - 20)
Arizona	196	(152 - 253)	45	(36 - 56)	152	(111 - 205)
Arkansas	93	(73 - 119)	20	(16 - 25)	73	(55 - 97)
California	968	(831 - 1,127)	271	(235 - 312)	697	(572 - 849)
Colorado	154	(120 - 197)	36	(29 - 45)	118	(88 - 159)
Connecticut	97	(75 - 123)	22	(17 - 28)	74	(55 - 100)
Delaware	26	(20 - 33)	7	(6 - 9)	19	(14 - 26)
District of Columbia	19	(15 - 24)	5	(4 - 6)	14	(11 - 19)
Florida	500	(423 - 591)	105	(90 - 122)	395	(323 - 483)
Georgia	221	(173 - 282)	59	(46 - 74)	162	(120 - 220)
Hawaii	38	(29 - 48)	10	(8 - 12)	28	(20 - 38)
Idaho	54	(42 - 69)	13	(11 - 16)	41	(30 - 55)
Illinois	387	(333 - 450)	100	(87 - 115)	287	(237 - 347)
Indiana	189	(151 - 237)	45	(36 - 56)	144	(109 - 190)
Iowa	88	(70 - 111)	22	(17 - 27)	67	(50 - 88)
Kansas	73	(58 - 92)	21	(16 - 26)	53	(39 - 70)
Kentucky	129	(102 - 162)	28	(22 - 35)	101	(76 - 133)
Louisiana	116	(91 - 147)	33	(26 - 41)	83	(62 - 112)
Maine	47	(37 - 60)	8	(7 - 10)	39	(29 - 51)
Maryland	150	(117 - 192)	39	(30 - 49)	112	(82 - 151)
Massachusetts	200	(160 - 251)	48	(38 - 60)	152	(116 - 200)
Michigan	311	(267 - 362)	76	(67 - 86)	236	(194 - 285)
Minnesota	154	(123 - 193)	39	(32 - 49)	115	(87 - 152)
Mississippi	79	(63 - 99)	21	(17 - 27)	58	(43 - 77)
Missouri	192	(150 - 244)	42	(34 - 52)	150	(111 - 200)
Montana	34	(27 - 43)	8	(7 - 10)	25	(19 - 34)

(continued)

Table 25. Had Serious Thoughts of Suicide in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Nebraska	53	(42 - 67)	13	(11 - 17)	40	(30 - 52)
Nevada	81	(63 - 104)	15	(12 - 18)	66	(49 - 89)
New Hampshire	44	(34 - 56)	10	(8 - 12)	34	(25 - 46)
New Jersey	238	(188 - 300)	63	(50 - 79)	175	(130 - 234)
New Mexico	52	(41 - 66)	15	(12 - 19)	37	(27 - 50)
New York	571	(487 - 668)	147	(127 - 170)	424	(347 - 517)
North Carolina	241	(188 - 308)	57	(45 - 72)	184	(136 - 248)
North Dakota	17	(13 - 22)	5	(4 - 7)	12	(9 - 16)
Ohio	358	(306 - 418)	85	(74 - 97)	273	(224 - 331)
Oklahoma	104	(81 - 133)	25	(20 - 32)	79	(58 - 106)
Oregon	121	(95 - 154)	25	(20 - 32)	96	(72 - 128)
Pennsylvania	355	(302 - 416)	90	(77 - 104)	265	(216 - 324)
Rhode Island	37	(29 - 47)	9	(7 - 11)	28	(21 - 38)
South Carolina	122	(95 - 156)	28	(22 - 35)	94	(70 - 127)
South Dakota	24	(19 - 30)	6	(5 - 7)	18	(13 - 24)
Tennessee	172	(133 - 220)	36	(29 - 45)	135	(100 - 182)
Texas	550	(468 - 647)	166	(144 - 191)	385	(311 - 475)
Utah	95	(76 - 119)	29	(23 - 36)	66	(49 - 88)
Vermont	22	(17 - 27)	5	(4 - 7)	16	(12 - 22)
Virginia	220	(174 - 277)	54	(43 - 67)	166	(124 - 220)
Washington	214	(169 - 269)	44	(35 - 55)	170	(128 - 224)
West Virginia	63	(49 - 81)	12	(10 - 15)	50	(37 - 68)
Wisconsin	173	(138 - 217)	45	(36 - 56)	128	(96 - 169)
Wyoming	16	(13 - 20)	4	(3 - 5)	12	(9 - 16)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

Table 26. Had at Least One Major Depressive Episode in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs

State	18+ (Estimate)	18+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Total U.S.	15,303	(14,725 - 15,903)	1,975	(1,895 - 2,059)	2,758	(2,643 - 2,877)	12,545	(11,975 - 13,141)
Northeast	2,820	(2,628 - 3,024)	333	(309 - 358)	501	(465 - 540)	2,318	(2,133 - 2,519)
Midwest	3,486	(3,299 - 3,683)	441	(417 - 466)	632	(596 - 669)	2,855	(2,674 - 3,046)
South	5,505	(5,209 - 5,818)	724	(685 - 766)	950	(896 - 1,007)	4,555	(4,267 - 4,862)
West	3,492	(3,240 - 3,762)	478	(442 - 517)	675	(624 - 730)	2,817	(2,573 - 3,082)
Alabama	262	(217 - 316)	29	(24 - 35)	41	(33 - 51)	221	(178 - 275)
Alaska	33	(28 - 40)	4	(4 - 5)	7	(5 - 8)	27	(21 - 34)
Arizona	330	(272 - 400)	45	(37 - 55)	60	(49 - 73)	270	(215 - 339)
Arkansas	167	(140 - 199)	20	(16 - 24)	26	(21 - 32)	141	(115 - 172)
California	1,659	(1,463 - 1,880)	244	(215 - 276)	347	(305 - 394)	1,312	(1,124 - 1,529)
Colorado	264	(219 - 316)	34	(28 - 41)	44	(36 - 54)	220	(177 - 272)
Connecticut	166	(137 - 201)	22	(18 - 26)	29	(23 - 36)	137	(109 - 172)
Delaware	46	(38 - 56)	6	(5 - 7)	10	(8 - 12)	36	(29 - 46)
District of Columbia	32	(26 - 38)	2	(2 - 3)	6	(5 - 8)	25	(20 - 32)
Florida	911	(807 - 1,027)	117	(104 - 132)	142	(124 - 162)	769	(668 - 884)
Georgia	441	(367 - 530)	66	(55 - 80)	82	(67 - 101)	359	(288 - 447)
Hawaii	57	(47 - 70)	7	(5 - 8)	10	(8 - 13)	47	(37 - 60)
Idaho	85	(71 - 100)	12	(10 - 14)	17	(14 - 21)	67	(55 - 82)
Illinois	651	(578 - 732)	79	(69 - 90)	130	(115 - 147)	521	(451 - 601)
Indiana	355	(296 - 425)	46	(38 - 55)	56	(45 - 68)	299	(243 - 368)
Iowa	158	(133 - 189)	18	(15 - 22)	27	(22 - 33)	131	(106 - 161)
Kansas	137	(116 - 163)	20	(16 - 23)	29	(24 - 35)	108	(88 - 133)
Kentucky	227	(188 - 272)	28	(23 - 33)	34	(28 - 42)	192	(155 - 237)
Louisiana	205	(168 - 249)	25	(21 - 31)	37	(30 - 46)	168	(133 - 210)
Maine	71	(59 - 85)	8	(7 - 10)	11	(9 - 14)	59	(48 - 73)
Maryland	260	(214 - 316)	34	(28 - 42)	47	(38 - 58)	213	(169 - 268)
Massachusetts	368	(308 - 438)	39	(32 - 47)	66	(54 - 81)	302	(245 - 370)
Michigan	517	(463 - 577)	69	(61 - 77)	101	(90 - 114)	416	(364 - 474)
Minnesota	271	(227 - 323)	32	(27 - 39)	50	(41 - 61)	221	(179 - 273)
Mississippi	135	(112 - 163)	19	(15 - 23)	22	(18 - 27)	114	(91 - 141)
Missouri	327	(273 - 392)	46	(38 - 55)	52	(42 - 63)	276	(224 - 339)
Montana	55	(46 - 65)	6	(5 - 7)	11	(9 - 13)	44	(36 - 54)

(continued)

Table 26. Had at Least One Major Depressive Episode in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2009 and 2010 NSDUHs (continued)

State	18+ (Estimate)	18+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Nebraska	94	(79 - 111)	12	(10 - 14)	17	(14 - 21)	77	(63 - 94)
Nevada	135	(112 - 163)	16	(13 - 20)	19	(15 - 23)	116	(94 - 144)
New Hampshire	74	(61 - 89)	9	(8 - 11)	13	(11 - 16)	61	(49 - 76)
New Jersey	400	(331 - 483)	50	(41 - 61)	75	(61 - 93)	325	(259 - 407)
New Mexico	87	(71 - 105)	14	(11 - 17)	18	(14 - 22)	69	(55 - 87)
New York	1,004	(888 - 1,135)	118	(105 - 134)	169	(148 - 192)	835	(723 - 963)
North Carolina	454	(376 - 548)	54	(44 - 65)	72	(58 - 89)	382	(307 - 474)
North Dakota	33	(28 - 39)	4	(3 - 5)	7	(6 - 9)	26	(21 - 31)
Ohio	605	(540 - 677)	76	(68 - 86)	100	(89 - 111)	505	(443 - 576)
Oklahoma	203	(169 - 242)	23	(19 - 28)	41	(33 - 50)	162	(130 - 200)
Oregon	220	(186 - 260)	26	(21 - 31)	39	(32 - 47)	181	(149 - 220)
Pennsylvania	637	(564 - 720)	76	(67 - 86)	118	(103 - 135)	519	(449 - 600)
Rhode Island	65	(54 - 78)	6	(5 - 8)	12	(10 - 15)	53	(42 - 65)
South Carolina	250	(207 - 301)	30	(25 - 36)	42	(34 - 52)	208	(167 - 257)
South Dakota	36	(29 - 44)	6	(5 - 7)	7	(5 - 9)	29	(23 - 36)
Tennessee	347	(288 - 418)	41	(34 - 49)	50	(41 - 62)	297	(239 - 367)
Texas	1,058	(933 - 1,198)	169	(149 - 191)	211	(185 - 240)	847	(727 - 985)
Utah	156	(133 - 182)	21	(17 - 25)	35	(29 - 42)	121	(99 - 147)
Vermont	34	(29 - 41)	4	(3 - 5)	7	(6 - 8)	28	(22 - 34)
Virginia	400	(336 - 475)	50	(42 - 60)	71	(58 - 87)	329	(267 - 403)
Washington	381	(320 - 453)	46	(38 - 55)	63	(52 - 77)	318	(259 - 388)
West Virginia	108	(90 - 129)	10	(9 - 13)	15	(12 - 18)	93	(76 - 114)
Wisconsin	302	(253 - 359)	34	(28 - 40)	56	(46 - 69)	246	(199 - 302)
Wyoming	29	(25 - 35)	4	(3 - 4)	6	(5 - 7)	24	(19 - 29)

NOTE: Major depressive episode is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. For details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.

NOTE: There are minor wording differences in the questions in the adult and adolescent major depressive episode modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to get an overall estimate (12 or older).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).