

Figure D1. Northeast Region Atlantic Cod – Georges Bank fishing mortality has not been controlled and biomass has not increased. Due to the periodic recalculation of  $F$  and  $B$  by stock assessment scientists, the initial estimates of  $F$  and  $B$  used in the overfished declaration are included to illustrate the uncertainty of stock assessment estimates.




Figure D2. Northeast Region Thorny Skate – Gulf of Maine fishing mortality has not been controlled and biomass has not increased as expected.  $B_{msy}$  proxy is in kg/tow. Overfishing occurs if there is greater than a 20% decrease in the 3-year moving average. Thus, a negative ratio or a ratio < 1 represents a stock that is not subject to overfishing.




Figure D3. Northeast Region White Hake – Gulf of Maine / Georges Bank fishing mortality has not been controlled and biomass has not increased.  $B_{msy}$  proxy is in kg/tow. Due to the periodic recalculation of F and B by stock assessment scientists, the initial estimates of F and B used in the overfished declaration are included to illustrate the uncertainty of stock assessment estimates.

## Winter Flounder - Southern New England/ Mid-Atlantic




Figure D4. Northeast Region Winter Flounder – Southern New England / Mid-Atlantic fishing mortality has not been controlled and biomass is not increasing. Due to the periodic recalculation of F and B by stock assessment scientists, the initial estimates of F and B used in the overfished declaration are included to illustrate the uncertainty of stock assessment estimates.




Figure D5. Northeast Region Yellowtail Flounder – Cape Cod / Gulf of Maine fishing mortality has not been controlled and biomass has not increased. Due to the periodic recalculation of F and B by stock assessment scientists, the initial estimates of F and B used in the overfished declaration are included to illustrate the uncertainty of stock assessment estimates.

## Yellowtail Flounder - Southern New England / Mid-Atlantic




Figure D6. Northeast Region Yellowtail Flounder – Southern New England / Mid-Atlantic fishing mortality has not been controlled and biomass has not increased. Due to the periodic recalculation of F and B by stock assessment scientists, the initial estimates of F and B used in the overfished declaration are included to illustrate the uncertainty of stock assessment estimates.

# Bluefin Tuna - Western Atlantic




Figure D7. Highly Migratory Species Bluefin Tuna - Western Atlantic fishing mortality is not controlled and biomass (SSB) is not increasing. Results from the LRS and HRS are not combined, but are presented separately.

\*FMP has not been internationally Implemented.