




Grain Inspection,
Packers and
Stockyards
Administration

Federal Grain
Inspection
Service

2010 U.S. Grain Exports: Quality Report



*27th
Edition*

Table of Contents

INTRODUCTION	1
WHEAT	2
WHEAT GRADES AND GRADE REQUIREMENTS	2
<i>Table 1. U.S. Wheat Exports: Number of lots and quantity exported by class and grade, 2008-2010.....</i>	<i>3</i>
<i>Table 2. Summary of export Hard Red Winter wheat quality, 2008-2010.....</i>	<i>6</i>
Figure 1. HRW Dockage Distribution, All Grades	8
Figure 2. HRW Protein Distribution, All Grades	9
<i>Table 3. Summary of export Hard Red Spring wheat quality, 2008-2010.....</i>	<i>10</i>
Figure 3. HRS Dockage Distribution, All Grades	12
Figure 4. HRS Protein Distribution, All Grades	13
<i>Table 4. Summary of export Soft Red Winter wheat quality, 2008-2010.....</i>	<i>14</i>
Figure 5. SRW Dockage Distribution, All Grades.....	16
Figure 6. SRW Protein Distribution, All Grades.....	17
<i>Table 5. Summary of export Durum wheat quality, 2008-2010.....</i>	<i>18</i>
Figure 7. DU Dockage Distribution, All Grades.....	20
Figure 8. DU Protein Distribution, All Grades.....	21
<i>Table 6. Summary of export Soft White wheat quality, 2008-2010.....</i>	<i>22</i>
Figure 9. SWH Dockage Distribution, All Grades	24
Figure 10. SWH Protein Distribution, All Grades	25
<i>Table 7. Summary of export Hard White wheat quality, 2008-2010.....</i>	<i>26</i>
<i>Table 8. Summary of export Mixed wheat quality, 2008-2010.....</i>	<i>28</i>
CORN	29
CORN GRADES AND GRADE REQUIREMENTS	29
<i>Table 9. U.S. Corn Exports: Number of lots and quantity exported by class and grade, 2008-2010.....</i>	<i>31</i>
<i>Table 10. Summary of export Yellow Corn quality, 2008-2010.....</i>	<i>32</i>
<i>Table 11. Summary of export White Corn quality, 2008-2010.....</i>	<i>34</i>
Figure 11. Corn BCFM Distribution, Grade 2	36
Figure 12. Corn BCFM Distribution, Grade 3	37
SOYBEANS.....	38
SOYBEAN GRADES AND GRADE REQUIREMENTS	38
<i>Table 12. U.S. Soybean Exports: Number of lots and quantity exported by grade, 2008-2010.....</i>	<i>39</i>
<i>Table 13. Summary of export Soybean quality, 2008-2010.....</i>	<i>40</i>
Figure 13. Soybean FM Distribution, Grade 2	42
Figure 14. Soybean Protein Distribution, All Grades	43
Figure 15. Soybean Oil Distribution, All Grades	44

OTHER GRAIN EXPORTS.....	45
SORGHUM.....	45
SORGHUM GRADES AND GRADE REQUIREMENTS.....	45
<i>Table 14. U.S. Sorghum Exports: Number of lots and quantity exported by class and grade, 2007-2009.....</i>	<i>47</i>
<i>Table 15. Summary of export Sorghum quality, 2008-2010.....</i>	<i>48</i>
Figure 16. Sorghum BNFM Distribution, Grade 2.....	50
Figure 17. Sorghum FM Distribution, Grade 2.....	51
BARLEY	52
BARLEY GRADES AND GRADE REQUIREMENTS	52
<i>Table 16. U.S. Barley Exports: Number of lots and quantity exported by grade, 2008-2010.....</i>	<i>54</i>
<i>Table 17. Summary of export Barley quality, 2008-2010.....</i>	<i>55</i>
CANOLA.....	56
CANOLA GRADES AND GRADE REQUIREMENTS.....	56
<i>Table 18. U.S. Canola Exports: Number of lots and quantity exported by grade, 2007-2009.....</i>	<i>57</i>
<i>Table 19. Summary of export Canola quality, 2008-2010.....</i>	<i>58</i>
FLAXSEED.....	59
FLAXSEED GRADES AND GRADE REQUIREMENTS.....	59
<i>Table 20. U.S. Flaxseed Exports: Number of lots and quantity exported by grade, 2008-2010.....</i>	<i>60</i>
<i>Table 21. Summary of export Flaxseed quality, 2008-2010.....</i>	<i>60</i>
APPENDIX	61

Introduction

The *2010 U.S. Grain Exports: Quality Report* is produced by the Federal Grain Inspection Service (FGIS) of the U.S. Department of Agriculture's Grain Inspection, Packers and Stockyards Administration. The report is the result of FGIS' efforts to determine, document, and disseminate critical information regarding U.S. export grain quality.

The 2010 report is the twenty-seventh edition of this annual summary of export grain quality. The report summarizes the quality of export wheat, corn, soybeans, sorghum, barley, canola, and flaxseed. Mixed grain, rye, and sunflower seed are not included in this year's report; no lots have been reported in the past 4 years.

Organization of the Report

The report contains chapters addressing export wheat, export corn, export soybeans, and other grains. Each chapter contains:

- Standards and definitions for each grain,
- Tables that clearly illustrate all factor result averages at each applicable U.S. grade level, and
- Factor quality distribution graphs for selected factors.

In addition, an appendix contains figures illustrating select quantity and quality trends over time.

Methodology

FGIS collects and documents information about export grain shipments in the automated Export Grain Information System (EGIS). This system contains one record for each export lot inspected and/or weighed. In the case of some railcar exports, each record may contain information from several lots which were aggregated to simplify internal reporting. For the purposes of this export quality report, only information from waterborne export shipments was used. Waterborne export shipments represented 90 percent of the total export lots in the EGIS database for 2010. Generally, each EGIS record contains the quantity of the lot and the average factor results certified for the lot. The tables in this report contain descriptive statistics which summarize these lot quantities and the weighted averages. Where appropriate, tables are provided which show the number of lots and the quantity of grain which was used to generate the descriptive statistics. Many of the tables summarize factor averages by grade.

A U.S. grade is determined by analyzing the physical and biological factors present in the sample. Limits for the grading factors are established for each numerical grade. Grades range from U.S. No. 1 (highest) to U.S. Sample grade (lowest). When a particular grade is cited in this report, it includes lots certified at that grade plus lots certified with the "or better" designation. For example, U.S. No. 2 grade includes lots which were certificated as "U.S. No. 2" and lots certificated as "U.S. No. 2 or better." Factors that exceed the established limits, except for test weight, lower the grade. The established limits for test weight represent minimum requirements for each grade.

This report does not contain data on the volume of export grain in bushels. Listed to the right are the equations for converting the approximate quantity of grain from metric tons to bushels.

Conversion Equation	
Bushels =	$\frac{\text{Metric Tons} \times 2204.622 \text{ Pounds}}{\text{Legal Test Weight/Bushel of Grain}}$
Legal Test Weights Per Bushel for Specific Grains	
Wheat =	60 pounds/bushel
Corn =	56 pounds/bushel
Soybeans =	60 pounds/bushel
Canola =	50 pounds/bushel
Sorghum =	56 pounds/bushel
Barley =	48 pounds/bushel
Sunflower Seed =	28 pounds/bushel
Rye =	56 pounds/bushel
Oats =	32 pounds/bushel

Wheat

Wheat Grades and Grade Requirements

Wheat is divided into eight classes: Hard Red Spring wheat, Hard Red Winter wheat, Soft Red Winter wheat, Durum wheat, Hard White wheat, Soft White wheat, Unclassed wheat, and Mixed wheat. The classes Hard Red Spring wheat, Soft White wheat, and Durum wheat are further divided into subclasses. There are no subclasses in the classes Hard Red Winter wheat, Soft Red Winter wheat, Hard White wheat, Unclassed wheat, and Mixed wheat. Each class and subclass is divided into five U.S. numerical grades and U.S. Sample grade. Special grades are provided to emphasize special qualities or conditions affecting the value of wheat. Special grades are added to and made a part of the grade designation. They do not affect the numerical or Sample grade designation.

U.S. Standards for Wheat

Grade	Minimum Limits of:		Maximum Limits of -						
	Test Weight per bushel		Damaged Kernels		Foreign material (percent)	Shrunken and broken kernels (percent)	Defects <u>3/</u> (percent)	Wheat of other classes <u>4/</u>	
	Hard Red Spring Wheat or White Club Wheat <u>1/</u> (pounds)	All other classes and subclasses (pounds)	Heat damage (part of total) (percent)	Total <u>2/</u> (percent)				Contrasting classes (percent)	Total <u>5/</u> (percent)
U.S. No. 1	58.0	60.0	0.2	2.0	0.4	3.0	3.0	1.0	3.0
U.S. No. 2	57.0	58.0	0.2	4.0	0.7	5.0	5.0	2.0	5.0
U.S. No. 3	55.0	56.0	0.5	7.0	1.3	8.0	8.0	3.0	10.0
U.S. No. 4	53.0	54.0	1.0	10.0	3.0	12.0	12.0	10.0	10.0
U.S. No. 5	50.0	51.0	3.0	15.0	5.0	20.0	20.0	10.0	10.0

U.S. Sample Grade:

U.S. Sample Grade is wheat that:

- (a) Does not meet the requirements for grades U.S. No. 1, 2, 3, 4, or 5; or
- (b) Contains 4 or more stones or any number of stones which have an aggregate weight in excess of 0.1 percent of the sample weight, 1 or more pieces of glass, 3 or more crotalaria seeds (*Crotalaria* spp.), 2 or more castor beans (*Ricinus communis* L.), 4 or more particles of an unknown foreign substance(s) or a commonly recognized harmful or toxic substance(s), 2 or more rodent pellets, bird dropping, or an equivalent quantity of other animal filth per 1,000 grams of wheat; or
- (c) Contains 5 or more animal filth, castor beans, crotalaria seeds, glass, stones, or unknown foreign substance(s) in any combination; or
- (d) Has a musty, sour, or commercially objectionable foreign odor (except smut or garlic odor); or
- (e) Is heating or otherwise of distinctly low quality.
- (f) Contains more than 31 insect-damaged kernels in 100 grams.

1/ These requirements also apply when Hard Red Spring or White Club wheat predominates in a sample of Mixed wheat.

2/ Includes heat-damaged kernels.

3/ Includes damaged kernels (total), foreign material, shrunken and broken kernels.

4/ Unclassed wheat of any grade may contain not more than 10.0 percent of wheat of other classes.

5/ Includes contrasting classes.

Wheat

Definitions

Test weight (lb/bu) is pounds of grain per Winchester bushel, determined by an approved device after the removal of dockage.

Test weight (kg/hl) is the metric system equivalent to pounds per bushel. Kilograms per hectoliter are calculated as follows:

For **Durum** wheat, multiply pounds per bushel by 1.292 and add 0.630. For **all other classes of wheat**, multiply pounds per bushel by 1.292 and add 1.419.

Heat-damaged kernels are kernels, pieces of wheat kernels, and other grains which have been materially discolored and damaged by heat.

Damaged kernels (total) are kernels, pieces of wheat kernels, and other grains that are badly ground-damaged, badly weather-damaged, diseased, frost-damaged, heat-damaged, insect bored, mold-damaged, sprout-damaged, or otherwise materially damaged.

Foreign material is all matter other than wheat which remains in a sample after removal of dockage and shrunken and broken kernels.

Shrunken and broken kernels are kernels, kernel pieces, and other matter that pass through a 0.064-by 3/8-inch oblong-hole sieve.

Total defects are the sum of three factors: damaged kernels (total), shrunken and broken kernels, and foreign material. In the factor summary tables, the average values listed for total defects may not equal the sum of the component factor averages due to rounding.

Dockage includes all matter other than wheat that can be removed from the original sample by use of an approved device. The percentage of dockage in a sample does not affect the numerical grade.

Moisture is the water content of grain as determined by an approved electronic moisture meter. The percentage of moisture in a sample does not affect the numerical grade.

Contrasting classes include:

* Durum, Soft White, and Unclassed wheats in the classes Hard Red Spring and Hard Red Winter wheats.

* Hard Red Spring, Hard Red Winter, Hard White, Soft Red Winter, Soft White, and Unclassed wheats in the class Durum wheat.

* Durum and Unclassed wheats in the class Soft Red Winter wheat.

* Durum, Hard Red Spring, Hard Red Winter, Soft Red Winter and Unclassed wheats in the classes Hard White wheat and Soft White wheat.

Wheat of other classes is any class that is mixed with the predominant class.

Protein is the protein content of grain as determined by an approved near infrared transmittance (NIRT) instrument calibrated against a Combustion Nitrogen Analyzer, or CNA (percent nitrogen multiplied by 5.7). The percentage of protein in a sample does not affect the numerical grade. Protein is certified on a 12 percent moisture basis.

Mixed wheat is a combination of classes of wheat which does not meet the minimum requirements of a specific class.

Table 1. U.S. Wheat Exports: Number of lots and quantity exported by class and grade, 2008-2010

Class	Grade	2008		2009		2010	
		No. of Lots	Metric Tons	No. of Lots	Metric Tons	No. of Lots	Metric Tons
Hard Red	U.S. No. 1	65	516,065	66	553,185	62	492,109
Winter Wheat	U.S. No. 2	671	12,740,901	522	7,346,859	686	11,875,031
	U.S. No. 3	8	94,776	3	12,471	2	39,376
	U.S. No. 4	--	--	--	--	1	9,163
	Not Inspected	2	21,419	--	--	1	8,412
	All lots	746	13,373,161	591	7,912,515	752	12,424,091
Hard Red	U.S. No. 1	85	911,258	80	892,620	89	997,293
Spring Wheat	U.S. No. 2	405	5,037,709	382	4,141,811	466	5,966,272
	U.S. No. 3	1	4,100	--	--	--	--
	U.S. No. 4	--	--	1	1,636	--	--
	Not Inspected	1	25,365	2	29,566	--	--
	All lots	492	5,978,432	465	5,065,633	555	6,963,555
Soft Red	U.S. No. 1	1	4,094	--	--	--	--
Winter Wheat	U.S. No. 2	309	4,542,990	262	2,877,143	206	1,658,794
	U.S. No. 3	8	61,375	24	175,993	42	311,853
	U.S. Sample Grade	--	--	--	--	1	4,249
	Not inspected	--	--	3	35,212	--	--
	All lots	318	4,608,459	289	3,088,348	249	1,975,076
Durum Wheat	U.S. No. 1	42	389,552	54	616,969	53	666,643
	U.S. No. 2	20	172,864	25	192,711	43	310,736
	U.S. No. 3	5	17,184	2	33,179	8	101,064
	U.S. No. 4	3	7,903	--	--	--	--
	U.S. No. 5	--	--	--	--	2	38,534
	U.S. Sample Grade	--	--	1	9,117	1	6,747
	All lots	70	587,503	82	851,976	107	1,123,724
Soft White Wheat	U.S. No. 1	126	701,876	110	653,765	120	722,996
	U.S. No. 2	151	2,660,816	197	3,407,776	200	3,331,900
	U.S. No. 3	1	7,875	--	--	1	969
	U.S. No. 4	--	--	--	--	4	212,622
	Not Inspected	--	--	--	--	1	735
	All lots	278	3,370,567	307	4,061,541	326	4,269,222
Hard White Wheat	U.S. No. 1	5	21,107	--	--	--	--
	U.S. No. 2	4	90,309	1	9,918	--	--
	U.S. No. 4	3	18,281	12	103,960	12	123,645
	All lots	12	129,697	13	113,878	12	123,645

-- = No lots reported in the category.

continued

Table 1. U.S. Wheat Exports: Number of lots and quantity exported by class and grade, 2008-2010, continued

Class	Grade	2008		2009		2010	
		No. of Lots	Metric Tons	No. of Lots	Metric Tons	No. of Lots	Metric Tons
Mixed Wheat	U.S. No 2	6	24,294	1	3,003	2	55,940
	U.S. Sample Grade	--	--	--	--	1	1,369
	All lots	6	24,294	1	3,003	3	57,309
All Classes	U.S. No. 1	324	2,543,952	310	2,716,539	324	2,879,031
	U.S. No. 2	1,566	25,269,883	1,390	17,979,221	1,603	23,198,853
	U.S. No. 3	23	185,310	29	221,643	53	453,262
	U.S. No. 4	6	26,184	13	105,596	17	345,430
	U.S. No. 5	--	--	--	--	2	38,534
	U.S. Sample Grade	--	--	3	35,212	3	12,365
	Not Inspected	3	46,784	3	38,683	2	9,147
	All lots	1,922	28,072,113	1,748	21,096,894	2,004	26,936,632

-- = No lots reported in the category.

Table 2. Summary of export Hard Red Winter wheat quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 1	60.0	65	62.0	60.5	63.7	66	61.7	60.5	62.9	62	61.7	60.3	63.0
	U.S. No. 2	58.0	671	60.7	58.2	64.0	522	60.7	58.1	63.7	686	80.4	76.5	83.0
	U.S. No. 3	56.0	8	59.1	58.2	61.2	3	59.8	59.5	61.0	2	81.7	76.1	81.8
	U.S. No. 4	54.0	--	--	--	--	--	--	--	--	1	61.0	61.0	61.0
	All lots	N/A	744	60.7	58.2	64.0	591	60.8	58.1	63.7	751	61.1	57.8	63.1
Test Weight (kg/hl)	U.S. No. 1	N/A	65	81.5	79.6	83.7	66	81.1	79.5	82.7	62	81.1	79.3	82.8
	U.S. No. 2	N/A	671	79.8	76.6	84.1	522	79.9	76.5	83.7	686	80.4	76.5	83.0
	U.S. No. 3	N/A	8	77.8	76.6	80.5	3	78.6	78.3	80.2	2	81.7	76.1	81.8
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	1	80.3	80.3	80.3
	All lots	N/A	744	79.8	76.6	84.1	591	80.0	76.5	83.7	751	80.4	76.1	83.0
Moisture	U.S. No. 1	N/A	65	9.9	8.6	11.8	66	10.6	9.2	12.4	62	11.0	8.6	12.2
	U.S. No. 2	N/A	671	11.4	8.8	12.8	522	11.3	9.0	13.1	686	11.3	8.7	12.7
	U.S. No. 3	N/A	8	12.0	11.8	12.5	3	11.8	11.5	13.3	2	11.2	11.2	12.4
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	1	9.2	9.2	9.2
	All lots	N/A	744	11.4	8.6	12.8	591	11.2	9.0	13.3	751	11.3	8.6	12.7
Heat-damaged Kernels	U.S. No. 1	0.2	65	0.0	0.0	0.1	66	0.0	0.0	0.0	62	0.0	0.0	0.0
	U.S. No. 2	0.2	671	0.0	0.0	0.2	522	0.0	0.0	0.1	686	0.0	0.0	0.1
	U.S. No. 3	0.5	8	0.0	0.0	0.0	3	0.0	0.0	0.0	2	0.0	0.0	0.0
	U.S. No. 4	1.0	--	--	--	--	--	--	--	--	1	0.0	0.0	0.0
	All lots	N/A	744	0.0	0.0	0.2	591	0.0	0.0	0.1	751	0.0	0.0	0.1
Damaged Kernels (Total)	U.S. No. 1	2.0	65	0.2	0.0	0.7	66	0.2	0.0	1.0	62	0.4	0.0	1.7
	U.S. No. 2	4.0	671	1.2	0.0	3.2	522	1.0	0.0	3.6	686	0.9	0.0	3.3
	U.S. No. 3	7.0	8	2.5	1.5	3.8	3	1.2	0.7	1.3	2	0.7	0.7	1.5
	U.S. No. 4	10.0	--	--	--	--	--	--	--	--	1	0.0	0.0	0.0
	All lots	N/A	744	1.1	0.0	3.8	591	0.9	0.0	3.6	751	0.9	0.0	3.3
Foreign Material	U.S. No. 1	0.4	65	0.1	0.0	0.2	66	0.1	0.0	0.3	62	0.1	0.0	0.3
	U.S. No. 2	0.7	671	0.2	0.0	0.7	522	0.2	0.0	0.7	686	0.2	0.0	0.7
	U.S. No. 3	1.3	8	0.4	0.2	1.0	3	0.5	0.1	0.6	2	0.1	0.1	0.2
	U.S. No. 4	3.0	--	--	--	--	--	--	--	--	1	0.1	0.1	0.1
	All lots	N/A	744	0.2	0.0	1.0	591	0.2	0.0	0.7	751	0.2	0.0	0.7
Shrunken and Broken	U.S. No. 1	3.0	65	1.6	0.7	2.2	66	1.5	0.7	2.5	62	1.1	0.6	1.8
	U.S. No. 2	5.0	671	1.6	0.5	2.9	522	1.5	0.3	2.5	686	1.4	0.8	2.3
	U.S. No. 3	8.0	8	2.2	1.7	2.7	3	1.4	0.6	1.6	2	1.3	0.8	1.3
	U.S. No. 4	12.0	--	--	--	--	--	--	--	--	1	1.0	1.0	1.0
	All lots	N/A	744	1.6	0.5	2.9	591	1.5	0.3	2.5	751	1.4	0.6	2.3
Total Defects ¹	U.S. No. 1	3.0	65	1.8	0.9	2.8	66	1.8	0.8	2.6	62	1.6	0.9	2.8
	U.S. No. 2	5.0	671	2.9	0.8	5.0	522	2.7	0.8	5.0	686	2.5	0.9	4.9
	U.S. No. 3	8.0	8	5.2	3.9	6.5	3	3.1	1.4	3.5	2	2.1	2.1	2.5
	U.S. No. 4	12.0	--	--	--	--	--	--	--	--	1	1.1	1.1	1.1
	All lots	N/A	744	2.9	0.8	6.5	591	2.6	0.8	5.0	751	2.5	0.9	4.9

continued

Table 2. Summary of export Hard Red Winter wheat quality, 2008-2010, continued

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Dockage	U.S. No. 1	N/A	65	0.3	0.2	0.3	66	0.3	0.1	0.3	62	0.3	0.2	0.9
	U.S. No. 2	N/A	671	0.6	0.1	1.4	522	0.6	0.1	1.1	686	0.6	0.1	1.5
	U.S. No. 3	N/A	8	0.8	0.6	0.9	3	0.7	0.7	0.7	2	0.6	0.6	1.3
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	1	0.7	0.7	0.7
	All lots	N/A	744	0.6	0.1	1.4	591	0.5	0.1	1.1	751	0.6	0.1	1.5
Wheat of Other Classes	U.S. No. 1	3.0	65	0.9	0.0	2.3	66	0.6	0.1	1.9	62	0.8	0.0	2.1
	U.S. No. 2	5.0	671	1.7	0.0	4.7	522	1.5	0.0	4.5	686	1.1	0.0	4.4
	U.S. No. 3	10.0	8	2.0	0.6	3.4	3	5.3	5.2	5.4	2	1.0	0.4	1.0
	U.S. No. 4	10.0	--	--	--	--	--	--	--	--	1	3.7	3.7	3.7
	All lots	N/A	744	1.6	0.0	4.7	591	1.4	0.0	5.4	751	1.1	0.0	4.4
Contrasting Classes	U.S. No. 1	1.0	65	0.3	0.0	0.9	66	0.3	0.0	1.0	62	0.3	0.0	1.0
	U.S. No. 2	2.0	671	0.1	0.0	1.6	522	0.1	0.0	1.6	686	0.1	0.0	1.4
	U.S. No. 3	3.0	8	0.0	0.0	0.0	3	0.0	0.0	0.0	2	0.0	0.0	0.0
	U.S. No. 4	10.0	--	--	--	--	--	--	--	--	1	3.7	3.7	3.7
	All lots	N/A	744	0.1	0.0	1.6	591	0.1	0.0	1.6	751	0.1	0.0	3.7
Protein (as is basis)	U.S. No. 1	N/A	65	12.4	11.5	14.1	66	12.6	11.6	13.3	62	12.3	11.5	13.1
	U.S. No. 2	N/A	668	11.9	10.2	13.6	522	12.0	10.6	14.4	685	11.8	10.3	13.4
	U.S. No. 3	N/A	8	11.7	11.4	11.9	3	12.1	11.5	12.2	1	12.1	12.1	12.1
	All lots	N/A	741	11.9	10.2	14.1	591	12.0	10.6	14.4	748	11.8	10.3	13.4
Protein (12% moisture)	U.S. No. 1	N/A	65	12.2	11.5	13.7	66	12.4	11.5	13.1	62	12.1	11.5	13.0
	U.S. No. 2	N/A	668	11.8	10.2	13.3	522	11.9	10.5	14.4	685	11.7	10.2	13.5
	U.S. No. 3	N/A	8	11.7	11.4	11.9	3	12.0	11.7	12.1	1	12.0	12.0	12.0
	All lots	N/A	741	11.8	10.2	13.7	591	11.9	10.5	14.4	748	11.7	10.2	13.5

N/A = Does not apply.

-- = No lots reported in this category.

¹The sum of the component factor averages may not equal the average for this factor due to rounding.

Figure 1. HRW Dockage Distribution, All Grades

U.S. WHEAT EXPORTED, 2010
DISTRIBUTION FOR DOCKAGE – ALL GRADES
HRW




Figure 2. HRW Protein Distribution, All Grades

U.S. WHEAT EXPORTED, 2010
DISTRIBUTION FOR PROTEIN (12% M) – ALL GRADES
HRW




Table 3. Summary of export Hard Red Spring wheat quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 1	58.0	85	61.6	59.9	63.5	80	61.6	60.2	63.3	89	62.0	60.2	63.5
	U.S. No. 2	57.0	405	61.7	59.1	63.8	382	62.0	59.8	63.9	466	62.2	59.9	64.0
	U.S. No. 3	55.0	1	61.4	61.4	61.4	--	--	--	--	--	--	--	--
	U.S. No. 4	53.0	--	--	--	--	1	63	63	63	--	--	--	--
	All lots	N/A	491	61.7	59.1	63.8	463	62.0	59.8	63.9	555	62.2	59.9	64.0
Test Weight (kg/hl)	U.S. No. 1	N/A	85	81.1	78.8	83.4	80	81.0	79.2	83.2	89	81.6	79.2	83.5
	U.S. No. 2	N/A	405	81.1	77.8	83.8	382	81.6	78.6	84.0	466	81.8	78.8	84.1
	U.S. No. 3	N/A	1	80.7	80.7	80.7	--	--	--	--	--	--	--	--
	U.S. No. 4	N/A	--	--	--	--	1	83	83	83	--	--	--	--
	All lots	N/A	491	81.1	77.8	83.8	463	81.5	78.6	84.0	555	81.7	78.8	84.1
Moisture	U.S. No. 1	N/A	85	11.4	8.9	13.1	80	11.3	9.8	13.3	89	11.6	8.9	13.2
	U.S. No. 2	N/A	405	12.1	9.0	13.5	382	12.3	9.3	13.7	466	12.5	9.1	13.5
	U.S. No. 3	N/A	1	12.8	12.8	12.8	--	--	--	--	--	--	--	--
	U.S. No. 4	N/A	--	--	--	--	1	13	13	13	--	--	--	--
	All lots	N/A	491	12.0	8.9	13.5	463	12.1	9.3	13.7	555	12.3	8.9	13.5
Heat-damaged Kernels	U.S. No. 1	0.2	85	0.0	0.0	0.1	80	0.0	0.0	0.0	89	0.0	0.0	0.1
	U.S. No. 2	0.2	405	0.0	0.0	0.1	382	0.0	0.0	0.2	466	0.0	0.0	0.1
	U.S. No. 3	0.5	1	0.0	0.0	0.0	--	--	--	--	--	--	--	--
	U.S. No. 4	1.0	--	--	--	--	1	0	0	0	--	--	--	--
	All lots	N/A	491	0.0	0.0	0.1	463	0.0	0.0	0.2	555	0.1	0.0	0.6
Damaged Kernels (Total)	U.S. No. 1	2.0	85	0.3	0.0	1.7	80	0.3	0.0	1.2	89	0.4	0.0	1.6
	U.S. No. 2	4.0	405	0.6	0.0	3.4	382	0.6	0.0	2.8	466	0.6	0.0	2.5
	U.S. No. 3	7.0	1	4.5	4.5	4.5	--	--	--	--	--	--	--	--
	U.S. No. 4	10.0	--	--	--	--	1	1	1	1	--	--	--	--
	All lots	N/A	491	0.6	0.0	4.5	463	0.6	0.0	2.8	555	0.6	0.0	2.5
Foreign Material	U.S. No. 1	0.4	85	0.1	0.0	0.3	80	0.1	0.0	0.4	89	0.1	0.0	0.2
	U.S. No. 2	0.7	405	0.1	0.0	0.5	382	0.1	0.0	0.5	466	0.1	0.0	0.6
	U.S. No. 3	1.3	1	0.1	0.1	0.1	--	--	--	--	--	--	--	--
	U.S. No. 4	3.0	--	--	--	--	1	0	0	0	--	--	--	--
	All lots	N/A	491	0.1	0.0	0.5	463	0.1	0.0	0.5	555	0.1	0.0	0.6
Shrunken and Broken	U.S. No. 1	3.0	85	1.4	0.4	2.1	80	1.2	0.5	2.0	89	1.0	0.5	1.5
	U.S. No. 2	5.0	405	1.2	0.5	2.2	382	1.0	0.4	1.9	466	0.8	0.4	2.0
	U.S. No. 3	8.0	1	1.4	1.4	1.4	--	--	--	--	--	--	--	--
	U.S. No. 4	12.0	--	--	--	--	1	1	1	1	--	--	--	--
	All lots	N/A	491	1.2	0.4	2.2	463	1.0	0.4	2.0	555	0.9	0.4	2.0

continued

Table 3. Summary of export Hard Red Spring wheat quality, 2008-2010, continued

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Total Defects ¹	U.S. No. 1	3.0	85	1.8	0.5	2.9	80	1.5	0.0	2.7	89	1.5	0.6	2.8
	U.S. No. 2	5.0	405	1.9	0.7	5.0	382	1.7	0.0	4.0	466	1.6	0.8	3.5
	U.S. No. 3	8.0	1	6.0	6.0	6.0	--	--	--	--	--	--	--	--
	U.S. No. 4	12.0	--	--	--	--	1	2	2	2	--	--	--	--
	All lots	N/A	491	1.9	0.5	6.0	463	1.7	0.0	4.0	555	1.6	0.6	3.5
Dockage	U.S. No. 1	N/A	85	0.3	0.2	0.8	80	0.3	0.1	0.7	89	0.3	0.1	0.8
	U.S. No. 2	N/A	405	0.5	0.2	1.9	382	0.4	0.1	1.0	465	0.4	0.1	1.0
	U.S. No. 3	N/A	1	0.5	0.5	0.5	--	--	--	--	--	--	--	--
	U.S. No. 4	N/A	--	--	--	--	1	1	1	1	--	--	--	--
	All lots	N/A	491	0.4	0.2	1.9	463	0.4	0.1	1.0	554	0.4	0.1	1.0
Wheat of	U.S. No. 1	3.0	85	0.8	0.0	2.7	80	0.7	0.0	2.5	89	0.5	0.0	1.8
Other Classes	U.S. No. 2	5.0	405	1.0	0.0	4.7	382	0.8	0.0	5.0	466	0.6	0.0	4.1
	U.S. No. 3	10.0	1	4.1	4.1	4.1	--	--	--	--	--	--	--	--
	U.S. No. 4	10.0	--	--	--	--	1	4	4	4	--	--	--	--
	All lots	N/A	491	1.0	0.0	4.7	463	0.8	0.0	5.0	555	0.6	0.0	4.1
Contrasting Classes	U.S. No. 1	1.0	85	0.2	0.0	0.8	80	0.3	0.0	1.0	89	0.2	0.0	0.9
	U.S. No. 2	2.0	405	0.2	0.0	1.4	382	0.2	0.0	1.6	466	0.2	0.0	1.2
	U.S. No. 3	3.0	1	0.1	0.1	0.1	--	--	--	--	--	--	--	--
	U.S. No. 4	10.0	--	--	--	--	1	3	3	3	--	--	--	--
	All lots	N/A	491	0.2	0.0	1.4	463	0.2	0.0	3.4	555	0.2	0.0	1.2
Protein (as is basis)	U.S. No. 1	N/A	85	14.4	13.9	15.2	80	14.4	13.3	14.9	89	13.9	12.9	14.5
	U.S. No. 2	N/A	405	14.1	12.4	15.4	378	13.8	11.9	14.9	462	13.5	11.9	14.7
	U.S. No. 3	N/A	1	15.2	15.2	15.2	--	--	--	--	--	--	--	--
	U.S. No. 4	N/A	--	--	--	--	1	12	12	12	--	--	--	--
	All lots	N/A	491	14.1	12.4	15.4	459	13.9	11.9	14.9	551	13.5	11.9	14.7
Protein (12% moisture)	U.S. No. 1	N/A	85	14.3	14.0	15.2	80	14.2	13.5	14.8	89	13.9	12.9	14.4
	U.S. No. 2	N/A	405	14.1	12.5	15.5	378	13.8	12.1	14.7	462	13.5	12.0	14.6
	U.S. No. 3	N/A	1	15.3	15.3	15.3	--	--	--	--	--	--	--	--
	U.S. No. 4	N/A	--	--	--	--	1	13	13	13	--	--	--	--
	All lots	N/A	491	14.1	12.5	15.5	459	13.9	12.1	14.8	551	13.6	12.0	14.6

N/A = Does not apply.

-- = No lots reported in this category.

¹The sum of the component factor averages may not equal the average for this factor due to rounding.

Figure 3. HRS Dockage Distribution, All Grades

**U.S. WHEAT EXPORTED, 2010
DISTRIBUTION FOR DOCKAGE – ALL GRADES
HRS**




Figure 4. HRS Protein Distribution, All Grades

U.S. WHEAT EXPORTED, 2010
DISTRIBUTION FOR PROTEIN (12% M) – ALL GRADES
HRS




Table 4. Summary of export Soft Red Winter wheat quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 1	60.0	1	60.1	60.1	60.1	--	--	--	--	--	--	--	--
	U.S. No. 2	58.0	309	59.8	0.0	62.9	262	58.9	58.0	60.6	206	58.5	58.0	60.2
	U.S. No. 3	56.0	8	59.7	58.7	60.1	24	57.8	56.8	60.5	42	57.6	56.1	58.9
	U.S. Sample Grade	N/A	--	--	--	--	3	58.9	58.7	59.1	1	59.2	59.2	59.2
	All lots	N/A	318	59.8	0.0	62.9	289	58.8	56.8	60.6	249	58.4	56.1	60.2
Test Weight (kg/hl)	U.S. No. 1	N/A	1	79.1	79.1	79.1	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	309	78.7	1.4	82.7	262	77.5	76.3	79.7	206	77.1	76.3	79.1
	U.S. No. 3	N/A	8	78.6	77.3	79.1	24	76.1	74.8	79.6	42	75.8	73.9	77.5
	U.S. Sample Grade	N/A	--	--	--	--	3	77.5	77.3	77.8	1	77.9	77.9	77.9
	All lots	N/A	318	78.7	1.4	82.7	289	77.4	74.8	79.7	249	76.9	73.9	79.1
Moisture	U.S. No. 1	N/A	1	12.7	12.7	12.7	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	308	12.6	11.1	13.5	262	12.5	11.9	13.4	206	12.5	11.9	13.4
	U.S. No. 3	N/A	8	12.3	12.0	12.9	24	12.5	11.8	13.3	42	12.4	11.7	13.5
	U.S. Sample Grade	N/A	--	--	--	--	3	13.0	12.9	13.2	1	12.1	12.1	12.1
	All lots	N/A	317	12.6	11.1	13.5	289	12.5	11.8	13.4	249	12.5	11.7	13.5
Heat-damaged Kernels	U.S. No. 1	0.2	1	0.0	0.0	0.0	--	--	--	--	--	--	--	--
	U.S. No. 2	0.2	309	0.0	0.0	0.2	262	0.0	0.0	0.2	206	0.0	0.0	0.2
	U.S. No. 3	0.5	8	0.0	0.0	0.0	24	0.0	0.0	0.1	42	0.0	0.0	0.1
	U.S. Sample Grade	N/A	--	--	--	--	3	0.4	0.2	0.6	1	2.0	2.0	2.0
	All lots	N/A	318	0.0	0.0	0.2	289	0.0	0.0	0.6	249	0.0	0.0	2.0
Damaged Kernels (Total)	U.S. No. 1	2.0	1	0.9	0.9	0.9	--	--	--	--	--	--	--	--
	U.S. No. 2	4.0	309	1.4	0.0	3.7	262	2.0	0.2	3.8	206	2.3	0.6	4.0
	U.S. No. 3	7.0	8	2.0	0.5	3.7	24	2.1	0.6	4.0	42	2.9	0.9	4.3
	U.S. Sample Grade	N/A	--	--	--	--	3	44.4	25.4	59.4	1	24.5	24.5	24.5
	All lots	N/A	318	1.4	0.0	3.7	289	2.5	0.2	59.4	249	2.5	0.6	24.5
Foreign Material	U.S. No. 1	0.4	1	0.3	0.3	0.3	--	--	--	--	--	--	--	--
	U.S. No. 2	0.7	309	0.1	0.0	0.7	262	0.1	0.0	0.4	206	0.1	0.0	0.5
	U.S. No. 3	1.3	8	0.3	0.1	0.4	24	0.1	0.0	0.2	42	0.1	0.0	0.5
	U.S. Sample Grade	N/A	--	--	--	--	3	0.1	0.1	0.1	1	0.1	0.1	0.1
	All lots	N/A	318	0.1	0.0	0.7	289	0.1	0.0	0.4	249	0.1	0.0	0.5
Shrunken and Broken	U.S. No. 1	3.0	1	0.6	0.6	0.6	--	--	--	--	--	--	--	--
	U.S. No. 2	5.0	309	0.7	0.0	1.5	262	0.7	0.2	1.5	206	0.8	0.3	1.5
	U.S. No. 3	8.0	8	0.8	0.4	1.1	24	0.8	0.3	1.0	42	1.0	0.6	1.8
	U.S. Sample Grade	N/A	--	--	--	--	3	1.0	0.9	1.0	1	0.8	0.8	0.8
	All lots	N/A	318	0.7	0.0	1.5	289	0.7	0.2	1.5	249	0.8	0.3	1.8

continued

Table 4. Summary of export Soft Red Winter wheat quality, 2008-2010, continued

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Total Defects ¹	U.S. No. 1	3.0	1	1.8	1.8	1.8	--	--	--	--	--	--	--	--
	U.S. No. 2	5.0	309	2.2	0.0	4.5	262	2.8	0.7	4.8	206	3.3	1.1	5.0
	U.S. No. 3	8.0	8	3.1	1.1	5.2	24	3.1	1.3	4.8	42	3.9	1.7	5.9
	U.S. Sample Grade	N/A	--	--	--	--	3	45.4	26.5	60.5	1	25.4	25.4	25.4
	All lots	N/A	318	2.2	0.0	5.2	289	3.3	0.7	60.5	249	3.4	1.1	25.4
Dockage	U.S. No. 1	N/A	1	0.7	0.7	0.7	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	307	0.7	0.3	1.7	262	0.7	0.4	1.3	203	0.8	0.3	1.5
	U.S. No. 3	N/A	8	1.1	0.5	1.4	24	0.8	0.3	1.2	42	0.8	0.3	1.2
	U.S. Sample Grade	N/A	--	--	--	--	3	1.1	1.0	1.2	1	0.7	0.7	0.7
	All lots	N/A	316	0.7	0.3	1.7	289	0.7	0.3	1.3	246	0.8	0.3	1.5
Wheat of Other Classes	U.S. No. 1	3.0	1	0.0	0.0	0.0	--	--	--	--	--	--	--	--
	U.S. No. 2	5.0	309	0.4	0.0	4.6	262	0.6	0.0	4.1	206	0.4	0.0	3.3
	U.S. No. 3	10.0	8	0.5	0.0	1.0	24	0.5	0.0	2.0	42	0.7	0.0	5.0
	U.S. Sample Grade	N/A	--	--	--	--	3	0.1	0.0	0.2	1	0.1	0.1	0.1
	All lots	N/A	318	0.4	0.0	4.6	289	0.6	0.0	4.1	249	0.5	0.0	5.0
Contrasting Classes	U.S. No. 1	1.0	1	0.0	0.0	0.0	--	--	--	--	--	--	--	--
	U.S. No. 2	2.0	309	0.0	0.0	0.3	262	0.0	0.0	0.9	206	0.0	0.0	0.9
	U.S. No. 3	3.0	8	0.0	0.0	0.0	24	0.0	0.0	0.0	42	0.0	0.0	0.5
	U.S. Sample Grade	N/A	--	--	--	--	3	0.0	0.0	0.1	1	0.0	0.0	0.0
	All lots	N/A	318	0.0	0.0	0.3	289	0.0	0.0	0.9	249	0.0	0.0	0.9
Protein (as is basis)	U.S. No. 1	N/A	1	9.9	9.9	9.9	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	283	10.0	8.9	11.8	248	9.9	8.9	10.6	197	9.9	0.4	10.9
	U.S. No. 3	N/A	3	10.4	10.2	10.6	18	10.0	9.4	10.5	39	9.9	9.0	10.9
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	0	--	--	--
	All lots	N/A	287	10.0	8.9	11.8	266	9.9	8.9	10.6	236	9.9	0.4	10.9
Protein (12% moisture)	U.S. No. 1	N/A	1	10.0	10.0	10.0	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	283	10.1	9.1	11.7	248	10.0	9.0	10.7	197	10.0	0.4	10.9
	U.S. No. 3	N/A	3	10.5	10.2	10.6	18	10.0	9.5	10.5	39	9.9	9.2	10.9
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	0	--	--	10.9
	All lots	N/A	287	10.1	9.1	11.7	266	10.0	9.0	10.7	236	10.0	0.4	10.9

N/A = Does not apply.

-- = No lots reported in this category.

¹The sum of the component factor averages may not equal the average for this factor due to rounding.

Figure 5. SRW Dockage Distribution, All Grades

**U.S. WHEAT EXPORTED, 2010
DISTRIBUTION FOR DOCKAGE – ALL GRADES
SRW**




Figure 6. SRW Protein Distribution, All Grades

U.S. WHEAT EXPORTED, 2010
DISTRIBUTION FOR PROTEIN (12% M) – ALL GRADES
SRW




Table 5. Summary of export Durum wheat quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of Lots	Avg.	Low	High	No. of Lots	Avg.	Low	High	No. of Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 1	60.0	42	62.2	60.1	64.3	54	62.2	60.8	63.5	53	62.1	61.0	63.3
	U.S. No. 2	58.0	20	60.9	59.7	63.4	25	61.3	58.6	62.8	43	61.9	60.1	62.7
	U.S. No. 3	56.0	5	61.4	60.2	63.3	2	62.2	62.0	62.6	8	60.1	59.0	61.3
	U.S. No. 4	54.0	3	62.2	60.6	62.7	--	--	--	--	--	--	--	--
	U.S. No. 5	51.0	--	--	--	--	--	--	--	--	2	59.4	59.0	59.7
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	62.2	62.2	62.2
	All lots	N/A	70	61.8	59.7	64.3	81	62.0	58.6	63.5	107	61.8	59.0	63.3
Test Weight (kg/hl)	U.S. No. 1	N/A	42	81.0	78.3	83.7	54	81.0	79.1	82.7	53	80.8	79.4	82.5
	U.S. No. 2	N/A	20	79.3	77.8	82.5	25	79.8	76.3	81.7	43	80.6	78.3	81.6
	U.S. No. 3	N/A	5	80.0	78.4	82.4	2	81.0	80.7	81.5	8	78.3	76.8	79.8
	U.S. No. 4	N/A	3	81.0	78.9	81.6	--	--	--	--	--	--	--	--
	U.S. No. 5	N/A	--	--	--	--	--	--	--	--	2	77.4	76.9	77.8
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	81.0	81.0	81.0
	All lots	N/A	70	80.4	77.8	83.7	81	80.7	76.3	82.7	107	80.4	76.8	82.5
Moisture	U.S. No. 1	N/A	42	8.3	6.4	12.7	54	10.4	6.6	12.6	53	10.2	6.5	12.6
	U.S. No. 2	N/A	20	11.8	8.5	12.5	25	12.0	7.9	12.8	43	12.0	7.5	12.8
	U.S. No. 3	N/A	5	10.3	7.4	12.0	2	11.3	8.1	12.5	8	12.6	12.3	13.2
	U.S. No. 4	N/A	3	8.9	7.9	11.6	--	--	--	--	--	--	--	--
	U.S. No. 5	N/A	--	--	--	--	--	--	--	--	2	12.4	12.4	12.5
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	7.9	7.9	7.9
	All lots	N/A	70	9.4	6.4	12.7	81	10.8	6.6	12.8	107	11.0	6.5	13.2
Heat-damaged Kernels	U.S. No. 1	0.2	42	0.0	0.0	0.1	54	0.0	0.0	0.1	53	0.0	0.0	0.1
	U.S. No. 2	0.2	20	0.0	0.0	0.1	25	0.0	0.0	0.0	43	0.0	0.0	0.0
	U.S. No. 3	0.5	5	0.0	0.0	0.1	2	0.0	0.0	0.1	8	0.0	0.0	0.0
	U.S. No. 4	1.0	3	0.1	0.0	0.1	--	--	--	--	--	--	--	--
	U.S. No. 5	3.0	--	--	--	--	--	--	--	--	2	0.0	0.0	0.0
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	0.0	0.0	0.0
	All lots	N/A	70	0.0	0.0	0.1	81	0.0	0.0	0.1	107	0.0	0.0	0.1
Damaged Kernels (Total)	U.S. No. 1	2.0	42	0.5	0.1	1.3	54	0.8	0.0	1.3	53	0.8	0.3	1.7
	U.S. No. 2	4.0	20	1.3	0.5	2.3	25	1.8	0.2	2.8	43	1.4	0.0	3.0
	U.S. No. 3	7.0	5	0.8	0.5	2.0	2	0.9	0.8	1.0	8	4.0	2.2	5.6
	U.S. No. 4	10.0	3	0.9	0.6	1.8	--	--	--	--	--	--	--	--
	U.S. No. 5	15.0	--	--	--	--	--	--	--	--	2	6.9	5.9	7.9
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	0.8	0.8	0.8
	All lots	N/A	70	0.8	0.1	2.3	81	1.0	0.0	2.8	107	1.5	0.0	7.9
Foreign Material	U.S. No. 1	0.4	42	0.1	0.0	0.3	54	0.1	0.0	0.3	53	0.1	0.1	0.3
	U.S. No. 2	0.7	20	0.2	0.0	0.5	25	0.2	0.1	0.7	43	0.2	0.0	0.5
	U.S. No. 3	1.3	5	0.4	0.1	1.0	2	0.4	0.1	1.1	8	0.2	0.1	0.3
	U.S. No. 4	3.0	3	0.3	0.1	0.7	--	--	--	--	--	--	--	--
	U.S. No. 5	5.0	--	--	--	--	--	--	--	--	2	0.2	0.2	0.2
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	0.7	0.7	0.7
	All lots	N/A	70	0.2	0.0	1.0	81	0.1	0.0	1.1	107	0.2	0.0	0.7

continued

Table 5. Summary of export Durum wheat quality, 2008-2010, continued

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of Lots	Avg.	Low	High	No. of Lots	Avg.	Low	High	No. of Lots	Avg.	Low	High
Shrunken and Broken	U.S. No. 1	3.0	42	0.8	0.3	1.9	54	0.9	0.3	1.6	53	0.9	0.4	1.2
	U.S. No. 2	5.0	20	1.6	0.5	2.1	25	1.3	0.8	1.8	43	1.0	0.5	1.7
	U.S. No. 3	8.0	5	1.4	1.0	1.8	2	0.9	0.8	1.0	8	1.3	1.1	1.6
	U.S. No. 4	12.0	3	1.3	1.1	2.1	--	--	--	--	--	--	--	--
	U.S. No. 5	20.0	--	--	--	--	--	--	--	--	2	1.5	1.4	1.6
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	1.1	1.1	1.1
	All lots	N/A	70	1.1	0.3	2.1	81	1.0	0.3	1.8	107	1.0	0.4	1.7
Total Defects ¹	U.S. No. 1	3.0	42	1.4	0.5	2.9	54	1.8	0.5	2.9	53	0.5	0.3	0.8
	U.S. No. 2	5.0	20	3.1	1.5	4.6	25	3.3	1.4	4.6	43	0.5	0.1	0.8
	U.S. No. 3	8.0	5	2.6	2.2	4.5	2	2.2	1.9	2.9	8	0.6	0.5	0.7
	U.S. No. 4	12.0	3	2.5	1.8	4.4	--	--	--	--	--	--	--	--
	U.S. No. 5	20.0	--	--	--	--	--	--	--	--	2	0.6	0.6	0.6
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	1.0	1.0	1.0
	All lots	N/A	70	2.0	0.5	4.6	81	2.2	0.5	4.6	107	0.5	0.1	1.0
Dockage	U.S. No. 1	N/A	42	0.5	0.3	0.7	54	0.5	0.3	0.9	53	0.5	0.3	0.8
	U.S. No. 2	N/A	20	0.6	0.3	1.0	25	0.6	0.3	0.9	43	0.5	0.1	0.8
	U.S. No. 3	N/A	5	0.6	0.5	0.9	2	0.6	0.5	1.0	8	0.6	0.5	0.7
	U.S. No. 4	N/A	3	0.6	0.5	1.0	--	--	--	--	--	--	--	--
	U.S. No. 5	N/A	--	--	--	--	--	--	--	--	2	0.6	0.6	0.6
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	1.0	1.0	1.0
	All lots	N/A	70	0.6	0.3	1.0	81	0.5	0.3	1.0	107	0.5	0.1	1.0
Contrasting Classes	U.S. No. 1	1.0	42	0.2	0.0	0.9	54	0.5	0.0	1.0	53	0.5	0.0	1.0
	U.S. No. 2	2.0	20	0.9	0.0	2.0	25	1.1	0.1	1.6	43	1.1	0.0	1.9
	U.S. No. 3	3.0	5	0.8	0.0	3.0	2	1.8	0.9	2.1	8	1.2	0.6	1.7
	U.S. No. 4	10.0	3	2.6	1.3	9.3	--	--	--	--	--	--	--	--
	U.S. No. 5	10.0	--	--	--	--	--	--	--	--	2	1.1	0.9	1.3
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	0.4	0.4	0.4
	All lots	N/A	70	0.5	0.0	9.3	81	0.7	0.0	2.1	107	0.7	0.0	1.9
Protein (as is basis)	U.S. No. 1	N/A	30	13.7	11.9	15.5	49	13.5	12.5	15.0	48	13.3	12.2	14.5
	U.S. No. 2	N/A	14	14.5	11.0	15.7	21	13.9	11.1	15.7	34	13.0	12.4	13.9
	U.S. No. 3	N/A	4	13.6	11.3	14.6	2	12.6	11.5	13.0	8	13.2	13.0	13.4
	U.S. No. 4	N/A	1	13.4	13.4	13.4	--	--	--	--	--	--	--	--
	U.S. No. 5	N/A	--	--	--	--	--	--	--	--	2	13.2	13.1	13.2
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	11.6	11.6	11.6
	All lots	N/A	49	14.0	11.0	15.7	72	13.6	11.1	15.7	93	13.2	11.6	14.5
Protein (12% moisture)	U.S. No. 1	N/A	30	13.1	11.4	15.4	49	13.3	12.0	14.9	48	13.0	11.6	14.2
	U.S. No. 2	N/A	14	14.5	10.6	15.6	21	14.0	10.7	15.6	34	13.1	11.8	13.9
	U.S. No. 3	N/A	4	13.4	10.8	14.5	2	12.5	11.0	13.1	8	13.3	13.0	13.5
	U.S. No. 4	N/A	1	12.8	12.8	12.8	--	--	--	--	--	--	--	--
	U.S. No. 5	N/A	--	--	--	--	--	--	--	--	2	13.2	13.2	13.3
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	11.1	11.1	11.1
	All lots	N/A	49	13.6	10.6	15.6	72	13.4	10.7	15.6	93	13.1	11.1	14.2

N/A=Does Not Apply, -- = No lots reported, 1. Sum of component factor avg. may not equal factor avg.

Figure 7. DU Dockage Distribution, All Grades

U.S. WHEAT EXPORTED, 2010
DISTRIBUTION FOR DOCKAGE – ALL GRADES
DU




Figure 8. DU Protein Distribution, All Grades

U.S. WHEAT EXPORTED, 2010
DISTRIBUTION FOR PROTEIN (12% M) – ALL GRADES
DU




Table 6. Summary of export Soft White wheat quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 1	60.0	126	61.4	60.2	62.8	110	61.1	60.4	63.2	120	61.2	60.0	62.8
	U.S. No. 2	58.0	151	61.2	58.4	62.7	197	60.9	58.5	63.2	200	61.3	60.0	62.9
	U.S. No. 3	56.0	1	60.8	60.8	60.8	--	--	--	--	1	59.3	59.3	59.3
	U.S. No. 4	54.0	--	--	--	--	--	--	--	--	4	61.3	61.0	61.5
	All lots	N/A	278	61.3	58.4	62.8	307	60.9	58.5	63.2	325	61.3	59.3	62.9
Test Weight (kg/hl)	U.S. No. 1	N/A	126	80.8	79.1	82.5	110	80.4	79.4	83.1	120	80.5	79.0	82.6
	U.S. No. 2	N/A	151	80.5	76.9	82.4	197	80.1	77.1	83.1	200	80.6	78.9	82.6
	U.S. No. 3	N/A	1	80.0	80.0	80.0	--	--	--	--	1	78.0	78.0	78.0
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	4	80.6	80.2	80.9
	All lots	N/A	278	80.6	76.9	82.5	307	80.1	77.1	83.1	325	80.6	78.0	82.6
Moisture	U.S. No. 1	N/A	126	9.7	8.8	12.0	110	10.1	8.9	12.0	120	10.0	8.5	12.0
	U.S. No. 2	N/A	151	9.5	8.8	11.1	197	9.8	8.6	12.5	200	9.8	9.0	10.9
	U.S. No. 3	N/A	1	9.1	9.1	9.1	--	--	--	--	1	11.5	11.5	11.5
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	4	9.2	9.0	10.0
	All lots	N/A	278	9.6	8.8	12.0	307	9.9	8.6	12.5	325	9.8	8.5	12.0
Heat-damaged Kernels	U.S. No. 1	0.2	126	0.0	0.0	0.0	110	0.0	0.0	0.1	120	0.0	0.0	0.1
	U.S. No. 2	0.2	151	0.0	0.0	0.0	197	0.0	0.0	0.0	200	0.0	0.0	0.0
	U.S. No. 3	0.5	1	0.0	0.0	0.0	--	--	--	--	1	0.0	0.0	0.0
	U.S. No. 4	1.0	--	--	--	--	--	--	--	--	4	0.0	0.0	0.0
	All lots	N/A	278	0.0	0.0	0.0	307	0.0	0.0	0.1	325	0.0	0.0	0.1
Damaged Kernels (Total)	U.S. No. 1	2.0	126	0.1	0.0	0.6	110	0.2	0.0	1.0	120	0.2	0.0	1.5
	U.S. No. 2	4.0	151	0.1	0.0	2.2	197	0.1	0.0	2.2	200	0.1	0.0	2.4
	U.S. No. 3	7.0	1	0.3	0.3	0.3	--	--	--	--	1	0.2	0.2	0.2
	U.S. No. 4	10.0	--	--	--	--	--	--	--	--	4	0.2	0.1	0.4
	All lots	N/A	278	0.1	0.0	2.2	307	0.2	0.0	2.2	325	0.2	0.0	2.4
Foreign Material	U.S. No. 1	0.4	126	0.1	0.0	0.3	110	0.1	0.0	0.3	120	0.1	0.0	0.2
	U.S. No. 2	0.7	151	0.1	0.0	0.3	197	0.1	0.0	0.4	200	0.1	0.0	0.2
	U.S. No. 3	1.3	1	0.1	0.1	0.1	--	--	--	--	1	0.1	0.1	0.1
	U.S. No. 4	3.0	--	--	--	--	--	--	--	--	4	0.0	0.0	0.1
	All lots	N/A	278	0.1	0.0	0.3	307	0.1	0.0	0.4	325	0.1	0.0	0.2
Shrunken and Broken	U.S. No. 1	3.0	126	0.9	0.4	1.4	110	0.9	0.4	1.5	120	0.9	0.4	1.3
	U.S. No. 2	5.0	151	1.0	0.5	1.4	197	1.0	0.5	1.9	200	0.9	0.4	1.7
	U.S. No. 3	8.0	1	0.9	0.9	0.9	--	--	--	--	1	1.1	1.1	1.1
	U.S. No. 4	12.0	--	--	--	--	--	--	--	--	4	1.0	0.9	1.0
	All lots	N/A	278	1.0	0.4	1.4	307	1.0	0.4	1.9	325	0.9	0.4	1.7
Total Defects ¹	U.S. No. 1	3.0	126	1.1	0.4	1.6	110	1.1	0.4	2.2	120	1.1	0.5	2.4
	U.S. No. 2	5.0	151	1.2	0.8	3.2	197	1.2	0.6	3.8	200	1.2	0.6	3.4
	U.S. No. 3	8.0	1	1.3	1.3	1.3	--	--	--	--	1	1.4	1.4	1.4
	U.S. No. 4	12.0	--	--	--	--	--	--	--	--	4	1.2	1.0	1.5
	All lots	N/A	278	1.2	0.4	3.2	307	1.2	0.4	3.8	325	1.2	0.5	3.4

continued

Table 6. Summary of export Soft White wheat quality, 2008-2010, continued

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Dockage	U.S. No. 1	N/A	126	0.3	0.1	0.3	110	0.3	0.1	0.3	120	0.3	0.1	0.5
	U.S. No. 2	N/A	151	0.3	0.1	0.7	197	0.3	0.1	0.8	200	0.3	0.1	0.7
	U.S. No. 3	N/A	1	0.3	0.3	0.3	--	--	--	--	1	0.4	0.4	0.4
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	4	0.5	0.4	0.5
	All lots	N/A	278	0.3	0.1	0.7	307	0.3	0.1	0.8	325	0.3	0.1	0.7
Wheat of Other Classes	U.S. No. 1	3.0	126	0.5	0.0	1.9	110	0.3	0.0	2.1	120	0.3	0.0	1.0
	U.S. No. 2	5.0	151	0.5	0.0	2.8	197	0.4	0.0	1.2	200	0.3	0.0	3.0
	U.S. No. 3	10.0	1	1.1	1.1	1.1	--	--	--	--	1	0.1	0.1	0.1
	U.S. No. 4	10.0	--	--	--	--	--	--	--	--	4	0.5	0.2	1.4
	All lots	N/A	278	0.5	0.0	2.8	307	0.3	0.0	2.1	325	0.3	0.0	3.0
Contrasting Classes	U.S. No. 1	1.0	126	0.4	0.0	1.0	110	0.3	0.0	1.0	120	0.3	0.0	1.0
	U.S. No. 2	2.0	151	0.4	0.0	1.5	197	0.3	0.0	1.1	200	0.3	0.0	1.2
	U.S. No. 3	3.0	1	1.1	1.1	1.1	--	--	--	--	1	0.1	0.1	0.1
	U.S. No. 4	10.0	--	--	--	--	--	--	--	--	4	0.5	0.2	1.4
	All lots	N/A	278	0.4	0.0	1.5	307	0.3	0.0	1.1	325	0.3	0.0	1.4
Protein (as is basis)	U.S. No. 1	N/A	126	10.4	8.3	10.9	110	10.3	8.3	10.8	120	10.1	8.3	10.8
	U.S. No. 2	N/A	148	10.8	9.4	11.9	197	10.9	9.3	12.6	200	10.4	9.3	10.9
	U.S. No. 3	N/A	1	11.7	11.7	11.7	--	--	--	--	1	10.4	10.4	10.4
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	0	--	--	--
	All lots	N/A	275	10.7	8.3	11.9	307	10.8	8.3	12.6	321	10.3	8.3	10.9
Protein (12% moisture)	U.S. No. 1	N/A	126	10.1	8.1	10.5	110	10.1	8.3	10.5	120	9.9	8.3	10.5
	U.S. No. 2	N/A	148	10.5	9.1	11.7	197	10.6	9.2	12.2	200	10.1	9.2	10.6
	U.S. No. 3	N/A	1	11.3	11.3	11.3	--	--	--	--	1	10.3	10.3	10.3
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	0	--	--	--
	All lots	N/A	275	10.4	8.1	11.7	307	10.5	8.3	12.2	321	10.1	8.3	10.6

N/A = Does not apply.

-- = No lots reported in this category.

¹The sum of the component factor averages may not equal the average for this factor due to rounding.

Figure 9. SWH Dockage Distribution, All Grades

**U.S. WHEAT EXPORTED, 2010
DISTRIBUTION FOR DOCKAGE – ALL GRADES
SWH**




Figure 10. SWH Protein Distribution, All Grades

U.S. WHEAT EXPORTED, 2010
DISTRIBUTION FOR PROTEIN (12% M) – ALL GRADES
SWH




Table 7. Summary of export Hard White wheat quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 1	60.0	5	62.6	61.7	63.0	--	--	--	--	--	--	--	--
	U.S. No. 2	58.0	4	62.5	62.4	63.3	1	61.5	61.5	61.5	--	--	--	--
	U.S. No. 4	54.0	3	62.8	62.5	63.1	12	62.8	62.3	63.4	12	62.8	61.8	63.5
	All lots	N/A	12	62.6	61.7	63.3	13	62.7	61.5	63.4	12	62.8	61.8	63.5
Test Weight (kg/hl)	U.S. No. 1	N/A	5	82.3	81.2	82.8	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	4	82.2	82.0	83.2	1	80.9	80.9	80.9	--	--	--	--
	U.S. No. 4	N/A	3	82.6	82.2	82.9	12	82.6	81.8	83.3	12	82.6	81.3	83.4
	All lots	N/A	12	82.3	81.2	83.2	13	82.5	80.9	83.3	12	82.6	81.3	83.4
Moisture	U.S. No. 1	N/A	5	10.2	9.1	10.9	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	4	11.1	9.5	11.4	1	10.1	10.1	10.1	--	--	--	--
	U.S. No. 4	N/A	3	11.4	11.2	11.5	12	11.2	11.1	11.4	12	11.2	10.8	11.4
	All lots	N/A	12	11.0	9.1	11.5	13	11.1	10.1	11.4	12	11.2	10.8	11.4
Heat-damaged Kernels	U.S. No. 1	0.2	5	0.0	0.0	0.0	--	--	--	--	--	--	--	--
	U.S. No. 2	0.2	4	0.0	0.0	0.0	1	0.0	0.0	0.0	--	--	--	--
	U.S. No. 4	1.0	3	0.0	0.0	0.0	12	0.0	0.0	0.0	12	0.0	0.0	0.1
	All lots	N/A	12	0.0	0.0	0.0	13	0.0	0.0	0.0	12	0.0	0.0	0.1
Damaged Kernels (Total)	U.S. No. 1	2.0	5	0.1	0.0	0.3	--	--	--	--	--	--	--	--
	U.S. No. 2	4.0	4	0.5	0.0	0.5	1	0.3	0.3	0.3	--	--	--	--
	U.S. No. 4	10.0	3	1.6	1.2	2.2	12	0.7	0.3	1.3	12	0.9	0.4	1.6
	All lots	N/A	12	0.6	0.0	2.2	13	0.6	0.3	1.3	12	0.9	0.4	1.6
Foreign Material	U.S. No. 1	0.4	5	0.1	0.1	0.1	--	--	--	--	--	--	--	--
	U.S. No. 2	0.7	4	0.1	0.1	0.1	1	0.1	0.1	0.1	--	--	--	--
	U.S. No. 4	3.0	3	0.2	0.1	0.2	12	0.2	0.1	0.3	12	0.2	0.1	0.5
	All lots	N/A	12	0.1	0.1	0.2	13	0.2	0.1	0.3	12	0.2	0.1	0.5
Shrunken and Broken	U.S. No. 1	3.0	5	1.6	1.4	2.1	--	--	--	--	--	--	--	--
	U.S. No. 2	5.0	4	1.8	1.0	1.9	1	2.6	2.6	2.6	--	--	--	--
	U.S. No. 4	12.0	3	1.3	1.3	1.4	12	1.4	1.2	1.8	12	1.6	1.3	2.0
	All lots	N/A	12	1.7	1.0	2.1	13	1.5	1.2	2.6	12	1.6	1.3	2.0
Total Defects ¹	U.S. No. 1	3.0	5	2.0	1.8	2.4	--	--	--	--	--	--	--	--
	U.S. No. 2	5.0	4	2.4	1.1	2.5	1	3.0	3.0	3.0	--	--	--	--
	U.S. No. 4	12.0	3	3.1	2.7	3.7	12	2.2	1.7	2.7	12	2.7	2.2	3.6
	All lots	N/A	12	2.4	1.1	3.7	13	2.3	1.7	3.0	12	2.7	2.2	3.6

continued

Table 7. Summary of export Hard White wheat quality, 2008-2010, continued

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Dockage	U.S. No. 1	N/A	5	0.3	0.3	0.3	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	4	0.5	0.2	0.5	1	0.6	0.6	0.6	--	--	--	--
	U.S. No. 4	N/A	3	0.7	0.7	0.8	12	0.8	0.6	0.9	12	0.9	0.7	1.1
	All lots	N/A	12	0.5	0.2	0.8	13	0.7	0.6	0.9	12	0.9	0.7	1.1
Wheat of Other Classes	U.S. No. 1	3.0	5	1.4	0.0	2.3	--	--	--	--	--	--	--	--
	U.S. No. 2	5.0	4	1.2	0.0	2.6	1	1.5	1.5	1.5	--	--	--	--
	U.S. No. 4	10.0	3	5.2	4.2	8.4	12	3.0	1.6	3.9	12	3.2	2.0	4.9
	All lots	N/A	12	1.8	0.0	8.4	13	2.9	1.5	3.9	12	3.2	2.0	4.9
Contrasting Classes	U.S. No. 1	1.0	5	0.7	0.6	0.9	--	--	--	--	--	--	--	--
	U.S. No. 2	2.0	4	1.2	0.0	1.6	1	1.5	1.5	1.5	--	--	--	--
	U.S. No. 4	10.0	3	5.2	4.2	8.4	12	3.0	1.6	3.9	12	3.2	2.0	4.9
	All lots	N/A	12	1.7	0.0	8.4	13	2.9	1.5	3.9	12	3.2	2.0	4.9
Protein (as is basis)	U.S. No. 1	N/A	5	12.4	11.8	13.6	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	4	10.7	10.5	11.4	1	10.7	10.7	10.7	--	--	--	--
	U.S. No. 4	N/A	3	11.7	11.7	11.8	12	11.8	11.2	12.4	12	11.6	11.0	12.2
	All lots	N/A	12	11.1	10.5	13.6	13	11.7	10.7	12.4	12	11.6	11.0	12.2
Protein (12% moisture)	U.S. No. 1	N/A	5	12.1	11.5	13.2	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	4	10.6	10.4	11.3	1	10.5	10.5	10.5	--	--	--	--
	U.S. No. 4	N/A	3	11.6	11.6	11.7	12	11.7	11.1	12.3	12	11.5	10.9	12.0
	All lots	N/A	12	11.0	10.4	13.2	13	11.6	10.5	12.3	12	11.5	10.9	12.0

N/A = Does not apply.

-- = No lots reported in this category.

¹The sum of the component factor averages may not equal the average for this factor due to rounding.

Table 8. Summary of export Mixed wheat quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of Lots	Avg.	Low	High	No. of Lots	Avg.	Low	High	No. of Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 1	60.0	6	60.2	59.8	60.9	--	--	--	--	--	--	--	--
	U.S. No. 2	58.0	--	--	--	--	1	61.3	61.3	61.3	2	62.6	62.3	62.7
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	59.0	59.0	59.0
	All lots	N/A	6	60.2	59.8	60.9	1	61.3	61.3	61.3	3	62.5	59.0	62.7
Test Weight (kg/hl)	U.S. No. 1	N/A	6	79.1	78.7	80.2	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	--	--	--	--	1	80.6	80.6	80.6	2	82.3	81.9	82.4
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	77.6	77.6	77.6
	All lots	N/A	6	79.1	78.7	80.2	1	80.6	80.6	80.6	3	82.2	77.6	82.4
Moisture	U.S. No. 1	N/A	6	12.1	11.5	12.5	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	--	--	--	--	1	11.9	11.9	11.9	2	11.0	10.0	11.4
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	12.7	12.7	12.7
	All lots	N/A	6	12.1	11.5	12.5	1	11.9	11.9	11.9	3	11.0	10.0	12.7
Heat-damaged Kernels	U.S. No. 1	0.2	6	0.0	0.0	0.0	--	--	--	--	--	--	--	--
	U.S. No. 2	0.2	--	--	--	--	1	0.0	0.0	0.0	2	0.0	0.0	0.0
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	0.0	0.0	0.0
	All lots	N/A	6	0.0	0.0	0.0	1	0.0	0.0	0.0	3	0.0	0.0	0.0
Damaged Kernels (Total)	U.S. No. 1	2.0	6	1.4	0.8	2.3	--	--	--	--	--	--	--	--
	U.S. No. 2	4.0	--	--	--	--	1	1.8	1.8	1.8	2	0.2	0.1	0.3
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	1.1	1.1	1.1
	All lots	N/A	6	1.4	0.8	2.3	1	1.8	1.8	1.8	3	0.2	0.1	1.1
Foreign Material	U.S. No. 1	0.4	6	0.2	0.1	0.4	--	--	--	--	--	--	--	--
	U.S. No. 2	0.7	--	--	--	--	1	0.2	0.2	0.2	2	0.1	0.1	0.1
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	0.3	0.3	0.3
	All lots	N/A	6	0.2	0.1	0.4	1	0.2	0.2	0.2	3	0.1	0.1	0.3
Shrunken and Broken	U.S. No. 1	3.0	6	1.3	1.0	1.5	--	--	--	--	--	--	--	--
	U.S. No. 2	5.0	--	--	--	--	1	1.2	1.2	1.2	2	1.1	0.9	1.2
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	1.2	1.2	1.2
	All lots	N/A	6	1.3	1.0	1.5	1	1.2	1.2	1.2	3	1.1	0.9	1.2
Total Defects ¹	U.S. No. 1	3.0	6	2.9	2.2	3.6	--	--	--	--	--	--	--	--
	U.S. No. 2	5.0	--	--	--	--	1	3.2	3.2	3.2	2	1.4	1.3	1.4
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	2.6	2.6	2.6
	All lots	N/A	6	2.9	2.2	3.6	1	3.2	3.2	3.2	3	1.4	1.3	2.6
Dockage	U.S. No. 1	N/A	6	0.9	0.8	0.9	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	--	--	--	--	1	1.0	1.0	1.0	2	0.5	0.3	0.6
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	0.7	0.7	0.7
	All lots	N/A	6	0.9	0.8	0.9	1	1.0	1.0	1.0	3	0.5	0.3	0.7
Protein (as is basis)	U.S. No. 1	N/A	6	12.7	12.6	12.9	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	--	--	--	--	1	13.2	13.2	13.2	0	--	--	--
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	11.9	11.9	11.9
	All lots	N/A	6	12.7	12.6	12.9	1	13.2	13.2	13.2	1	11.9	11.9	11.9
Protein (12% moisture)	U.S. No. 1	N/A	6	12.7	12.7	12.8	--	--	--	--	--	--	--	--
	U.S. No. 2	N/A	--	--	--	--	1	13.2	13.2	13.2	0	--	--	--
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	1	12.0	12.0	12.0
	All lots	N/A	6	12.7	12.7	12.8	1	13.2	13.2	13.2	1	12.0	12.0	12.0

N/A = Does not apply.

-- = No lots reported in this category.

¹The sum of the component factor averages may not equal the average for this factor due to rounding.

Corn

Corn Grades and Grade Requirements

Corn is divided into three classes: Yellow corn, White Corn, and Mixed corn. There are no subclasses of corn. Each class of corn is divided into five U.S. numerical grades and U.S. Sample grade. Special grades are provided to emphasize the qualities or conditions affecting the value of the corn. These special grades are made a part of the grade designation but do not affect the numerical or Sample grade designation.

U.S. Standards for Corn

Grade:	Min Limits of:	Max Limits of:		
		Damaged Kernels		Broken Corn and Foreign material (percent)
	Test Weight/bu (pounds)	Heat-Damaged kernels (percent)	Damaged kernels total (percent)	
U.S. No. 1	56.0	0.1	3.0	2.0
U.S. No. 2	54.0	0.2	5.0	3.0
U.S. No. 3	52.0	0.5	7.0	4.0
U.S. No. 4	49.0	1.0	10.0	5.0
U.S. No. 5	46.0	3.0	15.0	7.0

U.S. Sample Grade:

U.S. Sample Grade is corn that:

- (a) Does not meet the requirements for grades U.S. No.1, 2, 3, 4, or 5; or
- (b) Contains stones which have an aggregate weight in excess of 0.1 percent of the sample weight, 2 or more pieces of glass, 3 or more crotalaria seeds (*Crotalaria* spp.), 2 or more castor beans (*Ricinus communis* L.), 4 or more particles of an unknown foreign substance(s) or a commonly recognized harmful or toxic substance(s), 8 or more cockleburs (*Xanthium* spp.) or similar seeds singly or in combination, or animal filth in excess of 0.20 percent in 1,000 grams; or
- (c) Has a musty, sour, or commercially objectionable foreign odor; or
- (d) Is heating or otherwise of distinctly low quality.

Corn

Definitions

Test weight (lb/bu) is pounds of grain per Winchester bushel (2,150.42 cubic inches) as determined using an approved device. Test weight is determined before the removal of broken corn and foreign material.

Test weight (kg/hl) is the metric system equivalent to pounds per bushel. Kilograms per hectoliter are calculated by multiplying pounds per bushel by 1.287.

Moisture is the water content of grain as determined by an approved moisture meter. The percentage of moisture in a sample does not affect the numerical grade.

Broken corn is all matter that passes readily through a 12/64-inch round-hole sieve and over a 6/64-inch round-hole sieve. The percentage of broken corn by itself does not affect the numerical grade.

Foreign material is all matter that passes readily through a 6/64-inch round-hole sieve and all matter other than corn that remains on top of the 12/64-inch round-hole sieve. The percentage of foreign material by itself does not affect the numerical grade.

Broken corn and foreign material is all matter that passes readily through a 12/64-inch sieve, and all matter other than corn that remains in the sieved sample.

Damaged kernels (total) are kernels and pieces of corn kernels that are badly ground-damaged, badly weather-damaged, diseased, frost-damaged, germ-damaged, heat-damaged, insect-bored, mold-damaged, sprout-damaged, or otherwise materially damaged.

Heat-damaged kernels are kernels and pieces of corn kernels that are materially discolored by excessive respiration, with dark discoloration extending out of the germ, through the sides, and into the back of the kernel.

Mixed corn is corn that does not meet the color requirements for either of the classes Yellow corn or White corn, and which includes White-capped Yellow corn.

Oil, protein, and starch percentages in corn are determined by an approved near infrared transmittance (NIRT) instrument calibrated to approved methods. Percent corn oil, protein, or starch is reported on a dry matter basis unless other basis is requested. The level of oil, protein, or starch in a sample does not affect the numerical grade.

Table 9. U.S. Corn Exports: Number of lots and quantity exported by class and grade, 2008-2010

Class	Grade	2008		2009		2010	
		No. of Lots	Metric Tons	No. of Lots	Metric Tons	No. of Lots	Metric Tons
Yellow Corn	U.S. No. 1	162	2,107,452	53	877,676	38	579,610
	U.S. No. 2	1,037	21,600,994	971	19,255,649	906	20,766,858
	U.S. No. 3	602	19,439,069	661	19,255,592	696	19,992,886
	U.S. No. 4	4	65,375	4	33,199	13	144,092
	U.S. No. 5	1	508	1	3,414	10	118,312
	U.S. Sample Grade	--	--	1	8,551	9	96,424
	Not Inspected	2	78,505	1	3,026	3	166,325
	All lots	1,808	43,291,903	1,692	39,437,107	1,675	41,864,507
White Corn	U.S. No. 1	17	106,054	12	76,204	9	111,545
	U.S. No. 2	39	676,683	67	757,553	61	627,052
	U.S. No. 3	1	4,316	--	--	3	13,370
	U.S. No. 4	--	--	1	1,324	2	32,378
	Not Inspected	--	--	--	--	--	--
	All lots	57	787,053	80	835,081	75	784,345
All Classes	U.S. No. 1	179	2,213,506	65	953,880	47	691,155
	U.S. No. 2	1,076	22,277,677	1,038	20,013,202	967	21,393,910
	U.S. No. 3	603	19,443,385	661	19,255,592	699	20,006,256
	U.S. No. 4	4	65,375	5	34,523	15	176,470
	U.S. No. 5	1	508	1	3,414	10	118,312
	U.S. Sample Grade	--	--	1	8,551	9	96,424
	Not Inspected	2	78,505	1	3,026	3	166,325
	All lots	1,865	44,078,956	1,772	40,272,188	1,750	42,648,852

-- = No lots reported in the category.

Not inspected = These lots were sold without grade designation.

Table 10. Summary of export Yellow Corn quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 1	56.0	162	57.6	56.1	60.4	53	56.9	56.0	60.1	38	56.9	56.0	61.0
	U.S. No. 2	54.0	1,037	57.4	54.2	60.1	971	56.8	54.1	60.6	905	55.8	54.0	60.7
	U.S. No. 3	52.0	602	56.7	52.5	60.4	661	56.0	52.2	60.5	696	55.5	52.4	59.3
	U.S. No. 4	49.0	4	56.8	51.6	57.7	4	53.9	50.9	58.0	13	55.0	50.0	58.0
	U.S. No. 5	46.0	1	58.8	58.8	58.8	1	57.4	57.4	57.4	10	56.0	48.4	57.5
	U.S. Sample Grade	N/A	--	--	--	--	1	56.9	56.9	56.9	9	54.1	45.5	57.2
	All lots	N/A	1,806	57.1	51.6	60.4	1,691	56.4	50.9	60.6	1,671	55.7	45.5	61.0
Test Weight (kg/hl)	U.S. No. 1	N/A	162	74.1	72.2	77.7	53	73.3	72.0	77.3	38	73.2	72.1	78.5
	U.S. No. 2	N/A	1,037	73.8	69.8	77.3	971	73.1	69.6	78.1	905	71.8	69.5	78.1
	U.S. No. 3	N/A	602	73.0	67.6	77.8	661	72.1	67.2	77.9	696	71.5	67.5	76.3
	U.S. No. 4	N/A	4	73.1	66.5	74.3	4	69.4	65.5	74.7	13	70.8	64.3	74.7
	U.S. No. 5	N/A	1	75.7	75.7	75.7	1	73.9	73.9	73.9	10	72.1	62.2	74.0
	U.S. Sample Grade	N/A	--	--	--	--	1	73.3	73.3	73.3	9	69.6	58.6	73.6
	All lots	N/A	1,806	73.5	66.5	77.8	1,691	72.6	65.5	78.1	1,671	71.7	58.6	78.5
Moisture	U.S. No. 1	N/A	162	13.7	12.2	14.7	53	14.3	13.7	14.8	38	14.3	13.5	14.8
	U.S. No. 2	N/A	1,037	13.7	12.7	14.9	971	14.3	13.2	15.0	904	14.2	12.7	15.6
	U.S. No. 3	N/A	602	14.0	12.8	15.0	661	14.3	13.0	15.4	696	14.3	13.2	15.0
	U.S. No. 4	N/A	4	13.5	13.4	14.3	4	14.1	13.9	14.9	13	14.1	13.5	14.7
	U.S. No. 5	N/A	1	14.2	14.2	14.2	1	15.1	15.1	15.1	10	14.3	12.8	14.7
	U.S. Sample Grade	N/A	--	--	--	--	1	14.8	14.8	14.8	9	13.9	13.6	14.4
	All lots	N/A	1,806	13.8	12.2	15.0	1,691	14.3	13.0	15.4	1,670	14.3	12.7	15.6
Heat-damaged Kernels	U.S. No. 1	0.1	162	0.0	0.0	0.1	53	0.0	0.0	0.0	38	0.0	0.0	0.0
	U.S. No. 2	0.2	1,037	0.0	0.0	0.2	971	0.0	0.0	0.1	906	0.0	0.0	0.1
	U.S. No. 3	0.5	602	0.0	0.0	0.1	661	0.0	0.0	0.1	696	0.0	0.0	0.3
	U.S. No. 4	1.0	4	0.0	0.0	0.0	4	0.0	0.0	0.0	13	0.0	0.0	0.0
	U.S. No. 5	3.0	1	0.0	0.0	0.0	1	0.0	0.0	0.0	10	0.0	0.0	0.0
	U.S. Sample Grade	N/A	--	--	--	--	1	0.0	0.0	0.0	9	0.1	0.0	0.1
	All lots	N/A	1,806	0.0	0.0	0.2	1,691	0.0	0.0	0.1	1,672	0.0	0.0	0.3

continued

Table 10. Summary of export Yellow Corn quality, 2008-2010, continued

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Damaged	U.S. No. 1	3.0	162	1.6	0.3	3.0	53	2.6	0.9	3.0	38	2.5	0.6	3.0
Kernels	U.S. No. 2	5.0	1,037	2.1	0.4	5.0	971	3.3	0.5	5.0	906	3.9	0.0	5.0
(Total)	U.S. No. 3	7.0	602	1.7	0.2	6.3	661	3.4	0.4	7.0	696	4.2	0.4	7.0
	U.S. No. 4	10.0	4	2.1	0.3	8.3	4	3.1	2.6	7.5	13	4.6	0.2	8.9
	U.S. No. 5	15.0	1	10.4	10.4	10.4	1	11.5	11.5	11.5	10	7.1	3.3	15.0
	U.S. Sample Grade	N/A	--	--	--	--	1	5.2	5.2	5.2	9	19.1	3.6	27.0
	All lots	N/A	1,806	1.9	0.2	10.4	1,691	3.4	0.4	11.5	1,672	4.1	0.0	27.0
Broken Corn	U.S. No. 1	2.0	162	1.7	0.9	2.0	53	1.9	1.1	2.0	38	1.9	1.1	2.0
and Foreign	U.S. No. 2	3.0	1,036	2.5	0.7	3.0	971	2.7	1.1	3.0	905	2.7	1.0	3.0
Material	U.S. No. 3	4.0	601	2.9	0.8	4.0	661	3.4	0.9	4.0	696	3.4	0.9	4.0
	U.S. No. 4	5.0	4	2.8	2.2	4.9	4	4.0	1.5	4.3	13	4.3	1.4	4.9
	U.S. No. 5	7.0	1	1.9	1.9	1.9	1	1.6	1.6	1.6	10	3.8	1.4	4.9
	U.S. Sample Grade	N/A	--	--	--	--	1	3.3	3.3	3.3	9	3.7	1.5	7.4
	All lots	N/A	1,804	2.6	0.7	4.9	1,691	3.0	0.9	4.3	1,671	3.0	0.9	7.4
Broken Corn	U.S. No. 1	N/A	19	1.3	1.1	1.5	26	1.4	1.1	1.6	16	1.4	1.3	1.6
	U.S. No. 2	N/A	154	1.7	1.1	2.2	92	1.9	1.1	2.3	66	2.0	1.1	3.0
	U.S. No. 3	N/A	3	2.7	2.5	2.9	--	--	--	--	--	--	--	--
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	--	--	--	--
	U.S. No. 5	N/A	--	--	--	--	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	--	--	--	--
	All lots	N/A	176	1.7	1.1	2.9	118	1.8	1.1	2.3	82	1.8	1.1	3.0
Foreign	U.S. No. 1	N/A	19	0.5	0.4	0.7	26	0.5	0.4	0.7	16	0.5	0.3	0.6
Material	U.S. No. 2	N/A	167	0.7	0.3	1.1	95	0.7	0.4	1.1	67	0.7	0.4	1.2
	U.S. No. 3	N/A	3	0.4	0.2	0.9	--	--	--	--	--	--	--	--
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	--	--	--	--
	U.S. No. 5	N/A	--	--	--	--	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	--	--	--	--
	All lots	N/A	189	0.7	0.2	1.1	121	0.7	0.4	1.1	83	0.7	0.3	1.2

N/A = Does not apply.

-- = No lots reported in this category.

Table 11. Summary of export White Corn quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 1	56.0	17	58.8	57.2	60.3	12	59.2	56.1	59.7	9	59.4	58.7	60.2
	U.S. No. 2	54.0	39	59.0	57.9	60.5	67	59.4	58.2	61.0	61	59.0	57.9	60.0
	U.S. No. 3	52.0	1	59.3	59.3	59.3	--	--	--	--	3	58.0	52.8	58.9
	U.S. No. 4	49.0	--	--	--	--	1	57.9	57.9	57.9	2	57.8	57.7	58.5
	All lots	N/A	57	59.0	57.2	60.5	80	59.4	56.1	61.0	75	59.0	52.8	60.2
Test Weight (kg/hl)	U.S. No. 1	N/A	17	75.7	73.6	77.6	12	76.2	72.2	76.9	9	76.4	75.6	77.5
	U.S. No. 2	N/A	39	76.0	74.6	77.9	67	76.5	74.9	78.5	61	76.0	74.6	77.2
	U.S. No. 3	N/A	1	76.4	76.4	76.4	--	--	--	--	3	74.6	68.0	75.8
	U.S. No. 4	N/A	--	--	--	--	1	74.5	74.5	74.5	2	74.4	74.3	75.3
	All lots	N/A	57	75.9	73.6	77.9	80	76.5	72.2	78.5	75	75.9	68.0	77.5
Moisture	U.S. No. 1	N/A	17	13.6	12.3	14.4	12	13.8	13.6	14.3	9	13.9	13.2	14.6
	U.S. No. 2	N/A	39	13.6	12.7	14.4	67	14.0	13.2	14.7	61	14.0	13.4	14.4
	U.S. No. 3	N/A	1	13.7	13.7	13.7	--	--	--	--	2	14.0	14.0	14.0
	U.S. No. 4	N/A	--	--	--	--	1	14.2	14.2	14.2	2	14.2	14.0	14.2
	All lots	N/A	57	13.6	12.3	14.4	80	14.0	13.2	14.7	74	14.0	13.2	14.6
Heat-damaged Kernels	U.S. No. 1	0.1	17	0.0	0.0	0.0	12	0.0	0.0	0.1	9	0.0	0.0	0.0
	U.S. No. 2	0.2	39	0.0	0.0	0.0	67	0.0	0.0	0.0	61	0.0	0.0	0.1
	U.S. No. 3	0.5	1	0.0	0.0	0.0	--	--	--	--	3	0.0	0.0	0.0
	U.S. No. 4	1.0	--	--	--	--	1	0.0	0.0	0.0	2	0.0	0.0	0.0
	All lots	N/A	57	0.0	0.0	0.0	80	0.0	0.0	0.1	75	0.0	0.0	0.1

continued

Table 11. Summary of export White Corn quality, 2008-2010, continued

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Damaged	U.S. No. 1	3.0	17	1.7	0.7	3.0	12	1.5	1.1	2.2	9	1.3	0.6	1.7
Kernels	U.S. No. 2	5.0	39	1.4	0.4	3.5	67	1.9	0.6	4.4	61	2.8	0.7	4.9
(Total)	U.S. No. 3	7.0	1	0.9	0.9	0.9	--	--	--	--	3	3.3	2.5	5.8
	U.S. No. 4	10.0	--	--	--	--	1	2.0	2.0	2.0	2	3.1	2.5	3.2
	All lots	N/A	57	1.4	0.4	3.5	80	1.9	0.6	4.4	75	2.6	0.6	5.8
Broken Corn	U.S. No. 1	2.0	17	1.8	1.2	2.0	12	1.8	1.2	1.9	9	1.4	1.2	1.9
and Foreign	U.S. No. 2	3.0	39	2.2	1.2	2.9	67	2.5	0.9	3.0	61	2.4	1.5	3.0
Material	U.S. No. 3	4.0	1	3.7	3.7	3.7	--	--	--	--	3	3.3	2.6	3.5
	U.S. No. 4	5.0	--	--	--	--	1	4.5	4.5	4.5	2	4.5	4.3	4.5
	All lots	N/A	57	2.2	1.2	3.7	80	2.4	0.9	4.5	75	2.3	1.2	4.5
Broken Corn	U.S. No. 1	N/A	6	1.2	1.0	1.5	4	1.4	1.3	1.4	--	--	--	--
	U.S. No. 2	N/A	11	1.7	1.2	2.1	14	2.0	1.7	2.2	15	1.9	1.5	2.4
	U.S. No. 3	N/A	--	--	--	--	--	--	--	--	--	--	--	--
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	1	2.1	2.1	2.1
	All lots	N/A	17	1.6	1.0	2.1	18	1.9	1.3	2.2	16	2.0	1.5	2.4
Foreign	U.S. No. 1	N/A	6	0.5	0.4	0.6	4	0.5	0.4	0.6	--	--	--	--
Material	U.S. No. 2	N/A	11	0.7	0.6	0.9	14	0.7	0.6	0.9	15	0.5	0.4	0.7
	U.S. No. 3	N/A	--	--	--	--	--	--	--	--	--	--	--	--
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	1	2.4	2.4	2.4
	All lots	N/A	17	0.6	0.4	0.9	18	0.7	0.4	0.9	16	0.8	0.4	2.4

N/A = Does not apply.

-- = No lots reported in this category.

Figure 11. Corn BCFM Distribution, Grade 2

**U.S. CORN EXPORTED, 2010
DISTRIBUTION FOR BCFM – GRADE 2**





Figure 12. Corn BCFM Distribution, Grade 3

**U.S. CORN EXPORTED, 2010
DISTRIBUTION FOR BCFM – GRADE 3**



Soybeans

Soybean Grades and Grade Requirements

There are two classes of soybeans: Yellow soybeans and Mixed soybeans. There are no soybean subclasses. The class Yellow soybeans is the class most commonly exported by the U.S. market. Each class is divided into four U.S. numerical grades and U.S. Sample grade.

Special grades are provided to emphasize special qualities or conditions affecting the value of the soybeans. These special grades are a part of the grade designation but do not affect the numerical or Sample grade designation.

U.S. Standards for Soybeans

Grade:	Max Limits of:				
	Damaged Kernels		Foreign Material (percent)	Splits (percent)	Soybeans of other colors <u>1</u> / (percent)
	Heat (part of total) (percent)	Total (percent)			
U.S. No. 1	0.2	2.0	1.0	10.0	1.0
U.S. No. 2	0.5	3.0	2.0	20.0	2.0
U.S. No. 3	1.0	5.0	3.0	30.0	5.0
U.S. No. 4	3.0	8.0	5.0	40.0	10.0

U.S. Sample Grade:

U.S. Sample grade shall be soybeans which:

- (a) Do not meet the requirements for U.S. Nos. 1, 2, 3, or 4; or
- (b) Contains 4 or more stones which have an aggregate weight in excess of 0.1 percent of the sample weight, 1 or more pieces of broken glass, 3 or more crotalaria seeds, 2 or more castor beans, 4 or more particles of an unknown foreign substance(s) or a commonly recognized harmful or toxic foreign substance(s), 10 or more rodent pellets, bird droppings, or an equivalent quantity of other animal filth in 1,000 grams of soybeans; or
- (c) Contains 11 or more pieces, in any combination, of animal filth, castor beans, crotalaria seeds, glass, stones, or unknown foreign substances.
- (d) Have a musty, sour, or commercially objectionable foreign odor (except garlic odor); or
- (e) Are heating or otherwise of distinctly low quality.

1/ Disregard for Mixed Soybeans.

Soybeans

Definitions

Test weight (lb/bu) is pounds of grain per Winchester bushel determined by an approved device and is recorded to the nearest tenth pound for soybeans.

Test weight (kg/hl) is the metric system equivalent to pounds per bushel. Kilograms per hectoliter are calculated by multiplying pounds per bushel by 1.287.

Moisture is the water content of grain as determined by an approved electronic moisture meter. The percentage of moisture in a sample does not affect the numerical grade.

Splits are soybeans with more than one-fourth of the bean removed and which are not damaged.

Damaged kernels are soybeans and pieces of soybeans which are badly ground-damaged, badly weather-damaged, diseased, frost-damaged, heat-damaged, insect-bored, mold-damaged, sprout damaged, stinkbug-stung, or otherwise materially damaged.

Heat-damaged kernels are soybeans and pieces of soybeans which are materially discolored and damaged by heat.

Foreign material is all matter, including soybeans and pieces of soybeans that will pass readily through an 8/64-inch sieve and all matter other than soybeans remaining on the sieve after sieving.

Soybeans of other colors are soybeans which have green, black, brown, or bicolored seed coats. Before September 9, 1985, this factor was called "brown, black, and/or bicolored soybeans in yellow or green soybeans."

Mixed soybeans is a combination of classes of soybeans which does not meet the minimum requirements of the class Yellow soybeans.

Protein and oil percentages in soybeans are determined by an approved near infrared transmittance (NIRT) instrument calibrated to approved methods. Percent protein and oil is reported on a 13 percent moisture basis unless another basis is requested. The level of protein and oil in a sample does not affect the numerical grade.

Table 12. U.S. Soybean Exports: Number of lots and quantity exported by grade, 2008-2010

Grade	2008		2009		2010	
	No. of Lots	Metric Tons	No. of Lots	Metric Tons	No. of Lots	Metric Tons
U.S. No. 1	16	117,125	20	303,270	26	532,098
U.S. No. 2	792	24,698,359	995	35,135,738	1,029	36,215,801
U.S. No. 3	17	341,769	15	307,884	11	192,479
U.S. No. 4	8	240,536	2	45,111	1	8,811
U.S. Sample Grade	42	1,787,135	10	389,666	15	654,218
Not Inspected	2	131,615	1	57,750	1	62,995
All lots	877	27,316,539	1,043	36,239,419	1,083	37,666,402

-- = No lots reported in the category.

Not inspected = These lots were sold without grade designation.

Table 13. Summary of export Soybean quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 1	N/A	16	55.8	55.1	57.3	20	56.0	55.5	58.0	26	56.4	55.4	57.5
	U.S. No. 2	N/A	792	54.5	0.0	57.6	995	54.3	0.0	58.1	993	55.9	54.1	58.6
	U.S. No. 3	N/A	17	53.0	0.0	56.9	15	39.5	0.0	57.7	5	55.3	52.5	56.1
	U.S. No. 4	N/A	8	54.9	53.0	55.3	2	53.1	53.0	53.3	1	55.1	55.1	55.1
	U.S. Sample Grade	N/A	42	54.4	53.0	55.4	10	54.3	53.1	55.1	15	54.8	52.6	56.1
	All lots	N/A	875	54.4	0.0	57.6	1042	54.2	0.0	58.1	1,040	55.8	52.5	58.6
Test Weight (kg/hl)	U.S. No. 1	N/A	16	71.9	71.0	73.8	20	72.1	71.4	74.7	26	72.6	71.3	74.1
	U.S. No. 2	N/A	792	70.1	0.0	74.1	995	69.9	0.0	74.8	993	71.9	69.6	75.5
	U.S. No. 3	N/A	17	68.2	0.0	73.3	15	50.8	0.0	74.3	5	71.1	67.6	72.2
	U.S. No. 4	N/A	8	70.6	68.3	71.2	2	68.4	68.2	68.6	1	70.9	70.9	70.9
	U.S. Sample Grade	N/A	42	70.0	68.2	71.3	10	70.0	68.4	70.9	15	70.5	67.7	72.3
	All lots	N/A	875	70.1	0.0	74.1	1042	69.8	0.0	74.8	1,040	71.9	67.6	75.5
Moisture	U.S. No. 1	N/A	16	11.4	10.6	12.5	20	12.5	10.6	13.8	26	12.3	10.7	13.4
	U.S. No. 2	N/A	790	11.6	9.8	13.0	995	12.3	9.9	14.0	1,026	11.7	9.5	13.3
	U.S. No. 3	N/A	17	12.1	11.0	13.6	15	12.5	10.1	13.2	11	12.0	11.0	12.9
	U.S. No. 4	N/A	8	11.5	10.9	13.7	2	13.5	13.4	13.7	1	12.5	12.5	12.5
	U.S. Sample Grade	N/A	42	11.7	10.7	13.9	10	13.0	11.7	13.8	15	12.4	10.5	13.2
	All lots	N/A	873	11.6	9.8	13.9	1042	12.3	9.9	14.0	1,079	11.7	9.5	13.4
Heat-damaged Kernels	U.S. No. 1	0.2	16	0.0	0.0	0.0	20	0.0	0.0	0.1	26	0.0	0.0	0.1
	U.S. No. 2	0.5	792	0.1	0.0	0.5	995	0.1	0.0	0.5	1,029	0.1	0.0	0.5
	U.S. No. 3	1.0	17	0.3	0.0	0.8	15	0.2	0.0	0.4	11	0.3	0.0	0.6
	U.S. No. 4	3.0	8	0.5	0.3	0.5	2	0.2	0.1	0.3	1	0.5	0.5	0.5
	U.S. Sample Grade	N/A	42	0.7	0.1	1.3	10	0.9	0.4	1.4	15	0.6	0.1	2.1
	All lots	N/A	875	0.2	0.0	1.3	1042	0.1	0.0	1.4	1,082	0.1	0.0	2.1
Damaged Kernels (Total)	U.S. No. 1	2.0	16	0.7	0.1	1.1	20	0.7	0.1	1.3	26	0.7	0.2	1.0
	U.S. No. 2	3.0	792	1.3	0.1	3.0	995	1.5	0.1	3.0	1,029	1.4	0.0	3.2
	U.S. No. 3	5.0	17	2.8	0.2	4.9	15	2.9	0.1	5.0	11	2.8	0.8	4.8
	U.S. No. 4	8.0	8	3.0	1.9	7.7	2	7.2	6.9	7.7	1	5.5	5.5	5.5
	U.S. Sample Grade	N/A	42	4.5	1.9	10.8	10	8.0	4.8	9.2	15	5.5	0.8	13.2
	All lots	N/A	875	1.5	0.1	10.8	1042	1.6	0.1	9.2	1,082	1.5	0.0	13.2

continued

Table 13. Summary of export Soybean quality, 2008-2010,
continued

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Foreign	U.S. No. 1	1.0	16	0.9	0.3	1.0	20	0.8	0.5	1.0	26	0.8	0.2	1.0
Material	U.S. No. 2	2.0	792	1.5	0.4	2.0	995	1.3	0.4	2.0	1,029	1.4	0.0	2.0
	U.S. No. 3	3.0	17	1.7	1.2	2.1	15	1.3	0.8	2.4	11	1.3	1.1	1.8
	U.S. No. 4	5.0	8	1.7	1.5	2.0	2	1.3	1.1	1.6	1	1.5	1.5	1.5
	U.S. Sample Grade	N/A	42	2.0	1.3	3.0	10	1.6	1.3	2.0	15	1.7	1.1	2.2
	All lots	N/A	875	1.5	0.3	3.0	1042	1.3	0.4	2.4	1,082	1.4	0.0	2.2
Splits	U.S. No. 1	10.0	16	7.7	2.0	9.7	20	5.9	2.9	8.7	26	6.6	3.1	8.6
	U.S. No. 2	20.0	792	8.7	1.9	18.7	995	7.9	2.3	19.2	1,029	7.2	0.0	17.1
	U.S. No. 3	30.0	17	8.1	4.3	24.0	15	6.8	3.7	23.4	11	7.5	3.8	21.3
	U.S. No. 4	40.0	8	9.9	5.3	11.7	2	6.5	6.1	7.0	1	7.7	7.7	7.7
	U.S. Sample Grade	N/A	42	10.0	4.9	14.6	10	5.9	4.3	12.1	15	7.8	4.8	10.2
	All lots	N/A	875	8.7	1.9	24.0	1042	7.9	2.3	23.4	1,082	7.2	0.0	21.3
Soybeans of Other Colors	U.S. No. 1	1.0	16	0.0	0.0	0.2	20	0.0	0.0	0.2	26	0.0	0.0	0.2
	U.S. No. 2	2.0	792	0.0	0.0	0.7	995	0.0	0.0	0.5	1,029	0.0	0.0	0.4
	U.S. No. 3	5.0	17	0.1	0.0	0.2	15	0.1	0.0	0.1	11	0.0	0.0	0.2
	U.S. No. 4	10.0	8	0.0	0.0	0.1	2	0.1	0.0	0.1	1	0.0	0.0	0.0
	U.S. Sample Grade	N/A	42	0.1	0.0	0.2	10	0.0	0.0	0.1	15	0.0	0.0	0.1
	All lots	N/A	875	0.0	0.0	0.7	1042	0.0	0.0	0.5	1,082	0.0	0.0	0.4
Protein (adjusted to 13% moisture)	U.S. No. 1	N/A	10	35.2	34.2	36.9	15	35.2	34.7	35.9	20	35.6	35.1	36.6
	U.S. No. 2	N/A	416	34.9	33.4	38.7	503	34.7	32.7	38.0	514	34.9	33.2	39.9
	U.S. No. 3	N/A	6	35.3	34.4	38.6	3	35.7	33.9	36.1	--	--	--	--
	U.S. No. 4	N/A	6	35.3	35.2	35.4	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	33	35.2	34.5	35.7	10	35.2	35.1	35.5	12	35.2	34.9	35.5
	All lots	N/A	471	34.9	33.4	38.7	531	34.7	32.7	38.0	546	34.9	33.2	39.9
Oil (adjusted to 13% moisture)	U.S. No. 1	N/A	11	19.0	17.8	20.0	15	18.7	18.2	19.6	20	18.6	18.0	19.3
	U.S. No. 2	N/A	418	19.1	17.2	20.0	510	18.8	16.7	20.4	522	18.9	16.7	20.1
	U.S. No. 3	N/A	6	19.0	17.3	19.4	3	19.8	18.2	20.1	--	--	--	--
	U.S. No. 4	N/A	6	19.1	19.0	19.3	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	33	19.6	18.9	20.6	10	19.2	18.6	19.9	12	18.8	18.5	19.6
	All lots	N/A	474	19.2	17.2	20.6	538	18.8	16.7	20.4	554	18.8	16.7	20.1

N/A = Does not apply.

-- = No lots reported in this category.

Figure 13. Soybean FM Distribution, Grade 2

**U.S. SOYBEANS EXPORTED, 2010
DISTRIBUTION FOR FM – GRADE 2**




Figure 14. Soybean Protein Distribution, All Grades

U.S. SOYBEANS EXPORTED, 2010
DISTRIBUTION FOR PROTEIN (13% M) – ALL GRADES





Figure 15. Soybean Oil Distribution, All Grades

**U.S. SOYBEANS EXPORTED, 2010
DISTRIBUTION FOR OIL (13% M) – ALL GRADES**



Other Grain Exports

Sorghum

Sorghum Grades and Grade Requirements

Sorghum is divided into four classes: Sorghum, Tannin sorghum, White sorghum, and Mixed sorghum. There are no subclasses in sorghum. Each class is divided into four numerical grades and U.S. Sample grade. Special grades are provided to emphasize special qualities or conditions affecting the value of sorghum. Special grades are added to and made a part of the grade designation. They do not affect the numerical or Sample grade designation.

U.S. Standards for Sorghum

Grade:	Min Limits of:	Max Limits of:			
	Test weight per bushel (pounds)	Damaged Kernels		Broken kernels and foreign material	
		Heat (percent)	Total (percent)	Foreign material (percent)	Total (percent)
U.S. No. 1	57.0	0.2	2.0	1.0	3.0
U.S. No. 2	55.0	0.5	5.0	2.0	6.0
U.S. No. 3 <u>1</u> /	53.0	1.0	10.0	3.0	8.0
U.S. No. 4	51.0	3.0	15.0	4.0	10.0

U.S. Sample Grade:

U.S. Sample grade shall be sorghum which:

- (a) Do not meet the requirements for U.S. Nos. 1, 2, 3, or 4; or
- (b) Contains 8 or more stones which have an aggregate weight in excess of 0.2 percent of the sample weight, 2 or more pieces of glass, 3 or more crotalaria seeds (*Crotalaria* spp.), 2 or more castor beans (*Ricinus communis* L.), 4 or more particles of an unknown foreign substance(s) or a commonly recognized harmful or toxic substance(s), 8 or more cockleburs (*Xanthium* spp.) or similar seeds singly or in combination, 10 or more rodent pellets, bird droppings, or equivalent quantity of other animal filth; or
- (c) Have a musty, sour, or commercially objectionable foreign odor (except smut odor); or
- (d) Is badly weathered, heating, or otherwise of distinctly low quality.

1/ Sorghum which is distinctly discolored shall not grade any higher than U.S. No. 3.

Sorghum

Definitions

Test weight (lb/bu) is pounds of grain per Winchester bushel (2.150.42 cubic inches) as determined using an approved device before the removal of dockage.

Test weight (kg/hl) is the metric system equivalent to pounds per bushel. Kilograms per hectoliter are calculated by multiplying pounds per bushel by 1.287.

Heat-damaged kernels are kernels, pieces of sorghum kernels, and other grains that are materially discolored and damaged by heat.

Damaged kernels are kernels, pieces of sorghum kernels, and other grains that are badly ground-damaged, badly weather-damaged, diseased, frost-damaged, germ-damaged, heat-damaged, insect-bored, mold-damaged, sprout-damaged, or otherwise materially damaged.

Foreign material is all matter except sorghum, which passes over the number 6 riddle and all matter other than sorghum that remains on the top of the 5/64 triangular-hole sieve.

Broken kernels are all matter which passes through a 5/64 triangular-hole sieve and over a 2.5/64 round-hole sieve.

Broken kernels and foreign material consists of the combination of broken kernels and foreign material.

Dockage is all matter other than sorghum that can be removed from the original sample by use of an approved device. Also, underdeveloped, shriveled, and small pieces of sorghum kernels removed in properly separating the material other than sorghum.

Moisture is the water content of grain as determined by an approved moisture meter. The percentage of moisture in a sample does not affect the numerical grade.

Mixed sorghum is a sorghum which does not meet the minimum requirements for any of the classes of sorghum, Tannin sorghum or White sorghum.

Table 14. U.S. Sorghum Exports: Number of lots and quantity exported by class and grade, 2008-2010

Class	Grade	2008		2009		2010	
		No. of Lots	Metric Tons	No. of Lots	Metric Tons	No. of Lots	Metric Tons
Sorghum	U.S. No. 2	199	4,055,944	163	2,542,005	192	2,745,431
	U.S. No. 3	2	104,505	3	17,553	9	351,380
	U.S. No. 4	1	33,554	--	--	--	--
	U.S. Sample Grade	1	60,003	--	--	--	--
	Not Inspected	4	226,848	1	2,622	--	--
	All lots	207	4,480,854	167	2,562,180	201	3,096,811
All Classes	U.S. No. 2	199	4,055,944	163	2,542,005	192	2,745,431
	U.S. No. 3	2	104,505	3	17,553	9	351,380
	U.S. No. 4	1	33,554	--	--	--	--
	U.S. Sample Grade	1	60,003	--	--	--	--
	Not Inspected	4	226,848	1	2,622	--	--
	All lots	207	4,480,854	167	2,562,180	201	3,096,811

-- = No lots reported in the category.

Not inspected = These lots were sold without grade designation.

Table 15. Summary of export Sorghum quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 2	55.0	199	58.7	55.8	60.1	163	58.0	56.4	59.9	192	57.7	55.3	60.1
	U.S. No. 3	53.0	2	58.7	58.5	58.8	3	58.5	57.8	58.9	9	57.4	55.8	58.3
	U.S. No. 4	51.0	1	55.9	55.9	55.9	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	1	59.1	59.1	59.1	--	--	--	--	--	--	--	--
	All lots	N/A	203	58.7	55.8	60.1	166	58.0	56.4	59.9	201	57.7	55.3	60.1
Test Weight (kg/hl)	U.S. No. 2	N/A	199	75.5	71.9	77.3	163	74.7	72.6	77.1	192	74.3	71.2	77.4
	U.S. No. 3	N/A	2	75.5	75.3	75.7	3	75.4	74.4	75.8	9	74.0	71.8	75.0
	U.S. No. 4	N/A	1	71.9	71.9	71.9	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	1	76.1	76.1	76.1	--	--	--	--	--	--	--	--
	All lots	N/A	203	75.5	71.9	77.3	166	74.7	72.6	77.1	201	74.2	71.2	77.4
Moisture	U.S. No. 2	N/A	199	13.4	12.1	14.0	163	13.5	12.3	14.0	192	13.4	12.4	14.1
	U.S. No. 3	N/A	2	13.1	13.0	13.2	3	14.0	13.8	14.2	9	13.1	12.6	13.3
	U.S. No. 4	N/A	1	13.1	13.1	13.1	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	1	13.5	13.5	13.5	--	--	--	--	--	--	--	--
	All lots	N/A	203	13.3	12.1	14.0	166	13.5	12.3	14.2	201	13.4	12.4	14.1
Heat-damaged Kernels	U.S. No. 2	0.5	199	0.0	0.0	0.1	163	0.0	0.0	0.1	192	0.0	0.0	0.1
	U.S. No. 3	1.0	2	0.0	0.0	0.0	3	0.1	0.0	0.1	9	0.0	0.0	0.0
	U.S. No. 4	3.0	1	0.0	0.0	0.0	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	1	0.0	0.0	0.0	--	--	--	--	--	--	--	--
	All lots	N/A	203	0.0	0.0	0.1	166	0.0	0.0	0.1	201	0.0	0.0	0.1
Damaged Kernels (Total)	U.S. No. 2	5.0	199	1.9	0.2	4.7	163	1.3	0.2	4.5	192	1.9	0.2	5.0
	U.S. No. 3	10.0	2	3.6	2.7	5.1	3	1.9	1.3	2.3	9	6.5	2.7	9.4
	U.S. No. 4	15.0	1	9.1	9.1	9.1	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	1	1.5	1.5	1.5	--	--	--	--	--	--	--	--
	All lots	N/A	203	2.0	0.2	9.1	166	1.3	0.2	4.5	201	2.4	0.2	9.4

continued

Table 15. Summary of export Sorghum quality, 2008-2010, continued

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Broken Kernels and Foreign Material	U.S. No. 2	6.0	199	3.1	1.5	5.8	163	3.2	1.0	5.0	192	3.4	2.0	5.1
	U.S. No. 3	8.0	2	3.1	3.1	3.2	3	4.0	3.7	4.2	9	3.1	2.5	3.6
	U.S. No. 4	10.0	1	4.7	4.7	4.7	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	1	3.6	3.6	3.6	--	--	--	--	--	--	--	--
	All lots	N/A	203	3.1	1.5	5.8	166	3.2	1.0	5.0	201	3.4	2.0	5.1
Broken Kernels	U.S. No. 2	N/A	--	--	--	--	--	--	--	--	2	2.6	2.5	2.6
	U.S. No. 3	N/A	--	--	--	--	--	--	--	--	--	--	--	--
	U.S. No. 4	N/A	--	--	--	--	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	--	--	--	--	--	--	--	--	--	--	--	--
	All lots	N/A	--	--	--	--	--	--	--	--	2	2.6	2.5	2.6
Foreign Material	U.S. No. 2	2.0	199	1.0	0.0	2.4	163	1.2	0.3	2.0	192	1.2	0.0	2.0
	U.S. No. 3	3.0	2	0.9	0.8	0.9	3	2.6	2.5	2.8	9	0.9	0.7	1.4
	U.S. No. 4	4.0	1	1.6	1.6	1.6	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	1	1.2	1.2	1.2	--	--	--	--	--	--	--	--
	All lots	N/A	203	1.0	0.0	2.4	166	1.3	0.3	2.8	201	1.2	0.0	2.0
Dockage	U.S. No. 2	N/A	198	0.2	0.1	0.4	162	0.2	0.1	0.5	186	0.2	0.1	0.6
	U.S. No. 3	N/A	2	0.2	0.2	0.2	3	0.4	0.2	0.9	9	0.2	0.1	0.3
	U.S. No. 4	N/A	1	0.3	0.3	0.3	--	--	--	--	--	--	--	--
	U.S. Sample Grade	N/A	1	0.2	0.2	0.2	--	--	--	--	--	--	--	--
	All lots	N/A	202	0.2	0.1	0.4	165	0.2	0.1	0.9	195	0.2	0.1	0.6

N/A = Does not apply.

-- = No lots reported in this category.

Figure 16. Sorghum BNFM Distribution, Grade 2

U.S. SORGHUM EXPORTED, 2010
DISTRIBUTION FOR BNFM – GRADE 2





Figure 17. Sorghum FM Distribution, Grade 2

U.S. SORGHUM EXPORTED, 2010
DISTRIBUTION FOR FM – GRADE 2



Barley

Barley Grades and Grade Requirements

Barley is divided into two classes: Malting Barley and Barley. The class Malting Barley is divided into three subclasses: Six-Rowed Malting Barley, Six-Rowed Blue Malting Barley, and Two-Rowed Malting Barley. The class Barley is divided into three subclasses: Six Rowed Barley, Two-Rowed Barley, and Barley. The applicant for service may request either the malting standards or barley standards for malting types.

Grades and grade requirements for Six-rowed Malting barley and Six-rowed Blue Malting barley

Grade	Minimum Limits of:			Maximum Limits of -				
	Test weight per bushel (pounds)	Suitable malting type (percent)	Sound barley <u>1/</u> (percent)	Damaged kernels <u>1/</u> (percent)	Foreign material (percent)	Other grains (percent)	Skinned and broken kernels (percent)	Thin Barley <u>2/</u> (percent)
U.S. No. 1	47.0	95.0	97.0	2.0	0.5	2.0	4.0	7.0
U.S. No. 2	45.0	95.0	94.0	3.0	1.0	3.0	6.0	10.0
U.S. No. 3	43.0	95.0	90.0	4.0	2.0	5.0	8.0	15.0
U.S. No. 4	43.0	95.0	87.0	5.0	3.0	5.0	10.0	15.0

1/ Injured-by-frost kernels and injured-by-mold kernels are not considered damaged kernels or considered against sound barley.

2/ Using a 5.5/64 x 3/4 slotted-hole sieve.

NOTE: Malting barley shall not be infested, blighted, ergoty, garlicky, smutty, or contain any special grades. Upon request, malting barley varieties may be inspected and graded in accordance with standards established for the class Barley.

Grades and grade requirements for Two-rowed Malting barley

Grade	Minimum Limits of:			Maximum Limits of -			
	Test weight per bushel (pounds)	Suitable malting type (percent)	Sound barley <u>1/</u> (percent)	Wild Oats (percent)	Foreign material (percent)	Skinned and broken kernels (percent)	Thin Barley <u>2/</u> (percent)
U.S. No. 1	50.0	97.0	98.0	1.0	0.5	5.0	5.0
U.S. No. 2	48.0	97.0	98.0	1.0	1.0	7.0	7.0
U.S. No. 3	48.0	95.0	96.0	2.0	2.0	10.0	10.0
U.S. No. 4	48.0	95.0	93.0	3.0	3.0	10.0	10.0

1/ Injured-by-frost kernels and injured-by-mold kernels are not considered damaged kernels or considered against sound barley.

2/ Using a 5.5/64 x 3/4 slotted-hole sieve.

NOTE: Malting barley shall not be infested, blighted, ergoty, garlicky, smutty, or contain any special grades. Upon request, malting barley varieties may be inspected and graded in accordance with standards established for the class Barley.

Two-rowed barley that does not meet the requirements for U.S. Nos. 1, 2, 3, or 4 Malting shall be graded under the Barley standards.

Grades and grade requirements for Barley

Grade	Minimum Limits of:		Maximum Limits of -				
	Test weight per bushel (pounds)	Sound barley (percent)	Damaged kernels <u>1/</u> (percent)	Heat damaged kernels (percent)	Foreign material (percent)	Broken kernels (percent)	Thin Barley <u>2/</u> (percent)
U.S. No. 1	47.0	97.0	2.0	0.2	1.0	4.0	10.0
U.S. No. 2	45.0	94.0	4.0	0.3	2.0	8.0	15.0
U.S. No. 3	43.0	90.0	6.0	0.5	3.0	12.0	25.0
U.S. No. 4	40.0	85.0	8.0	1.0	4.0	18.0	35.0
U.S. No. 5	36.0	75.0	10.0	3.0	5.0	28.0	75.0

U.S. Sample Grade:

U.S. Sample Grade is wheat that:

- (a) Does not meet the requirements for grades U.S. No. 1, 2, 3, 4, or 5; or
- (b) Contains 8 or more stones or any number of stones which have an aggregate weight in excess of 0.2 percent of the sample weight, 2 or more pieces of glass, 3 or more crotalaria seeds (*Crotalaria* spp.), 2 or more castor beans (*Ricinus communis* L.), 4 or more particles of an unknown foreign substance(s) or a commonly recognized harmful or toxic substance(s), 8 or more cocklebur (*Xanthium* spp.) or similar seeds singly or in combination, 10 or more rodent pellets, bird droppings, or equivalent quantity of other animal filth per 1-1/8 to 1-1/4 quarts of barley; or
- (c) Has a musty, sour, or commercially objectionable foreign odor (except smut or garlic odor); or
- (d) Is heating or otherwise of distinctly low quality.

1/ Injured-by-frost kernels and injured-by-mold kernels are not considered damaged kernels or considered against sound barley.

2/ Using a 5.5/64 x 3/4 slotted-hole sieve.

Barley

Definitions

Test weight (lb/bu) is pounds of grain per Winchester bushel as determined using an approved device on a dockage-free barley sample.

Test weight (kg/hl) is the metric system equivalent to pounds per bushel. Kilograms per hectoliter are calculated by multiplying pounds per bushel by 1.287.

Heat-damaged kernels are kernels and pieces of barley kernels, other grains, and wild oats that are materially discolored and damaged by heat.

Damaged kernels are kernels, pieces of barley kernels, other grains, and wild oats that are badly ground-damaged, badly weather-damaged, diseased, frost-damaged, germ-damaged, heat-damaged, injured-by-heat, insect-bored, mold-damaged, sprout-damaged, or otherwise materially damaged.

Foreign material is all matter other than barley, other grains, and wild oats that remains in the sample after removal of dockage.

Skinned and broken kernels are barley kernels that have one-third or more of the hull removed, or that the hull is loose or missing over the germ, or broken kernels, or whole kernels that have a part or all of the germ missing.

Dockage is all matter other than barley that can be removed from the original sample by use of an approved device. Also, underdeveloped, shriveled, and small pieces of barley kernels removed by properly separating the material other than barley and that cannot be recovered by properly rescreening or re-cleaning.

Moisture is the water content of grain as determined by an approved moisture meter. The percentage of moisture in a sample does not affect the numerical grade.

Suitable malting type are varieties of malting barley that are recommended by the American Malting Barley Association and any other proprietary malting type(s) used by the malting and brewing industries.

Sound barley is kernels and pieces of barley kernels that are not damaged.

Thin barley is: Six-rowed Malting barley that passes through a 5/64 x 3/4 slotted-hole sieve and Two-rowed Malting barley that passes through a 5.5/64 x 3/4 slotted-hole sieve; Six-rowed barley, Two-rowed barley, or Barley that passes through a 5/64 x 3/4 slotted-hole sieve.

Table 16. U.S. Barley Exports: Number of lots and quantity exported by grade, 2007-2009

Grade	2008		2009		2010	
	No. of Lots	Metric Tons	No. of Lots	Metric Tons	No. of Lots	Metric Tons
U.S. No. 2	23	412,099	3	38,606	5	69,771
All lots	23	412,099	3	38,606	5	69,771

Table 17. Summary of export Barley quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 2	45.0	23	51.2	49.3	53.5	3	52.2	51.4	52.6	5	51.0	49.0	54.4
	All lots	N/A	23	51.2	49.3	53.5	3	52.2	51.4	52.6	5	51.0	49.0	54.4
Test Weight (kg/hl)	U.S. No. 2	N/A	23	65.9	63.5	68.9	3	67.2	66.2	67.7	5	65.6	63.1	70.1
	All lots	N/A	23	65.9	63.5	68.9	3	67.2	66.2	67.7	5	65.6	63.1	70.1
Moisture	U.S. No. 2	N/A	23	10.9	9.9	12.7	3	11.2	11.0	11.7	5	12.7	10.8	13.4
	All lots	N/A	23	10.9	9.9	12.7	3	11.2	11.0	11.7	5	12.7	10.8	13.4
Heat-damaged Kernels	U.S. No. 2	0.3	23	0.0	0.0	0.1	3	0.1	0.0	0.1	5	0.2	0.0	0.3
	All lots	N/A	23	0.0	0.0	0.1	3	0.1	0.0	0.1	5	0.2	0.0	0.3
Damaged Kernels (Total)	U.S. No. 2	4.0	23	0.4	0.0	1.4	3	0.4	0.1	0.5	5	1.3	0.0	2.2
	All lots	N/A	23	0.4	0.0	1.4	3	0.4	0.1	0.5	5	1.3	0.0	2.2
Foreign Material	U.S. No. 2	2.0	23	0.1	0.0	0.3	3	0.2	0.0	0.5	5	0.1	0.1	0.2
	All lots	N/A	23	0.1	0.0	0.3	3	0.2	0.0	0.5	5	0.1	0.1	0.2
Sound Barley	U.S. No. 2	94.0	23	26.2	0.0	99.3	3	0.0	0.0	0.0	5	28.2	0.0	98.4
	All lots	N/A	23	26.2	0.0	99.3	3	0.0	0.0	0.0	5	28.2	0.0	98.4
Thin Barley	U.S. No. 2	15.0	23	7.7	2.6	10.3	3	5.6	5.0	7.2	5	3.8	1.1	7.5
	All lots	N/A	23	7.7	2.6	10.3	3	5.6	5.0	7.2	5	3.8	1.1	7.5
Broken Kernels	U.S. No. 2	8.0	23	0.6	0.0	1.0	3	0.9	0.4	1.2	5	0.3	0.0	0.7
	All lots	N/A	23	0.6	0.0	1.0	3	0.9	0.4	1.2	5	0.3	0.0	0.7
Dockage	U.S. No. 2	N/A	23	0.9	0.5	1.3	3	1.0	0.5	1.2	4	1.0	0.4	1.5
	All lots	N/A	23	0.9	0.5	1.3	3	1.0	0.5	1.2	4	1.0	0.4	1.5

N/A = Does not apply.

Canola

Canola Grades and Grade Requirements

There are no classes of canola. Canola is divided into three numerical grades and U.S. Sample grade. Special grades are provided to emphasize special qualities or conditions affecting the value, and are added to and made a part of the grade designation. They do not affect the numerical or sample grade designation.

U.S. Standards for Canola

Grade	Maximum percent limits of:								Maximum count limits of:		
	Damaged Kernels			Conspicuous Admixture				Inconspicuous admixture	Other Material		
	Heat Damage (percent)	Distinctly green (percent)	Total (percent)	Ergot (percent)	Sclerotinia (percent)	Stones (percent)	Total (percent)		Animal filth	Glass	Unknown foreign substances
U.S. No. 1	0.1	2.0	3.0	0.05	0.05	0.05	1.0	5.0	3	0	1
U.S. No. 2	0.5	6.0	10.0	0.05	0.10	0.05	1.5	5.0	3	0	1
U.S. No. 3	2.0	20.0	20.0	0.05	0.15	0.05	2.0	5.0	3	0	1

U.S. Sample Grade:

U.S. Sample Grade is canola that:

- (a) Does not meet the requirements for U.S. Nos. 1, 2, or 3; or
- (b) Has a musty, sour, or commercially objectionable foreign odor; or
- (c) Is heating or otherwise of distinctly low quality.

Canola

Definitions

Conspicuous admixture is all matter other than canola including, but not limited to, ergot, sclerotinia, and stones, which is conspicuous and readily distinguishable from canola and which remains in the sample after the removal of machine separated dockage. Conspicuous admixture is added to machine separated dockage in the computation of total dockage.

Damaged kernels are canola and pieces of canola that are heat-damaged, sprout-damaged, mold-damaged, distinctly green-damaged, frost-damaged, rime-damaged, or otherwise materially damaged.

Distinctly green kernels are canola and pieces of canola which, after being crushed, exhibit a distinctly green color.

Dockage is all matter other than canola that can be removed from the original sample by use of an approved device according to procedures described in FGIS instructions, as well as underdeveloped, shriveled, and small pieces of canola kernels that cannot be recovered by properly screening or re-cleaning. Machine separated dockage is added to conspicuous admixture in the computation of total dockage.

Ergot is sclerotia of the fungus, *Claviceps* species, which are associated with some seeds other than canola where the fungal organism has replaced the seed.

Heat-damaged kernels are canola and pieces of canola which, after being crushed, exhibit that they are discolored and damaged by heat.

Inconspicuous admixture is any seed which is difficult to distinguish from canola. This includes, but is not limited to, common wild mustard (*Brassica kaber* and *B. juncea*), domestic brown mustard (*Brassica juncea*), yellow mustard (*B. hirta*), and seed other than the mustard group.

Sclerotia are dark colored or black resting bodies of the *Sclerotinia* and *Claviceps*.

Sclerotinia is the genus name which includes the fungus *Sclerotinia sclerotiorum* which produces sclerotia. Canola is only infrequently infected, and the sclerotia, unlike sclerotia of ergot, are usually associated with the stem of the plants.

Table 18. U.S. Canola Exports: Number of lots and quantity exported by grade, 2007-2009

Grade	2008		2009		2010	
	No. of Lots	Metric Tons	No. of Lots	Metric Tons	No. of Lots	Metric Tons
U.S. No. 1	1	21,078	3	61,785	4	65,878
All lots	1	21,078	3	61,785	4	65,878

-- = No lots reported in this category.

Table 19. Summary of export Canola quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of				No. of				No. of			
			Lots	Avg.	Low	High	Lots	Avg.	Low	High	Lots	Avg.	Low	High
Dockage	U.S. No. 1	N/A	1	1.9	1.9	1.9	3	2.3	2.3	2.4	4	2.6	2.3	2.7
	All lots	N/A	1	1.9	1.9	1.9	3	2.3	2.3	2.4	4	2.6	2.3	2.7
Moisture	U.S. No. 1	N/A	1	8.1	8.1	8.1	3	7.7	7.5	7.8	4	7.7	7.3	8.2
	All lots	N/A	1	8.1	8.1	8.1	3	7.7	7.5	7.8	4	7.7	7.3	8.2
Heat-damaged Kernels	U.S. No. 1	0.1	1	0.0	0.0	0.0	3	0.0	0.0	0.0	4	0.0	0.0	0.0
	All lots	N/A	1	0.0	0.0	0.0	3	0.0	0.0	0.0	4	0.0	0.0	0.0
Distinctly green Kernels	U.S. No. 1	2.0	1	0.3	0.3	0.3	3	0.5	0.4	0.6	4	0.5	0.3	0.7
	All lots	N/A	1	0.3	0.3	0.3	3	0.5	0.4	0.6	4	0.5	0.3	0.7
Total Damaged Kernels	U.S. No. 1	3.0	1	0.4	0.4	0.4	3	0.7	0.6	0.7	4	0.6	0.4	0.8
	All lots	N/A	1	0.4	0.4	0.4	3	0.7	0.6	0.7	4	0.6	0.4	0.8
Ergot	U.S. No. 1	0.05	1	0.0	0.0	0.0	3	0.0	0.0	0.0	4	0.0	0.0	0.0
	All lots	N/A	1	0.0	0.0	0.0	3	0.0	0.0	0.0	4	0.0	0.0	0.0
Sclerotinia	U.S. No. 1	0.05	1	0.0	0.0	0.0	3	0.0	0.0	0.0	4	0.0	0.0	0.0
	All lots	N/A	1	0.0	0.0	0.0	3	0.0	0.0	0.0	4	0.0	0.0	0.0
Stones	U.S. No. 1	0.05	1	0.0	0.0	0.0	3	0.0	0.0	0.0	4	0.0	0.0	0.0
	All lots	N/A	1	0.0	0.0	0.0	3	0.0	0.0	0.0	4	0.0	0.0	0.0
Total Conspicuous Admixture	U.S. No. 1	1.0	1	0.1	0.1	0.1	3	0.2	0.1	0.2	4	0.2	0.1	0.3
	All lots	N/A	1	0.1	0.1	0.1	3	0.2	0.1	0.2	4	0.2	0.1	0.3
Inconspicuous Admixture	U.S. No. 1	5.0	1	0.0	0.0	0.0	3	0.1	0.0	0.1	4	0.0	0.0	0.1
	All lots	N/A	1	0.0	0.0	0.0	3	0.1	0.0	0.1	4	0.0	0.0	0.1

N/A = Does not apply.

-- = No lots reported in this category.

Flaxseed

Flaxseed Grades and Grade Requirements

There are no classes of flaxseed. Flaxseed is divided into two numerical grades and U.S. Sample grade. Other determinations not specifically provided for under the general provisions are made on the basis of the grain when free from dockage, except the determination of odor is made on either the basis of the grain as a whole or the grain when free from dockage.

U.S. Standards for Flaxseed

Grade:	Minimum Test Weight per bushel (pounds)	Max Limits of damaged kernels:	
		Heat-Damaged kernels (percent)	Damaged kernels total (percent)
U.S. No. 1	49.0	0.2	10.0
U.S. No. 2	47.0	0.5	15.0

U.S. Sample Grade:

U.S. Sample Grade is flaxseed that:

- (a) Does not meet the requirements for the grades U.S. Nos. 1 or 2; or
- (b) Contains 8 or more stones which have an aggregate weight in excess of 0.2 percent of the sample weight, 2 or more pieces of glass, 3 or more crotalaria seeds (*Crotalaria* spp.), 2 or more castor beans (*Ricinus communis* L.), 4 or more particles of an unknown foreign substance(s) or a commonly recognized harmful or toxic substance(s), 10 or more rodent pellets, bird dropping, or equivalent quantity of other animal filth per 1-1/8 to 1-1/4 quarts of flaxseed; or
- (c) Has a musty, sour, or commercially objectionable foreign odor (except smut or garlic odor); or
- (d) Is heating or otherwise of distinctly low quality.

Definitions

Flaxseed. Grain that, before the removal of dockage, consists of 50 percent or more of common flaxseed (*Linum usitatissimum* L.) and not more than 20 percent of other grains for which standards have been established under the United States Grain Standards Act and which, after the removal of dockage, contains 50 percent or more of whole flaxseed.

Damaged kernels. Kernels and pieces of flaxseed kernels that are badly ground damaged, badly weather-damaged, diseased, frost-damaged, germ-damaged, heat damaged, insect-bored, mold-damaged, sprout-damaged, or otherwise materially damaged.

Dockage. All matter other than flaxseed that can be removed from the original sample by use of an approved device according to procedures prescribed in FGIS instructions. Also, underdeveloped, shriveled, and small pieces of flaxseed kernels removed in properly separating the material other than flaxseed and that cannot be recovered by properly rescreening or re-cleaning.

Heat-damaged kernels. Kernels and pieces of flaxseed kernels that are materially discolored and damaged by heat.

Other grains. Barley, corn, cultivated buckwheat, einkorn, emmer, guar, hull-less barley, non-grain sorghum, oats, Polish wheat, popcorn, poulard wheat, rice, rye, safflower, sorghum, soybeans, spelt, sunflower seed, sweet corn, triticale, wheat, and wild oats.

Table 20. U.S. Flaxseed Exports: Number of lots and quantity exported by grade, 2008-2010

Grade	2008		2009		2010	
	No. of Lots	Metric Tons	No. of Lots	Metric Tons	No. of Lots	Metric Tons
U.S. No. 1	3	21,112	2	14,754	3	46,550
All lots	3	21,112	2	14,754	3	46,550

Table 21. Summary of export Flaxseed quality, 2008-2010

Factor	Grade	Grade Limit	2008				2009				2010			
			No. of Lots	Avg.	Low	High	No. of Lots	Avg.	Low	High	No. of Lots	Avg.	Low	High
Test Weight (lb/bu)	U.S. No. 1	49.0	3	50.1	49.7	50.7	2	50.0	50.0	50.1	3	49.8	49.6	50.2
	All lots	N/A	3	50.1	49.7	50.7	2	50.0	50.0	50.1	3	49.8	49.6	50.2
Test Weight (kg/hl)	U.S. No. 1	N/A	3	64.5	64.0	65.3	2	64.4	64.3	64.5	3	64.2	63.9	64.6
	All lots	N/A	3	64.5	64.0	65.3	2	64.4	64.3	64.5	3	64.2	63.9	64.6
Moisture	U.S. No. 1	N/A	3	7.1	6.7	7.3	2	7.9	7.7	8.0	3	7.5	7.4	7.8
	All lots	N/A	3	7.1	6.7	7.3	2	7.9	7.7	8.0	3	7.5	7.4	7.8
Heat-damaged kernels	U.S. No. 1	0.2	3	0.0	0.0	0.0	2	0.0	0.0	0.0	3	0.0	0.0	0.0
	All lots	N/A	3	0.0	0.0	0.0	2	0.0	0.0	0.0	3	0.0	0.0	0.0
Damaged Flaxseed (Total)	U.S. No. 1	10.0	3	0.0	0.0	0.0	2	0.1	0.0	0.1	3	0.0	0.0	0.1
	All lots	N/A	3	0.0	0.0	0.0	2	0.1	0.0	0.1	3	0.0	0.0	0.1
Dockage	U.S. No. 1	N/A	3	5.2	4.7	5.5	2	4.6	4.0	5.4	3	4.6	4.3	5.1
	All lots	N/A	3	5.2	4.7	5.5	2	4.6	4.0	5.4	3	4.6	4.3	5.1


N/A = Does not apply.

Appendix


U.S. Grain Quantity Exported, 2001-2010.....	62
U.S. Grain Exported, 2001-2010, Average Dockage by Grain.....	63
U.S. Wheat Exported, 2001-2010, Average Dockage by Class.....	64
U.S. Wheat Exported, 2001-2010, Average Dockage by Destination Country.....	65
U.S. Wheat Exported, 2001-2010, Average Protein (12% Moisture) by Class.....	66
U.S. Soybeans Exported, 2001-2010, Average Protein and Oil (13% Moisture).....	67
U.S. Grains Exported, 2001-2010, Average FM, BNFM, and BCFM.....	68

U.S. GRAIN QUANTITY EXPORTED, 2001-2010

METRIC TONS IN MILLIONS




U.S. GRAIN EXPORTED, 2001–2010
AVG DOCKAGE BY GRAIN




U.S. WHEAT EXPORTED, 2001-2010


AVG DOCKAGE BY CLASS




U.S. WHEAT EXPORTED, 2001-2010
AVG DOCKAGE BY DESTINATION COUNTRY



U.S. WHEAT EXPORTED, 2001-2010
AVG PROTEIN(12% M) BY CLASS




U.S. SOYBEANS EXPORTED, 2001–2010
AVG PROTEIN AND OIL (13% M)



U.S. GRAINS EXPORTED, 2001-2010

AVG FM BNFM BCFM



The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.