Leading Causes of Death by Age Group, White Females-United States, 2006*

| | | Age Group in Years | | | | | | | | | | | | | |
|------|------------------------------|--|------------------------------|------------------------------|------------------------------------|------------------------------------|--|--|--|--|--|--|---|--|--|
| Rank | 1-4 | 5-9 | 10-14 | 15-19 | 20-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65 + | 65-74 | 75-84 | 85 + | All Ages | |
| 1 | Unintentional injuries 34.1% | Unintentional injuries 35.2% | Unintentional injuries 36.3% | Unintentional injuries 56.9% | Unintentional injuries 45.9% | Unintentional injuries 32.2% | Cancer 27.0% | Cancer 37.5% | Cancer 42.6% | Heart disease 28.7% | Cancer 37.1% | Heart disease 25.6% | Heart disease 34.0% | Heart disease 26.0% | |
| 2 | Birth defects 12.7% | Cancer 18.8% | Cancer 16.4% | Suicide 8.1% | Suicide 8.0% | Cancer 15.5% | Unintentional injuries 19.1% | Heart disease 14.0% | Heart disease 17.0% | Cancer 19.2% | Heart disease 19.8% | Cancer 23.3% | Cancer 10.2% | Cancer 22.0% | |
| 3 | Cancer 8.5% | Birth defects 7.4% | Suicide 5.0% | Cancer 6.8% | Cancer 7.7% | Suicide 9.0% | Heart disease 11.0% | Unintentional injuries 9.3% | Chronic lower respiratory diseases 6.0% | Stroke 7.5% | Chronic lower respiratory diseases 9.1% | Chronic lower respiratory diseases 7.6% | Stroke 8.5% | Stroke 6.7% | |
| 4 | Homicide 5.6% | Homicide 6.2% | Birth defects 4.9% | Homicide 5.2% | Homicide 6.3% | Heart disease 6.7% | Suicide 6.3% | Stroke 3.5% | Diabetes 3.8% | Chronic lower respiratory diseases 6.2% | Stroke 4.8% | Stroke 7.3% | Alzheimer's disease 7.6% | Chronic lower respiratory diseases 5.8% | |
| 5 | Heart disease 3.4% | Heart disease 3.2% | Heart disease (5) 4.5% | Heart disease 2.5% | Heart disease 4.1% | Homicide 4.2% | Chronic liver disease 3.2% | Chronic liver disease 3.5% | Stroke 3.7% | Alzheimer's disease 5.5% | Diabetes 3.7% | Alzheimer's disease 4.5% | Chronic lower respiratory diseases 4.2% | Alzheimer's disease 4.5% | |
| 6 | Influenza & pneumonia 2.8% | Influenza & pneumonia 1.8% | Homicide (5) 4.5% | Birth defects 2.2% | Pregnancy complications 2.5% | Diabetes 2.0% | Stroke 2.9% | Suicide 3.3% | Unintentional injuries 3.4% | Influenza & pneumonia 2.9% | Unintentional injuries 1.9% | Diabetes 3.0% | Influenza & pneumonia 3.6% | Unintentional injuries 3.5% | |
| 7 | Benign neoplasms 2.2% | Septicemia (7) 1.4% | Benign neoplasms 1.4% | Influenza & pneumonia 1.0% | Birth defects 1.9% | Pregnancy complications 2.0% | Diabetes 2.5% | Chronic lower respiratory diseases 3.1% | Chronic liver disease 2.0% | Diabetes 2.6% | Kidney disease 1.7% | Influenza & pneumonia 2.5% | Unintentional injuries 2.0% | Diabetes 2.7% | |
| 8 | Septicemia 2.1% | Benign neoplasms (7) 1.4% | Septicemia 1.3% | Stroke 0.9% | Stroke 1.1% | Stroke 1.7% | Homicide 2.0% | Diabetes 2.8% | Septicemia 1.5% | Unintentional injuries 2.0% | Septicemia 1.6% | Unintentional injuries 2.0% | Diabetes 1.9% | Influenza & pneumonia 2.6% | |
| 9 | Stroke 1.6% | Stroke 1.3% | Stroke 1.0% | Pregnancy complications 0.8% | Diabetes 0.9% | Birth defects 1.5% | HIV disease 1.5% | Septicemia 1.2% | Kidney disease 1.4% | Kidney disease 1.9% | Influenza & pneumonia 1.6% | Kidney disease 2.0% | Kidney disease 1.8% | Kidney disease 1.7% | |
| 10 | Perinatal conditions 1.0% | Chronic lower respiratory diseases 0.7% | Influenza & pneumonia 0.7% | Septicemia 0.7% | Influenza & pneumonia 0.9% | Influenza & pneumonia 1.2% | Chronic lower respiratory diseases 1.4% | Viral Hepatitis 1.2% | Suicide 1.2% | Septicemia 1.4% | Alzheimer's disease 1.3% | Septicemia 1.6% | Hypertension 1.5% | Septicemia 1.4% | |

^{*}Percentages represent total deaths in the age group due to the cause indicated. Numbers in parentheses indicate tied rankings. The white, black, American Indian/Alaska Native, and Asian/Pacific Islander race groups include persons of Hispanic and non-Hispanic origin may be of any race. Some terms have shortened from those used in the National Vital Statistics Report. See the next page for a listing of the shortened terms in the table and their full unabridged equivalents used in the report. To learn more, visit *Mortality Tables* at http://www.cdc.gov/nchs/nvss/mortality_tables.htm (HHS, CDC, NCHS).

Short and Full Terms for Leading Causes of Death

Some terms in the leading causes of death table have been shortened from those used in the National Vital Statistics Report. Below is a listing of the shortened terms used in the table and their full, unabridged equivalents used in the report.

SHORT TERMS FULL TERMS

Aortic aneurism Aortic aneurism and dissection

Benign neoplasms In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior

Birth defects Congenital malformations, deformations and chromosomal abnormalities

Bronchitis Acute bronchitis and bronchiolitis

Cancer Malignant neoplasms

Chronic liver disease Chronic liver disease and cirrhosis

Diabetes Diabetes mellitus

Gallbladder disorders Cholethiasis and other disorders of gallbladder

Heart disease Diseases of heart

HIV disease Human immunodeficiency virus (HIV) disease

Homicide Assault (homicide)

Hypertension Essential (primary) hypertension and hypertensive renal disease

Kidney disease Nephritis, nephrotic syndrome and nephrosis

Kidney infection Infections of kidney

Medical & surgical care complications Complications of medical and surgical care

Perinatal conditions Certain conditions originating in the perinatal period

PreumonitisPregnancy complications

Pregnancy, childbirth and the puerperium

Stroke Cerebrovascular diseases
Suicide Intentional self-harm

Unintentional injuries Accidents (unintentional injuries)