

# **Management of Food Allergies**

**Federal Bureau of Prisons**

**Clinical Practice Guidelines**

**September 2012**

Clinical guidelines are made available to the public for informational purposes only. The Federal Bureau of Prisons (BOP) does not warrant these guidelines for any other purpose, and assumes no responsibility for any injury or damage resulting from the reliance thereof. Proper medical practice necessitates that all cases are evaluated on an individual basis and that treatment decisions are patient specific. Consult the BOP Clinical Practice Guideline Web page to determine the date of the most recent update to this document: <http://www.bop.gov/news/medresources.jsp>.

## Table of Contents

<b>1. Purpose .....</b>	<b>1</b>
<b>2. Food Allergy Overview .....</b>	<b>1</b>
<b>3. Food Allergy Assessment .....</b>	<b>3</b>
<b>4. Evaluation and Management of Potential Anaphylactic Food Allergies.....</b>	<b>4</b>
<b>5. Evaluation and Management of Potential Non-Anaphylactic Food Allergies .....</b>	<b>4</b>
<b>6. Diet Orders .....</b>	<b>4</b>
<b>Medical Diet Orders/Self-Selection.....</b>	<b>4</b>
<b>Special Diet Orders .....</b>	<b>5</b>
<b>7. Dietitian Referral.....</b>	<b>5</b>
<b>8. Nutrition Education .....</b>	<b>5</b>
<b>9. Satellite Feeding .....</b>	<b>6</b>
<b>10. Work and Housing Detail.....</b>	<b>6</b>
<b>General Definitions .....</b>	<b>7</b>
<b>References.....</b>	<b>8</b>
<b>Appendix 1: Definitions of Specific Food-Induced Allergic Conditions .....</b>	<b>9</b>
<b>Appendix 2: BOP Food Allergy Questionnaire.....</b>	<b>11</b>
<b>Appendix 3: Diagnostic Criteria for Anaphylaxis.....</b>	<b>13</b>
<b>Appendix 4: Algorithm for Patients with Suspected Food-Induced Anaphylaxis.....</b>	<b>15</b>
<b>Appendix 5: Pharmacological Treatment of Anaphylaxis.....</b>	<b>16</b>
<b>Appendix 6: Algorithm for Patients Without History of Suspected Food-Induced Anaphylaxis.....</b>	<b>17</b>
<b>Appendix 7: Inmate Handouts.....</b>	<b>18</b>
<b>Inmate Factsheet: An Overview of Food Allergies.....</b>	<b>19</b>
<b>Inmate Factsheet: Lactose Intolerance.....</b>	<b>20</b>
<b>Inmate Factsheet: Food Avoidance and Self-Selection from the BOP National Menu .....</b>	<b>21</b>

## 1. Purpose

The Federal Bureau of Prisons (BOP) Clinical Practice Guidelines for the *Management of Food Allergies* provide recommendations for the diagnosis and management of federal inmates with suspected food allergies.

## 2. Food Allergy Overview

### *Food Allergy vs. Food Intolerance*

Food allergy has no basic universally accepted definition. The National Institutes of Health (NIH) defines food allergy as “an adverse immune response that occurs *reproducibly* on exposure to a given food and is distinct from other adverse responses to food, such as food intolerance, pharmacologic reactions, and toxin-mediated reactions.” However, in published articles on food allergy, definitions frequently vary, thereby confounding the recommendations on diagnosing and managing patients with food allergies. Nevertheless, the distinction between a *food allergy* with an allergic response and *food intolerance*, such as the inability to digest the sugar lactose, is clinically relevant. (See the [General Definitions](#) section in these guidelines.)

### *Prevalence of Food Allergies*

The prevalence of food allergies is poorly defined, and estimates range from 0.2% to 3.5% in the general population. Estimates of peanut allergy prevalence range from 0.3% to 0.9%.

Although childhood food allergies tend to wane with aging, a subset of these patients will have food allergies that persist into adulthood. Furthermore, some adults develop allergies *de novo* from sensitization to food allergens encountered after childhood.

### *IgE-Mediated and Non-IgE-Mediated Food Allergic Reactions*

The distinction between IgE-mediated reactions and non-IgE-mediated reactions to food allergens is clinically important. *IgE-mediated food allergic reactions* are rapid in onset, typically beginning within minutes to two hours from the time of ingestion. Presentations include circulatory collapse, dyspnea, wheezing, stridor, angioedema, oropharyngeal symptoms, and urticarial rash. The most common foods associated with anaphylaxis are peanuts, tree nuts, and crustacean shellfish; however, milk and eggs can also induce IgE-mediated allergic responses. *Non-IgE-mediated reactions* are much more subacute or chronic and are usually isolated to the gastrointestinal tract and/or skin.

### *Diagnosis and Treatment*

There are no well-accepted criteria for diagnosing food allergies. However, the following diagnostics tests are NOT recommended for evaluating food allergies: intradermal allergen testing, total serum IgE quantification, and atopy patch testing.

*Skin prick tests* and *serum food-specific IgE assays* are potentially valuable diagnostic tests for food allergies; however, neither one is superior to the other, and both are considered nonconfirmatory of a specific food allergy—thus limiting their diagnostic efficacy.

- Skin prick testing for a given food allergy is not very specific diagnostically, as patients with a positive test still have a 40% chance of being able to eat the food in question without difficulty.
- Food-specific IgE assays (commonly known as RAST tests) are sensitive tests diagnostically, but also are not very specific. If negative, the specific food allergy is unlikely. If positive, the patient still may not have a true food allergy. They are most useful for confirming the diagnosis of a suspected specific food allergy. In the BOP, the primary role for RAST testing is in confirming allergy to milk, wheat, or baked egg in inmates with a history of anaphylaxis. IgE assays to a panel of potential antigens are usually not helpful or indicated—except possibly when ruling out claims of multiple food allergens—and should only be ordered in consultation with the Central Office Registered Dietitian (see [Appendix 4, Algorithm for Patients with Suspected Food-Induced Anaphylaxis](#)).

The gold standard for diagnosing a food allergy is a *placebo-controlled oral food challenge*. However, this testing requires specialized personnel, time, expense, and the risk of anaphylaxis, limiting the use of this diagnostic test in the community, let alone within the correctional setting.

*Therefore, in evaluating inmates for food allergies, the focus should be on identifying inmates at risk for anaphylaxis:* providing them epinephrine, if indicated, and pursuing diagnostic testing on a very limited basis—primarily for those inmates with questionable IgE-mediated food allergies. The vast majority of other inmates with non-IgE-mediated food allergies should be provided education on targeted food selection.

*Elimination diets* are the mainstay of therapy for patients with food allergies, although the effectiveness of this strategy is poorly studied. Immunotherapy for food allergies is unproven and not recommended.

*Within the BOP, a diagnosis of a food allergy should not be confirmed and documented as a patient's health problem unless:*

1. The food allergy was previously diagnosed by an outside medical provider and documented in the patient's medical records.
2. The patient was diagnosed while in the custody of the BOP, using standards indicated in these guidelines, including:
  - A thorough assessment has been conducted with the use of [Appendix 2](#), and the patient has been identified as having a history of reproducible food allergy-related symptoms upon exposure to an identified allergen.
  - The specific food allergen has been positively confirmed with RAST testing, as outlined in [Appendix 4](#) or [Appendix 6](#).

### 3. Food Allergy Assessment

#### *Medical History*

The medical history should focus on:

- Any past history of food allergy evaluations.
  - Anaphylactic episodes (including emergency room visits hospitalizations and prescriptions for hand-carried epinephrine).
  - History of poor outcomes from anaphylaxis therapy related to the use of beta-blocker or ACE inhibitor therapy.
  - History of asthma (particularly poorly controlled) or coronary artery disease.
  - The timing and descriptions of symptoms relative to ingestion of specific foods, e.g., wheezing, voice change related to laryngeal edema, urticarial, rashes.
  - The association of allergic symptoms with exercise or other complementary factors such as the use of aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs), or alcohol.
  - A history of asthma, dysphagia, or eosinophilic esophagitis.
  - A personal or family history of atopic dermatitis.
- Clinicians should familiarize themselves with [Appendix 1](#), which outlines the wide range of specific food-induced allergic conditions that may be diagnostically relevant. Key questions for evaluating food allergies are outlined in [Appendix 2](#), *BOP Food Allergy Questionnaire*.

#### *Physical Exam*

The physical exam should include:

- Vital signs
- Pulse oximetry
- Weight measurement
- Auscultation of the lungs, HEENT, CV
- Thorough examination of the skin for signs of atopic dermatitis

Other examinations should be conducted to evaluate for other co-morbidities that are indicated by the patient's medical history.

#### *Assessment*

An assessment should be made as to whether or not anaphylactic food allergy is a concern, based on the inmate's medical history of IgE-mediated allergic episodes and the specific offending food allergens.

- See the diagnostic criteria and related information outlined in [Appendix 3](#), *Diagnostic Criteria for Anaphylaxis*.

## 4. Evaluation and Management of Potential Anaphylactic Food Allergies

Inmates with suspected anaphylactic food allergies should be evaluated and managed in accordance with the stepwise approach outlined in [Appendix 4, Algorithm for Patients with Suspected Food-Induced Anaphylaxis](#). Inmates who have bona fide medical history of anaphylactic food allergies (e.g., history of hospitalization or EpiPen™ prescriptions) should be:

- Prescribed an epinephrine auto-injector (to be self-carried at all times).
- Given a copy of the [Inmate Factsheet: An Overview of Food Allergies](#), which includes information on potential anaphylactic symptoms and the use of self-administered epinephrine (see [Appendix 7](#)).
- Provided appropriate information on food selections, such as that provided in the [Inmate Fact Sheet: Food Avoidance and Self-Selection from the BOP National Menu](#) (see [Appendix 7](#)),
- Given information on reading food labels, as provided by the Food Allergy and Anaphylaxis Network at <http://www.foodallergy.org/page/patient-handouts>. Handouts are available for identifying foods containing milk, egg, peanuts, tree nuts, wheat, soy, and shellfish.

Guidance for health care providers on the pharmacologic prevention and treatment of anaphylaxis is outlined in [Appendix 5](#).

## 5. Evaluation and Management of Potential Non-Anaphylactic Food Allergies

Inmates with suspected non-IgE-mediated food allergies should be evaluated and managed in accordance with the stepwise approach shown in [Appendix 6, Algorithm for Patients Without History of Suspected Food-Induced Anaphylaxis](#). The algorithm outlines how patients with suspected food allergies must be managed differently for different allergens: baked egg or wheat, milk, another individual food, or multiple foods.

Clinicians should also be aware of the potential association of certain diseases and syndromes with non-IgE-mediated food allergies, as outlined in [Appendix 1, Definitions of Specific Food-Induced Allergic Conditions](#). Inmates diagnosed with lactose intolerance should be provided the patient information contained in the [Inmate Fact Sheet: Lactose Intolerance](#) (see [Appendix 7](#)).

## 6. Diet Orders

### Medical Diet Orders/Self-Selection

Diet orders for *food allergies* are to be offered only when medically necessary—and not for *food intolerance* or preference. In all cases when a diet order is being considered, the first option should be the inmate's simple avoidance of the item, with guidance provided by the [Inmate Fact Sheet: Food Avoidance and Self-Selection from the BOP National Menu](#) (see [Appendix 7](#)). For all individual food allergies except apple, citrus fruit, banana, baked egg, wheat, or milk, the

inmate may simply avoid the item or self-select the no-flesh option or heart-healthy alternative option. In accordance with the BOP *Guidelines for Medical Diets*, medical diets that will be provided through self-selection may be ordered by any mid-level practitioner (MLP), clinical director (CD), staff physician, staff psychiatrist, or staff dentist.

### **Special Diet Orders**

*A Special Diet should not be considered for allergic avoidance, unless:*

- The food allergy is reported for apple, citrus fruit, or banana.
- The individual has a confirmed diagnosis of baked egg, wheat, or milk, or a confirmed diagnosis of multiple-food allergies.

*In accordance with the Program Statement PS6031.01, Patient Care:*

- Special Diets will be prescribed *only* by the CD or by a staff physician, staff psychiatrist, or staff dentist.
- MLPs at Medical Referral Centers (MRC) may prescribe a Special Diet, but it must be countersigned by the primary physician.

*All Medical and Special Diets related to food allergies must be:*

- Documented in the patient's medical record.
- Furnished in writing via e-mail to the Food Service Administrator (FSA).
- Rewritten annually or more often if indicated.

## **7. Dietitian Referral**

For individuals at Medical Referral Centers (MRCs), if a Special Diet for a food allergy is ordered, a consultation should be placed to an MRC staff registered dietitian. For all other institutions, if a Special Diet for a food allergy is ordered a consultation should be placed to a Central Office registered dietitian per BOP Tele-Nutrition Standard Operating Procedures ([http://sallyport.bop.gov/co/hsd/food\\_svc/docs/Tele-Nutrition%20Info.jsp](http://sallyport.bop.gov/co/hsd/food_svc/docs/Tele-Nutrition%20Info.jsp)).

The registered dietitian will work with the institution Food Service Administrator to make sure appropriate precautions are taken in elimination of causative foods, preparation of meals, and if needed appropriate dietary substitutions to the National Menu to maintain nutritional adequacy.

## **8. Nutrition Education**

If indicated, nutrition education will be provided to assist patients in their ability to:

- Define the allergen-free nutrition prescription.
- List common foods that contain the allergen.
- List foods that are allowed in the allergen-free prescribed diet.
- List nutrient dense alternatives to foods that do not contain allergens.
- Read a food label and be able to identify hidden sources of allergens.

- Understand the risk involved in packaged foods with precautionary labeling.
- Identify problem solving avoidance skills.
- Describe symptoms of a food-induced allergic reaction and what should be done during a reaction.

Standardized patient handouts on reading food labels and understanding hidden sources of major food allergens are available at <http://www.foodallergy.org/page/patient-handouts>.

If local medical staff would like assistance in providing patients with nutrition education or in counseling patients with diagnosed food allergies, they can contact a Central Office Registered Dietitian for assistance by following the BOP Tele-Nutrition Standard Operating Procedures ([http://sallyport.bop.gov/co/hsd/food\\_svc/docs/Tele-Nutrition%20Info.jsp](http://sallyport.bop.gov/co/hsd/food_svc/docs/Tele-Nutrition%20Info.jsp)).

Guidelines for follow-up nutrition monitoring and evaluation of adherence to diet and avoidance will be provided as needed per the discretion of the referring provider.

## **9. Satellite Feeding**

If inmates in satellite service areas are not provided the opportunity to select a tray without identified or diagnosed allergens, the Food Service Administrator will develop procedures that ensure that inmates with Medical and Special Diet orders receive the proper diet.

## **10. Work and Housing Detail**

Consideration should be given on a case-by-case basis regarding appropriateness of alternative accommodations related to work assignments involving the handling of food. Consideration of housing detail should also be given on a case-by-case basis to reduce the potential for inadvertent exposure to the food allergen.



## General Definitions

**Food Allergens** are specific components of food or ingredients within food that are recognized by allergen-specific immune cells and elicit specific immunologic reactions, resulting in characteristic symptoms.

**Food Allergy** is an adverse health effect arising from a specific immune response that occurs reproducibly on exposure to a given food.

→ See also [Appendix 1](#), *Definitions of Specific Food-Induced Allergic Conditions*.

**Food Intolerance** occurs with foods or food components that elicit reproducible adverse reactions, but do not have established or likely immunologic mechanisms.

**Lactose Intolerance** is the onset of gastrointestinal symptoms (diarrhea, abdominal pain, flatulence, and/or bloating) following a blinded, single-dose challenge of ingested lactose by an individual with lactose malabsorption, which is not observed when the person ingests an indistinguishable placebo.

**Tolerance** is used in these guidelines to denote that an individual is symptom-free after consumption of that food or upon oral food challenge weeks, months, or even years after the cessation of treatment.

## References

Academy of Nutrition and Dietetics. *Nutrition Care Manual*. [homepage on the Internet]. 2012. Available from:

[http://nutritioncaremanual.org/topic.cfm?ncm\\_heading=Nutrition%20Care&ncm\\_toc\\_id=19659](http://nutritioncaremanual.org/topic.cfm?ncm_heading=Nutrition%20Care&ncm_toc_id=19659).

Accessed January 26, 2012.

Boyce JA, Assa'ad A, Burks AW, et al. Guidelines for the diagnosis and management of food allergy in the United States: report of the NIAID-Sponsored Expert Panel. *J Allergy Clin Immunol*. 2010;126(6)(suppl):S1–S58. Available from:

[http://www.jacionline.org/article/S0091-6749\(10\)01566-6/fulltext](http://www.jacionline.org/article/S0091-6749(10)01566-6/fulltext). Accessed January 26, 2012.

Chafen JJS, Newberry SJ, Riedl MA. Diagnosing and managing common food allergies: a systematic review. *JAMA*. 2010;303(18):1848–1856. Available from:

<http://jama.jamanetwork.com/article.aspx?articleid=185820>. Accessed April 26, 2012.

Keller, Jeff. *In-Depth Food Allergy Protocol*. 2012. Located at: Ada County Jail, Boise, ID.

Mayo Clinic Staff. Lactose Intolerance. [homepage on the Internet]. 2010. Available from:

<http://www.mayoclinic.com/health/lactose-intolerance/DS00530>. Accessed January 26, 2012.

Suchy FJ, Brannon PM, Carpenter TO, et al. NIH Consensus Development Conference Statement: lactose intolerance and health. *NIH Consensus State Sci Statements*. 2010;27(2):1–27.

Available from: [http://consensus.nih.gov/2010/docs/LI\\_CDC\\_2010\\_Final%20Statement.pdf](http://consensus.nih.gov/2010/docs/LI_CDC_2010_Final%20Statement.pdf).

Accessed January 26, 2012.

## Appendix 1: Definitions of Specific Food-Induced Allergic Conditions

**A number of specific clinical syndromes that may occur as a result of food allergy are defined below.**

*Source:* Guidelines for the Diagnosis and Management of Food Allergy in the United States: Report of the NIAID-Sponsored Expert Panel. (See [References](#) section for full citation.)

### FOOD-INDUCED ANAPHYLAXIS

Food-induced anaphylaxis is a serious allergic reaction that is rapid in onset and may cause death. Typically, IgE-mediated food-induced anaphylaxis is believed to involve systemic mediator release from sensitized mast cells and basophils. In some cases, such as food-dependent, exercise-induced anaphylaxis, the ability to induce reactions depends on the temporal association between food consumption and exercise, usually within two hours.

### GASTROINTESTINAL FOOD ALLERGIES

Gastrointestinal food allergies include a spectrum of disorders that result from adverse immunologic responses to dietary antigens. Although significant overlap may exist between these conditions, several specific syndromes have been defined as follows:

**Immediate GI hypersensitivity** refers to an IgE-mediated food allergy in which upper GI symptoms may occur within minutes, and lower GI symptoms may occur either immediately or with a delay of up to several hours. This is commonly seen as a manifestation of anaphylaxis. Among the GI conditions, acute, immediate vomiting is the most common reaction and the one that is best documented as immunologic and IgE mediated.

**Eosinophilic esophagitis (EoE)** involves localized eosinophilic inflammation of the esophagus. In some patients, avoidance of specific foods will result in normalization of histopathology. Although EoE is commonly associated with the presence of food-specific IgE, the precise causal role of the food allergy in its etiology is not well defined. Both IgE- and non-IgE-mediated mechanisms appear to be involved. In adults, EoE most often presents with dysphagia and esophageal food impactions.

**Eosinophilic gastroenteritis (EG)** also is both IgE- and non-IgE-mediated and is commonly linked to food allergy. EG describes a constellation of symptoms that vary depending on the portion of the GI tract involved and a pathologic infiltration of the GI tract by eosinophils, which may be localized or widespread. EoE is a common manifestation of EG.

**Oral allergy syndrome (OAS)**, also referred to as pollen-associated food allergy syndrome, is a form of localized IgE mediated allergy, usually to raw fruits or vegetables, with symptoms confined to the lips, mouth, and throat. OAS most commonly affects patients who are allergic to certain pollens. Symptoms include itching of the lips, tongue, roof of the mouth, and throat—with or without swelling, and/or tingling of the lips, tongue, roof of the mouth, and throat, or anaphylaxis. Isolated oral allergy syndrome (not systemic or GI symptoms) is the most common presentation and in >95% of patients is not associated with the later development of anaphylactic reactions. Patients should generally not be given epinephrine auto-injectors.

**A number of specific clinical syndromes that may occur as a result of food allergy are defined below.**

*Source:* Guidelines for the Diagnosis and Management of Food Allergy in the United States: Report of the NIAID-Sponsored Expert Panel. (See [References](#) section for full citation.)

### CUTANEOUS REACTIONS

Cutaneous reactions to foods are some of the most common presentations of food allergy and include IgE-mediated (urticaria, angioedema, flushing, pruritus), cell-mediated (contact dermatitis, dermatitis herpetiformis), and mixed IgE- and cell-mediated (atopic dermatitis) reactions, as follows:

**Acute urticaria** is a common manifestation of IgE-mediated food allergy, although food allergy is not the most common cause of acute urticaria and is rarely a cause of chronic urticaria. Lesions develop rapidly after ingesting the problem food and appear as polymorphic, round, or irregular-shaped pruritic wheals, ranging in size from a few millimeters to several centimeters.

**Angioedema** most often occurs in combination with urticaria and, if food-induced, is typically IgE-mediated. It is characterized by nonpitting, nonpruritic, well-defined edematous swelling that involves subcutaneous tissues (for example, face, hands, buttocks, and genitals), abdominal organs, or the upper airway.

→ *When the upper airway is involved, laryngeal angioedema is a medical emergency requiring prompt assessment. Both acute angioedema and urticaria are common features of anaphylaxis.*

**Atopic dermatitis (AD)**, also known as atopic eczema, is linked to a complex interaction between skin barrier dysfunction and environmental factors such as irritants, microbes, and allergens. Null mutations of the skin barrier protein filaggrin may increase the risk for transcutaneous allergen sensitization and the development of food allergy in subjects with AD. Although the Expert Panel does not mean to imply that AD results from food allergy, the role of food allergy in the pathogenesis and severity of this condition remains controversial. In some sensitized patients, particularly infants and young children, food allergens can induce urticarial lesions, itching, and eczematous flares, all of which may aggravate AD.

**Allergic contact dermatitis (ACD)** is a form of eczema caused by cell-mediated allergic reactions to chemical haptens that are additives to foods or occur naturally in foods, such as mango. Clinical features include marked pruritus, erythema, papules, vesicles, and edema. Contact urticaria can be either immunologic (IgE-mediated reactions to proteins) or non-immunologic (caused by direct histamine release).

### RESPIRATORY MANIFESTATIONS

Respiratory manifestations of IgE-mediated food allergies occur frequently during systemic allergic reactions and are an important indicator of severe anaphylaxis. However, food allergy is an uncommon cause of isolated respiratory symptoms, namely those of rhinitis and asthma.

## Appendix 2: BOP Food Allergy Questionnaire

PATIENT:		PATIENT NUMBER:	
<p>This 2-page questionnaire is a guideline for information gathering purposes:</p> <ul style="list-style-type: none"> <li>All relevant information should be written in a BEMR clinical encounter note.</li> <li>All diagnoses should be entered on the patient's problem list.</li> <li>All medication allergies should be entered into the BEMR allergy section.</li> </ul>			
<b>WHAT TRIGGERS ALLERGIES OR ANAPHYLAXIS?</b>			
<input type="checkbox"/> Eggs (V15.03)	<input type="checkbox"/> Milk (V15.02)	<input type="checkbox"/> Peanuts (V15.01)	<input type="checkbox"/> Soy (V15.05)
<input type="checkbox"/> Fish (V15.04)	<input type="checkbox"/> Nuts from trees (V15.05)	<input type="checkbox"/> Shellfish (V15.04)	<input type="checkbox"/> Wheat (V15.05)
Other (please list):			
Do you have Celiac Disease (adverse food reaction to wheat, oats, barley, and rye)? (579.0)		<input type="checkbox"/> Yes	<input type="checkbox"/> No
Do you have Oral Allergy Syndrome (mouth itching after eating foods such as raw fruit and vegetables)? (692.5)		<input type="checkbox"/> Yes	<input type="checkbox"/> No
Do you react if the food is ingested (eaten)?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
Do you react if you just touch the food?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
Do you react if the food is just close to you?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
<b>HISTORY AND ASSESSMENT DATA</b>			
Do you know what kinds of foods you cannot eat?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
If so, please list:			
Have you been diagnosed with a food allergy by a health care provider?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
How were you diagnosed with allergies before?			
At what age were you first diagnosed with allergies/anaphylaxis?			
When was the last time that you had a reaction (approximate date)?			
Do you recognize any <u>early</u> warning signs that indicate an allergic reaction?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
If so, please describe:			
How soon do symptoms occur after exposure to a particular food?			
Can the food ever be eaten without these symptoms occurring?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
Have other factors ever been involved, such as exercise, alcohol, or use of aspirin or NSAIDs?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
If so, please describe:			
Have the symptoms been present at times other than after exposure to the food?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
If so, please describe:			
Have you ever been issued an EpiPen?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
Have you been treated for a major reaction in the emergency room?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
If so, what was the approximate date and the hospital:			
Is there a history of poor outcomes from anaphylaxis treatment related to beta-blocker or ACE inhibitor therapy?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
If so, please describe:			
Is there a history of asthma (particularly, poorly controlled) or coronary artery disease?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
If so, please describe:			
<i>Appendix 2 — page 1 of 2</i>			

<b>PATIENT:</b>		<b>PATIENT NUMBER:</b>	
<b>WHAT SYMPTOMS HAVE YOU HAD BECAUSE OF FOOD ALLERGY EXPOSURE?</b>			
<input type="checkbox"/> Tightness of throat and/or chest	<input type="checkbox"/> Swelling of eyes, lips, tongue, throat, or neck	<input type="checkbox"/> Anxiety	
<input type="checkbox"/> Wheezing/difficulty breathing	<input type="checkbox"/> Flushing	<input type="checkbox"/> Irritability	
<input type="checkbox"/> Coughing or sneezing	<input type="checkbox"/> Blue or gray discoloration of lips or fingernails	<input type="checkbox"/> Dizziness	
<input type="checkbox"/> All over tingling or itching	<input type="checkbox"/> Nausea, vomiting, stomach cramping, or diarrhea	<input type="checkbox"/> Fainting	
<input type="checkbox"/> All over rash or hives	<input type="checkbox"/> Sudden mood change	<input type="checkbox"/> Voice change	
<input type="checkbox"/> Other:			
What medication(s) are used to treat your reaction symptoms? (See <a href="#">Appendix 4</a> or <a href="#">Appendix 6</a> .)			
<b>HAVE YOU BEEN DIAGNOSED WITH ANY OF THE FOLLOWING?</b>			
<input type="checkbox"/> Asthma (493.90)	<input type="checkbox"/> Latex allergy (V15.07)	<input type="checkbox"/> Medication allergies	
<input type="checkbox"/> Rhinitis (472.0)	<input type="checkbox"/> Eczema (692.9)	<input type="checkbox"/> Urticaria/angioedema (due to food – 708.0)	
If you have asthma, how often do you need a rescue medicine (albuterol)?			
Have you been hospitalized for asthma?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
<b>OTHER</b>			
Name of the regular physician prior to incarceration who can give an overall history of the patient's food allergies and past treatment plans:			
Name of any allergy specialist that the patient has seen about food allergies:			
<b>DISPOSITION</b>			
Food allergy suspected?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
Food elimination diet recommended?		<input type="checkbox"/> Yes	<input type="checkbox"/> No
Allergen-specific serum IgE testing indicated? (See <a href="#">Appendix 4</a> or <a href="#">Appendix 6</a> .)		<input type="checkbox"/> Yes	<input type="checkbox"/> No
<i>Appendix 2 — page 2 of 2</i>			

## Appendix 3: Diagnostic Criteria for Anaphylaxis

DIAGNOSIS
<p><b>The presence of <u>any one</u> of these criteria, occurring over minutes to two hours after exposure, indicates that anaphylaxis is highly likely:</b></p> <ol style="list-style-type: none"><li>1. Acute onset of an illness involving skin, mucosal tissue, or both (i.e., generalized hives, pruritus or flushing, swollen lips-tongue-uvula), <b>and</b> at least one of the following:<ul style="list-style-type: none"><li>• Respiratory compromise (i.e., dyspnea, wheeze-bronchospasm, stridor, reduced peak expiratory flow rate, hypoxemia).</li><li>• Reduced blood pressure (BP) or associated symptoms of end-organ dysfunction (i.e., hypotonia [circulatory collapse], syncope, incontinence).</li></ul><p style="text-align: center;"><b>OR</b></p></li><li>2. <b>Two or more</b> of the following that occur rapidly after exposure to a likely allergen for that patient:<ul style="list-style-type: none"><li>• Involvement of the skin-mucosal tissue (i.e., generalized hives, itch-flush, swollen lips-tongue-uvula).</li><li>• Respiratory compromise (i.e., dyspnea, wheeze, bronchospasm, stridor, reduced peak expiratory flow rate, hypoxemia).</li><li>• Reduced BP or associated symptoms of end-organ dysfunction (i.e., hypotonia, syncope, incontinence).</li><li>• Persistent GI symptoms (i.e., crampy abdominal pain, vomiting).</li></ul><p style="text-align: center;"><b>OR</b></p></li><li>3. Reduced BP after exposure to a known allergen for that patient. In adults, this is defined as:<ul style="list-style-type: none"><li>• a systolic BP of less than 90 mm Hg.</li></ul><p style="text-align: center;"><b>or</b></p><ul style="list-style-type: none"><li>• &gt; 30% decrease from that person's baseline.</li></ul></li></ol>
SIGNS AND SYMPTOMS OF ANAPHYLAXIS
<p>Usually, anaphylaxis involves more than one organ system, which helps distinguish it from other acute reactions such as asthma exacerbations, respiratory symptoms, urticaria/angioedema, or GI symptoms. In general, the signs and symptoms for anaphylaxis are the same for food-induced anaphylaxis and include:</p> <ul style="list-style-type: none"><li>• <b>Cutaneous symptoms:</b> Occur in the majority of patients, and include flushing, pruritus, urticaria, and angioedema. However, 10–20% of cases have no cutaneous manifestations.</li><li>• <b>Respiratory symptoms:</b> Occur in up to 70% of cases, and include nasal congestion and rhinorrhea, throat pruritus and laryngeal edema, stridor, choking, voice change related to laryngeal edema, wheezing, coughing, and dyspnea.</li><li>• <b>GI symptoms:</b> Occur in up to 40% of cases, and include cramping, abdominal pain, nausea, emesis, and diarrhea.</li><li>• <b>Cardiovascular symptoms:</b> Occur in up to 35% of cases, and include dizziness, tachycardia, hypotension, and hypotonia.</li><li>• <b>Other symptoms:</b> May include anxiety, mental confusion, lethargy, and seizures.</li></ul>
<i>Appendix 3 — page 1 of 2</i>

#### TIME COURSE

The time course of an anaphylactic reaction may be uniphasic, biphasic, or protracted, defined as follows:

- **A uniphasic reaction** occurs immediately after exposure and resolves—with or without treatment—within the first minutes to hours, and then does not recur during that anaphylactic episode.
- **A biphasic reaction** includes a recurrence of symptoms that develops after apparent resolution of the initial reaction. Biphasic reactions have been reported to occur in 1%–20% of anaphylaxis episodes, and typically occur about 8 hours after the first reaction, although recurrences have been reported up to 72 hours later.
- **A protracted reaction** is any anaphylaxis episode that lasts for hours or days following the initial reaction.

#### FATAL ANAPHYLAXIS


Fatalities associated with food-induced anaphylaxis are most commonly associated with peanut or tree nut ingestion. Such fatalities are associated with delayed use of epinephrine or a lack of proper dosing. The highest risk groups for fatal anaphylaxis associated with food ingestion are:

- Adolescents and young adults.
- Individuals with known food allergy and a prior history of anaphylaxis.
- Individuals with asthma, especially those with poor control (although fatal reactions may occur even in individuals with mild asthma).
- Individuals with a history of poor outcomes from anaphylaxis treatment related to the use of beta-blocker or ACE inhibitor therapy, or individuals with a history of coronary heart disease.

*Appendix 3 — page 2 of 2*




### Appendix 4: Algorithm for Patients with Suspected Food-Induced Anaphylaxis



## Appendix 5: Pharmacological Treatment of Anaphylaxis

<p><b>Note:</b> Preferred route is listed first is in <b>bold</b>. <i>Italicized</i> route can be used if preferred route is unavailable (IV = intravenous, IM= intramuscular, SQ = subcutaneous).</p>		
<b>TAKE IMMEDIATE STEPS</b>		
<p><b>Initial management should begin with the following <u>concurrent</u> steps:</b></p> <ul style="list-style-type: none"> <li>• Eliminate additional allergen exposure.</li> <li>• Administer epinephrine 0.3mg <b>IM/SQ</b> via auto-injector. Can use a 1:1000 solution <b>IM/SQ</b> at 0.01mg/kg per dose (max 0.5mg per dose).</li> <li>• Arrange for transport to the nearest emergency facility, although attempts to summon help should not delay use of epinephrine.</li> </ul>		
<b>QUICKLY FOLLOWED BY ...</b>		
<p><b>The initial actions should be quickly followed by these additional steps:</b></p> <ul style="list-style-type: none"> <li>• Place the patient in a recumbent position (if tolerated), with the lower extremities elevated.</li> <li>• Provide supplemental oxygen.</li> <li>• Administer <b>IV</b> fluid (volume resuscitation).</li> <li>• Consider administration of the following: (see <a href="#">Pharmacological Management</a> section at bottom of table) <ul style="list-style-type: none"> <li>• H<sub>1</sub> antihistamine – Diphenhydramine 50mg <b>IV/IM</b></li> <li>• Corticosteroid – Methylprednisolone sodium succinate (Solu-Medrol) up to 80mg <b>IV/IM</b></li> <li>• Bronchodilator – Albuterol nebulizer solution, as needed</li> <li>• H<sub>2</sub> antihistamine – Ranitidine 50mg <b>IV/IM</b> (if available locally)</li> </ul> </li> </ul> <p>➔ <b>Reassess and repeat epinephrine every 5–15 minutes until respiratory and cardiovascular status is stable.</b></p>		
<b>IMPORTANT NOTES</b>		
<ul style="list-style-type: none"> <li>• Prompt assessment and treatment are critical. <b>Failure to respond promptly can result in rapid decline and death within 30–60 minutes.</b></li> <li>• <b>Epinephrine 0.3mg injection IM, preferably via an auto-injector, is the first-line treatment in all cases of anaphylaxis.</b> All other drugs have a delayed onset of action. When there is suboptimal response to the initial dose of epinephrine, or if symptoms progress, repeated epinephrine dosing remains first-line therapy over adjunctive treatments.</li> <li>• Inmates will likely return from the emergency facility with orders for an H<sub>1</sub> antihistamine (e.g., diphenhydramine), an H<sub>2</sub> antihistamine (e.g., ranitidine), and a corticosteroid (e.g., prednisone) to be used for up to 3 days. These medications should be continued, or substituted with formulary equivalents, to prevent a biphasic or protracted reaction.</li> <li>• Inmates who have had severe or anaphylactic allergic reactions to food should have all allergies documented in BEMR, carry at least one epinephrine auto-injector with them at all times, and be provided with education including, at a minimum, the <a href="#">Inmate Factsheet: An Overview of Food Allergies</a> (see <a href="#">Appendix 7</a>).</li> <li>• Procedures for allowing the inmate to carry an epinephrine auto-injector should be coordinated locally. Refer to epinephrine auto-injector guidance in the BOP National Formulary (Part 1).</li> </ul>		
<b>PHARMACOLOGICAL MANAGEMENT OF FOOD ALLERGY ANAPHYLAXIS IN THE CORRECTIONAL SETTING</b>		
<i>Medication</i>	<i>Adult Dose</i>	<i>Comments</i>
Epinephrine auto-injector <i>OR</i> Epinephrine 1:1000	0.3mg <b>IM/SQ</b> 0.01mg/kg <b>IM/SQ</b> up to 0.5mg <i>per dose</i>	Give ASAP. Can repeat dose every 5–15 minutes.
Diphenhydramine*	50mg <b>IV/IM</b> <i>once</i>	
Methylprednisolone sod succ*	1mg/kg up to 80mg <b>IV/IM</b> <i>once</i>	
Albuterol*	3ml nebulizer solution <i>every 20 minutes</i>	
Ranitidine*	50mg <b>IV/IM</b>	
* <i>If clinically appropriate and available, give while awaiting arrival of EMS.</i>		

### Appendix 6: Algorithm for Patients Without History of Suspected Food-Induced Anaphylaxis



## Appendix 7: Inmate Handouts

Attached are three handouts for inmates:

- ***Inmate Fact Sheet: An Overview of Food Allergies***
- ***Inmate Fact Sheet: Lactose Intolerance***
- ***Inmate Fact Sheet: Food Avoidance and Self-Selection from the BOP National Menu***

**Note:** In addition to the above-listed handouts, inmates can be given information on **reading food labels**, which is available at the Food Allergy and Anaphylaxis Network website at <http://www.foodallergy.org/page/patient-handouts>. Printable handouts are available for identifying foods containing milk, egg, peanuts, tree nuts, wheat, soy, and shellfish.

## Inmate Factsheet: An Overview of Food Allergies

### What is a food allergy?

A food allergy to a certain food is when a person's immune system responds as if the food is a harmful substance. The response may be mild or, in rare cases, it may be a life-threatening reaction called "anaphylaxis." An allergic reaction to food is most likely to take place within 5 minutes to 1 hour after eating or touching the food.

### What are the most common foods that adults are allergic to?

Peanuts, tree nuts (e.g., almonds, cashews), fish, and shellfish (e.g., oysters, crab)

### What are the symptoms of a food allergy?

#### ***Mild symptoms include:***

- Hives
- Itching in your mouth
- Swelling of the lips and tongue
- Vomiting, diarrhea, or abdominal cramps and pain
- Red, swollen skin or worsening of eczema
- Itchy, watery, or swollen eyes
- Runny nose and/or sneezing

#### ***Severe, life-threatening symptoms include:***

- Swelling of the throat or trouble breathing
- Drop in blood pressure
- Wheezing or difficulty breathing
- Feeling dizzy or passing out

### Can food allergy reactions be prevented?

Yes! Reactions can be prevented by completely avoiding any foods you are allergic to. Talk to your Primary Care Provider or Registered Dietitian about:

- How to read food labels to see if there are ingredients you are allergic to
- How to avoid the foods you are allergic to when they are offered by the Food Service or available in the commissary
- When and how to get help for an allergic reaction, whether it is mild or severe
- Whether you might need to carry an epinephrine auto-injector at all times

### What should you do for a severe allergic reaction?

The best treatment for a severe allergic reaction is a medicine called epinephrine. If you feel you are having a severe allergic reaction to food, **immediately** let the nearest staff member know you need medical assistance. If you carry an epinephrine auto-injector, use it immediately.

**Inmate Factsheet: Lactose Intolerance**

**What is lactose intolerance?**

Lactose intolerance means that a person’s body cannot produce enough of the enzyme that helps digest the lactose found in dairy products. Lactose intolerance can affect anyone, but is most common in Native Americans, Asians, African Americans, and older individuals.

**Which foods contain lactose?**

Dairy products such as milk, yogurt, ice cream, cheese, cream, and butter all contain lactose. If you are highly sensitive to lactose, you will also need to be aware of other foods that are made with dairy products—such as some baked goods, for example.

**What are the symptoms of lactose intolerance?**

Any or all of the following symptoms can occur after eating dairy products:

- Cramps or stomach pain
- Gas
- Bloating feeling
- Diarrhea
- Vomiting

**How is lactose intolerance treated?**

There are two ways to treat lactose intolerance. (1) One approach is to eat only small, tolerable amounts of foods that contain lactose, and not on an empty stomach, i.e., eat them together with other foods. (2) The other is to use an over-the-counter enzyme supplement that you can purchase from the commissary (Lactaid™, Dairy Ease™, etc.). If you decide to use this supplement, you must take it right before each meal or snack that contains dairy products.

**Which common foods on the BOP menu or in the commissary contain lactose?**

<i>High in Lactose</i>	<i>Medium</i>	<i>Low in Lactose</i>
<ul style="list-style-type: none"> <li>• Milk – whole, 2%, and skim</li> <li>• Ice cream</li> </ul>	<ul style="list-style-type: none"> <li>• Cottage cheese</li> <li>• Sherbet</li> </ul>	<ul style="list-style-type: none"> <li>• Butter</li> <li>• Processed cheese</li> <li>• Mozzarella cheese</li> <li>• Cheddar cheese</li> </ul>

## **Inmate Factsheet: Food Avoidance and Self-Selection from the BOP National Menu**

### **Prevention**

The best way to prevent an allergic reaction to food is to know which foods cause the signs and symptoms—and avoid them!

### **Self-Selection**

In all cases of food allergy, the first option is to simply avoid the item through self-selection from the BOP National Menu. If an item, even an entrée, is on the National Menu only once or twice in any given week, you may decline the item or self-select the no-flesh option or the heart-healthy option, if appropriate. Your health care provider and Food Service Administrator can help to educate you to choose the items you need to meet your prescribed diet from the items available on the menu.

Only the individuals with allergies to wheat, egg, milk, or fruits will have specific menu substitutions made for them, since these items are offered so often on the menu that avoiding them creates a nutritional risk.

### **Satellite Meal Service**

In satellite service areas, it is unlikely that you will have the opportunity to select specific options from the National Menu. In these situations, let staff know of your allergy. The food service administrator will develop procedures so that your diet order is processed properly, including items that you would normally self-select in a cafeteria setting.

### **Commissary Food Items**

Always read food labels to make sure that they do not contain an ingredient you are allergic to. Even if you think you know what is in a food—check the label! Ingredients sometimes change. Food labels are required to clearly list whether they contain any common food allergens, including: milk, eggs, peanuts, tree nuts, fish, shellfish, soy, and wheat.